

ADAMS' NEW  
**ARITHMETIC,**  
SUITED TO HALIFAX CURRENCY;

IN WHICH THE  
PRINCIPLES OF OPERATING BY NUMBERS

ARE  
**ANALYTICALLY EXPLAINED,**  
AND  
**SYNTHETICALLY APPLIED:**

THUS  
COMBINING THE ADVANTAGES  
TO BE DERIVED BOTH FROM  
**THE INDUCTIVE AND SYNTHETIC MODE**  
**OF INSTRUCTING:**

THE WHOLE  
MADE FAMILIAR BY A GREAT VARIETY OF USEFUL AND INTERESTING  
EXAMPLES, CALCULATED AT ONCE TO ENGAGE THE PUPIL IN  
THE STUDY, AND TO GIVE HIM A FULL KNOWLEDGE  
OF FIGURES IN THEIR APPLICATION TO  
ALL THE PRACTICAL PURPOSES  
OF LIFE.

DESIGNED FOR THE USE OF  
SCHOOLS AND ACADEMIES IN THE  
**BRITISH PROVINCES.**

BY DANIEL ADAMS, M. D.

STANSTEAD, L. C.  
PUBLISHED BY WALTON & GAYLORD.

1833.



R  
QA

101

A3

1833

T  
In th  
view  
of w  
pupil  
certa  
gene  
TH  
is a f  
The  
ment  
intro  
Swit  
by M  
Th  
ledge  
it. I  
ful.  
occup  
forbo  
tinuin  
ronap  
shoul  
scienc  
to ma

In  
conta  
follov  
pleme  
till it  
vious  
the r  
Th  
ginni  
those  
and v  
mann  
figur  
plical  
At  
to co  
rules  
Thes  
appli  
to int  
appli

## PREFACE.

THERE are two methods of teaching: the *synthetic* and the *analytic*. In the synthetic method, the pupil is first presented with a *general* view of the science he is studying, and afterwards with the particulars of which it consists. The analytic method *reverses* this order: the pupil is first presented with the *particulars*, from which he is led, by certain natural and easy gradations, to those views which are more general and comprehensive.

The Scholar's Arithmetic published in 1801, is synthetic. If that is a *fault* of the work, it is a fault of the *times* in which it appeared. The analytic or inductive method of teaching, as now applied to elementary instruction, is among the improvements of later years. Its introduction is ascribed to PESTALOZZI, a distinguished teacher in Switzerland. It has been applied to arithmetic, with great ingenuity, by Mr. COLBURN, in our own country.

The analytic is unquestionably the best method of *acquiring* knowledge; the synthetic is the best method of *recapitulating* or *reviewing* it. In a treatise designed for school education, *both* methods are useful. Such is the plan of the present undertaking which the author, occupied as he is with other objects and pursuits, would willingly have forborne, but that, the demand for the Scholar's Arithmetic still continuing, an obligation, incurred by long-continued and extended patronage, did not allow him to decline the labor of a revision, which should adapt it to the present more enlightened views of teaching this science in our schools. In doing this, however, it has been necessary to make it a new work.

In the execution of this design, an *analysis* of each rule is first given, containing a familiar explanation of its various principles; after which follows a *synthesis* of these principles, with questions in form of a supplement. Nothing is taught *dogmatically*; no technical term is used till it has first been defined, nor any principle inculcated without a previous developement of its truth; and the pupil is made to understand the *reason* of each process as he proceeds.

The examples under each rule are mostly of a practical nature, beginning with those that are very easy, and gradually advancing to those more difficult, till one is introduced containing larger numbers, and which is not easily solved in the mind; then in a plain, familiar manner, the pupil is shown how the solution may be facilitated by *figures*. In this way he is made to see at once their *use* and their *application*.

At the close of the fundamental rules, it has been thought advisable to collect into one clear view the distinguishing properties of those rules, and to give a number of examples involving one or more of them. These exercises will prepare the pupil more readily to understand the application of these to the succeeding rules; and, besides, will serve to interest him in the science, since he will find himself able, by the application of a very few principles, to solve many curious questions.



The arrangement of the subjects is that, which to the author has appeared most natural. *Fractions*, have received all that consideration which their importance demands. The principles of a rule called *Practice* are exhibited, but its detail of cases omitted, as unnecessary since the adoption and general use of federal money. The *Rule of Three, or Proportion*, is retained and the solution of questions involving the principles of proportion, by *analysis*, is distinctly shown.

The articles *Alligation, Arithmetical and Geometrical Progression, Annuities and Permutation*, were prepared by Mr. ISA YOUNG, a member of Dartmouth College, from whose knowledge of the subject, and experience in teaching, I have derived important aid in other parts of the work.

The numerical paragraphs are chiefly for the purpose of *reference*; these references the pupil should not be allowed to neglect. His attention also ought to be particularly directed, by his instructor, to the illustration of each particular principle, from which general rules are deduced; for this purpose, recitations by classes ought to be instituted in every school where arithmetic is taught.

The supplements to the rules, and the geometrical demonstrations of the extraction of the square and cube roots, are the only traits of the *old work* preserved in the *new*.

DANIEL ADAMS.

Mount Vernon, (N. H.) Sept. 29, 1827.

The  
metie  
States  
great  
of Ac  
metie  
versa  
view  
estab  
ry ed  
this t  
This  
The  
law i  
quain  
arriv  
are t  
man  
are c  
prud  
meth  
mig  
amp  
the  
ject  
the  
I  
met  
din  
wh  
I  
ada  
ces  
lot  
red  
am  
Ac

## PUBLISHERS' PREFACE.

THE author of the following practical treatise upon Arithmetic, has made himself favorably known in the United States and to a considerable extent in the Canadas, for a great number of years, by his works, designed for the use of Academies and primary schools. The "Scholars' Arithmetic," published in the year 1801, continued in almost universal use, until within a very short time past. But juster views beginning to prevail, and sounder principles becoming established in the public mind, upon the subject of elementary education, a revision of the work seemed necessary. At this time, "Adams' New Arithmetic," was published.— This seems evidently to have been prepared with much care. The author has recognized in it throughout, this important law in relation to the mind, that it must first be made acquainted with particular facts, or there will be no ability to arrive at correct general conclusions. Particular examples are therefore given upon each subject, and from them, in a manner obvious to the young mind, all the general rules are deduced. In other words, the author has carefully and prudently pursued, in his book, what is called the *analytic* method. The care used in defining necessary terms, which might not be quite clear, the practical character of the examples given under each rule, the methodical disposition of the different parts of each subject, and of the different subjects, the general perspicuity, simplicity and accuracy of the work, render it invaluable to the pupil.

It is due the author to observe, that "Adams' New Arithmetic," for its adaptation to the capacities of young and ordinary minds, is justly considered the best practical treatise which has been offered to the public.

In the present edition, the main purpose in view was to adapt Adam's work to the currency of the British Provinces. No separate article, as in the original, has been allotted to Federal Money; for this the pupil has been referred to Decimal Fractions, in which also almost all the examples will be found in the money of the United States.— Additional examples in the compound rules have been given.

en, and the old ones retained, under the title of Halifax currency; and generally, throughout the book, where denominations of money occur, Halifax currency has been substituted for Federal Money.

The rules and examples in Reduction of Currencies have been essentially changed; and in Reduction, after the Table of English Money, which is called the Table of Halifax Currency, a list of the Gold and Silver Coins current in the Province, has been inserted. This may be depended upon as entirely accurate. The tables of French, and Dry, Long, Square, and Solid Measure, have been given—and what are the weights and measures established by law in this Province is also stated.

The most novel feature in the book will be found in the rule of Interest. Certainly an innovation, but it is believed, an improvement, has been made. The pounds in any given sum upon which interest is to be cast, are left to stand as the units, and the shillings and pence are reduced to decimal parts of a pound. The interest is then obtained the same as in Federal Money, and the decimal parts in the result reduced to shillings and pence. It is considered, that this method is more simple, and concise, and will be found in practice to be more convenient than any other. But setting aside considerations of temporary convenience, if this change and attempted amelioration, shall assist in some very slight degree in turning men's minds toward the *Decimal Ratio*, and inducing them to look forward to a period when all the denominations of money, weights and measures, throughout the world, shall be expressed in DECIMALS, it cannot be affirmed that no benefit has been obtained.

The importance of the *principal and essential* alteration in the book, viz. the adaptation of it to the currency of the country, will not fail to be observed by every one. It is indeed singular, that hitherto, no *Canadian Arithmetic* in the English language, has been published. Mercantile, agricultural, and generally the business men of the country, will be aware of a benefit to be realized, and it is considered that something also bearing a relation to political advantage, may be in the result.

*Stanstead, L. C. Aug. 1, 1833.*

Ad  
All  
Ari  
Co

Con  
Co  
Cul  
Div  
Duc

Equ  
Evo  
Fed  
Fra

Fel  
Hal  
Ins  
Int

Inv  
Mu  
Nu  
Pro

# INDEX.

	Page.
Addition, . . . . .	14
Alligation, . . . . .	192
Arithmetical Progression, . . . . .	218
Compound Numbers, . . . . .	58
Addition of, . . . . .	80
Subtraction of, . . . . .	85
Multiplication and Division of, . . . . .	89
Commission, . . . . .	152
Coins, Table of, . . . . .	59
Cube Root, Extraction of, . . . . .	211
Division, . . . . .	39
Duodecimals, . . . . .	197
Multiplication of, . . . . .	198
Equation of Payments, . . . . .	171
Evolution, . . . . .	202
Federal Money, . . . . .	64
Fractions, . . . . .	98
Proper and Improper, . . . . .	100
To change an improper fraction to a whole or mixed number, . . . . .	100
To reduce a whole or mixed number to an improper fraction, . . . . .	101
To reduce a fraction to its lowest terms, . . . . .	102
To divide a fraction by a whole number, . . . . .	105
To multiply a fraction by a whole number, . . . . .	107
whole number by a fraction, . . . . .	109
one fraction by another, . . . . .	110
To divide a whole number by a fraction, . . . . .	112
one fraction by another, . . . . .	114
Addition and subtraction of fractions, . . . . .	116
Reduction of . . . . . do. . . . .	121
Decimal Fractions, . . . . .	129
Addition and subtraction of, . . . . .	132
Multiplication of, . . . . .	135
Division of, . . . . .	137
Reduction of, . . . . .	143
Reduction of vulgar fractions to decimals, . . . . .	140
Fellowship, . . . . .	188
Halifax Currency, . . . . .	59
Insurance, . . . . .	152
Interest, . . . . .	150
Compound, . . . . .	164
Involution, . . . . .	201
Multiplication, Simple, . . . . .	28
Numeration, . . . . .	9
Proportion, or the Rule of Three, . . . . .	174
Compound, . . . . .	182

Permutation, . . . . .	232
Reduction, . . . . .	58
----- Tables of Money, Weights, Measures, &c., . . . . .	59—78
Reduction of Currencies, . . . . .	149
Ratio, or the Relation of Numbers, . . . . .	172
Subtraction, Simple, . . . . .	21
Square Root, Extraction of, . . . . .	203

### MISCELLANEOUS EXMAPLES.

Barter, ex. 21—32. . . . .	Position, ex. 89—108.
To find the area of a Square or Parallelogram, ex. 148—154.	
----- a Triangle, ex. 155—159.	
Having the Diameter of a Circle, to find the Circumference; or hav-	
ing the Circumference, to find the Diameter, ex. 171—175.	
To find the Area of a Circle, ex. 176—179.	
----- a Globe, ex. 180, 181.	
To find the Solid Contents of a Globe, ex. 182, 184.	
----- Cylinder, ex. 185—187.	
----- Pyramid, or Cone, ex. 188, 189.	
----- any Irregular Body, ex. 202.	
Gauging, ex. 190, 191. . . . .	Mechanical Powers, ex. 192—201.
Forms of Notes, Receipts, Orders and Bills of Parcels, page, . . . . .	258
Book-Keeping, . . . . .	259



# ARITHMETIC.

## NUMERATION.

¶ 1. A SINGLE or individual thing is called a *unit*, *unity*, or *one*; one and one more are called *two*; two and one more are called *three*; three and one more are called *four*; four and one more are called *five*; five and one more are called *six*; six and one more are called *seven*; seven and one more are called *eight*; eight and one more are called *nine*; nine and one more are called *ten*, &c.

These terms, which are expressions for quantities, are called *numbers*. There are two methods of expressing numbers shorter than writing them out in words; one called the *Roman* method by letters,\* and the other the *Arabic* method by figures. The latter is that in general use.

In the Arabic method, the nine first numbers have each an appropriate character to represent them. Thus,

\*In the Roman method by letters, I represents *one*; V, *five*; X, *ten*; L, *fifty*; C, *one hundred*; D, *five hundred*; and M, *one thousand*.

As often as any letter is repeated, so many times is its value repeated, unless it be a letter representing a *less* number, placed before one representing a *greater*; then the less number is taken from the greater, thus IV represents *four*, IX *nine*, &c., as will be seen in the following

TABLE.			
One	I.	Ninety.	LXXXX, or XC.
Two	II.	One hundred	C.
Three	III.	Two hundred	CC.
Four	IIII, or IV.	Three hundred	CCC.
Five	V.	Four hundred	CCCC.
Six	VI.	Five hundred	D, or I <sub>0</sub> .*
Seven	VII.	Six hundred	DC.
Eight	VIII.	Seven hundred	DCC.
Nine	VIII, or IX.	Eight hundred	DCCC.
Ten	X.	Nine hundred	DCCCC.
Twenty	XX.	One thousand	M, or C <sub>10</sub> .†
Thirty	XXX.	Five thousand	I <sub>00</sub> . or V <sub>1</sub> ‡
Forty	XXXX, or XL.	Ten thousand	CCI <sub>100</sub> , or X̄.
Fifty	L.	Fifty thousand	I <sub>000</sub> .
Sixty	LX.	Hundred thousand	CCCI <sub>000</sub> , or C̄.
Seventy	LXX.	One million	M̄.
Eighty	LXXX.	Two millions	MM̄.

\* I<sub>0</sub> is used instead of D to represent five hundred, and for every additional 0 annexed at the right hand, the number is increased *ten times*.

† C<sub>10</sub> is used to represent one thousand, and for every C and 0 put at each end, the number is increased *ten times*.

‡ A line drawn over any number increases its value *one thousand times*.

A unit, unity, or one, is represented by this character

<i>Two</i>	1.
<i>Three</i>	2.
<i>Four</i>	3.
<i>Five</i>	4.
<i>Six</i>	5.
<i>Seven</i>	6.
<i>Eight</i>	7.
<i>Nine</i>	8.
	9

*Ten* has no appropriate character to represent it ; but is considered as forming a unit of a second or higher order, consisting of *tens*, represented by the same character (1) as a unit of the first or lower order, but is written in the *second place* from the right hand, that is, on the left hand side of units ; and as, in this case, there are no units to be written with it, we write, in the place of units, a cipher, 0, which of itself signifies nothing ; thus, *Ten*,

One ten and one unit are called	<i>Eleven</i>	10.
One ten and two units are called	<i>Twelve</i>	11.
One ten and three units are called	<i>Thirteen</i>	12.
One ten and four units are called	<i>Fourteen</i>	13.
One ten and five units are called	<i>Fifteen</i>	14.
One ten and six units are called	<i>Sixteen</i>	15.
One ten and seven units are called	<i>Seventeen</i>	16.
One ten and eight units are called	<i>Eighteen</i>	17.
One ten and nine units are called	<i>Nineteen</i>	18.
Two tens are called	<i>Twenty</i>	19.
Three tens are called	<i>Thirty</i>	20.
Four tens are called	<i>Forty</i>	30.
Five tens are called	<i>Fifty</i>	40.
Six tens are called	<i>Sixty</i>	50.
Seven tens are called	<i>Seventy</i>	60.
Eight tens are called	<i>Eighty</i>	70.
Nine tens are called	<i>Ninety</i>	80.

*Ten* tens are called a *hundred*, which forms a unit of a still higher order, consisting of *hundreds*, represented by the same character (1) as a unit of each of the foregoing orders, but is written one place further toward the left hand, that is on the left hand side of *tens* ; thus, *one hundred* 100.

One hundred, one ten, and one unit, are called *One hundred and eleven* 111.

In this viz. un occupy place from number exhibit order, 3 ; but, ing from thus, 30 are six, but, tha of tens, The low charact

We r ing dov then th then th tens, th three h three h

ue acco right h

Take figure t the first ond pla creased signifie ten time ately p fundam wards t

Ten order.

ter 1.  
2.  
3.  
4.  
5.  
6.  
7.  
8.  
9.  
it is  
ther  
ame  
der,  
ght  
and  
ten  
, 0,  
en 10.  
ve 11.  
teen 12.  
teen 13.  
en 14.  
n 15.  
n 16.  
teen 17.  
teen 18.  
teen 19.  
y 20.  
y 30.  
40.  
50.  
60.  
y 70.  
80.  
90.  
f a  
at-  
of  
h-  
nd  
100.  
111.

¶ 2. There are three hundred sixty-five days in a year. In this number are contained all the orders now described, viz. units, tens, and hundreds. Let it be recollected, *units* occupy the *first place* on the right hand; *tens*, the *second place* from the right hand; *hundreds*, the *third place*. This number may now be *decomposed*, that is, *separated into parts*, exhibiting each order by itself, as follows:—The highest order, or hundreds, are *three*, represented by this character, 3; but, that it may be made to occupy the third place, counting from the right hand, it must be followed by two ciphers, thus, 300. (three hundred.) The next lower order, or *tens*, are six, (six tens are sixty,) represented by this character, 6; but, that it may occupy the second place, which is the place of tens, it must be followed by one cipher, thus 60, (sixty.) The lowest order, or *units*, are five, represented by a single character, thus, 5, (five.)

We may now combine all these parts together, first writing down the five units for the right hand figure, thus, 5; then the six tens (60) on the left hand of the units, thus 65; then the three hundreds (300) on the left hand of the six tens, thus, 365, which number, so written, may be read three hundred, six tens, and five units; or, as is more usual, three hundred and sixty-five.

¶ 3. Hence it appears, that figures have a different value according to the *place* they occupy, counting from the right hand towards the left.

Hund.  
Tens.  
Units.

Take for example the number 3 3 3, made by the same figure three times repeated. The 3 on the right hand, or in the *first place*, signifies 3 units; the same figure, in the *second place*, signifies 3 *tens*, or thirty; its value is now increased ten times. Again, the same figure in the *third place*, signifies neither 3 *units*, nor 3 *tens*, but 3 *hundreds*, which is *ten times* the value of the same figure in the place immediately preceding, that is, in the place of *tens*; and this is a fundamental law in notation, that a *removal of one place towards the left increases the value of a figure TEN TIMES*.

Ten hundred make a *thousand*, or a unit of the *fourth* order. Then follow tens and hundreds of thousands, in the







5. Twenty-seven hundred and nineteen.
6. Forty-nine hundred and sixty.
7. Ninety-two thousand and forty-five.
8. One hundred thousand.
9. Two millions, eighty thousands, and seven hundreds.
10. One hundred millions, one hundred thousand, one hundred and one.
11. Fifty-two millions, six thousand, and twenty.
12. Six billions, seven millions, eight thousand, and nine hundred.
13. Ninety-four billions, eighteen thousand, one hundred and seventeen.
14. One hundred thirty-two billions, two hundred millions, and nine.
15. Five trillions, sixty billions, twelve millions, and ten thousand.
16. Seven hundred trillions, eighty-six billions, and seven millions.

**ADDITON**

OF SIMPLE NUMBERS.

14. 1. James had five peaches, his mother gave him 3 peaches more ; how many peaches had he then ?
2. John bought one book for 9 pence, and another for 6 pence ; how many pence did he give for both ?
3. Peter bought a waggon for 10 shillings, and sold it so as to gain 4 shillings ; how many shillings did he get for it ?
4. Frank gave 15 walnuts to one boy, 8 to another, and had 7 left ; how many walnuts had he at first ?
5. A man bought a carriage for 54 pounds ; he expended 8 pounds in repairs, and then sold it so as to gain 5 pounds ; how many pounds did he get for the carriage ?
6. A man bought 3 yoke of oxen ; for the first he gave 16 pounds, for the second he gave 18 pounds, and for the third he gave 20 pounds ; how many pounds did he give for the three ?
7. Samuel bought an orange for four pence, and some walnuts for three pence ; then he bought a knife for 1 shilling, and a book for 4 shillings ; how many shillings did he spend, and how many pence ?

8. A man had 3 calves worth 10 shillings each, 4 calves worth 15 shillings each, and 7 calves worth 2 pounds each; how many calves had he?

9. A man sold a cow for 4 pounds, some corn for 5 pounds, wheat for 7 pounds, and butter for 2 pounds; how many pounds must he receive?

The putting together two or more numbers, (as in the foregoing examples,) so as to make one *whole number*, is called *Addition*, and the whole number is called the *sum*, or *amount*.

10. One man owes me 5 pounds, another 6 pounds, another 14 pounds, and another 3 pounds; what is the amount due to me?

11. What is the amount of 4, 3, 7, 2, 8, and 9 pounds?

12. In a certain school 9 study grammar, 15 study arithmetic, 20 attend to writing, and 12 study geography; what is the whole number of scholars?

**SIGNS.** A cross,  $+$ , one line horizontal and the other perpendicular, is the sign of addition. It shows that numbers, with this sign between them, are to be added together. It is sometimes read *plus*, which is a Latin word, signifying *more*.

Two parallel, horizontal lines,  $=$ , are the sign of equality. It signifies that the number *before* it is equal to the number *after* it. Thus,  $5+3=8$  is read 5 and 3 are 8; or, 5 plus (that is, more) 3 is equal to 8.

In this manner let the pupil be instructed to commit the following

ADDITION TABLE.

$2+0=2$	$3+0=3$	$4+0=4$	$5+0=5$
$2+1=3$	$3+1=4$	$4+1=5$	$5+1=6$
$2+2=4$	$3+2=5$	$4+2=6$	$5+2=7$
$2+3=5$	$3+3=6$	$4+3=7$	$5+3=8$
$2+4=6$	$3+4=7$	$4+4=8$	$5+4=9$
$2+5=7$	$3+5=8$	$4+5=9$	$5+5=10$
$2+6=8$	$3+6=9$	$4+6=10$	$5+6=11$
$2+7=9$	$3+7=10$	$4+7=11$	$5+7=12$
$2+8=10$	$3+8=11$	$4+8=12$	$5+8=13$
$2+9=11$	$3+9=12$	$4+9=13$	$5+9=14$

6 + 0 = 6	7 + 0 = 7	8 + 0 = 8	9 + 0 = 9
6 + 1 = 7	7 + 1 = 8	8 + 1 = 9	9 + 1 = 10
6 + 2 = 8	7 + 2 = 9	8 + 2 = 10	9 + 2 = 11
6 + 3 = 9	7 + 3 = 10	8 + 3 = 11	9 + 3 = 12
6 + 4 = 10	7 + 4 = 11	8 + 4 = 12	9 + 4 = 13
6 + 5 = 11	7 + 5 = 12	8 + 5 = 13	9 + 5 = 14
6 + 6 = 12	7 + 6 = 13	8 + 6 = 14	9 + 6 = 15
6 + 7 = 13	7 + 7 = 14	8 + 7 = 15	9 + 7 = 16
6 + 8 = 14	7 + 8 = 15	8 + 8 = 16	9 + 8 = 17
6 + 9 = 15	7 + 9 = 16	8 + 9 = 17	9 + 9 = 18

---


$$5 + 9 = \text{how many ?}$$

$$8 + 7 = \text{how many ?}$$

$$4 + 3 + 2 = \text{how many ?}$$

$$6 + 4 + 5 = \text{how many ?}$$

$$2 + 0 + 4 + 6 = \text{how many ?}$$

$$7 + 1 + 0 + 8 = \text{how many ?}$$

$$3 + 0 + 9 + 5 = \text{how many ?}$$

$$9 + 2 + 6 + 4 + 5 = \text{how many ?}$$

$$1 + 3 + 5 + 7 + 8 = \text{how many ?}$$

$$1 + 2 + 3 + 4 + 5 + 6 = \text{how many ?}$$

$$8 + 9 + 0 + 2 + 4 + 5 = \text{how many ?}$$

$$6 + 2 + 5 + 0 + 8 + 3 = \text{how many ?}$$

---

¶ 5. When the numbers to be added are *small*, the addition is readily performed in the *mind*; but it will frequently be more convenient, and even necessary, to write the numbers down before adding them.

13. Harry had 43 books in his little library, his father gave him 25 volumes more; how many volumes had he then?

One of these numbers contains 4 tens and 3 units. The other number contains 2 tens and 5 units. To unite these two numbers together into one, write them down one under the other, placing the *units* of one number directly under *units* of the other, and the *tens* of one number directly under *tens* of the other, thus :

43 volumes,      Having written the numbers in this  
25 volumes.      manner, draw a line underneath.

---

Ans.

It

6 ten

14

horse

the v

W

units

C

Carr

Hors

Sadd

Ans

A

amp

15

past

the

10

ano

in t

1

so a

625

I

has

sin

of

ure

0 = 9  
1 = 10  
2 = 11  
3 = 12  
4 = 13  
5 = 14  
6 = 15  
7 = 16  
8 = 17  
9 = 18

43 volumes, We then begin at the right hand, and  
25 volumes, add the 5 units of the lower number to  
— the 3 units of the upper number, making  
8 8 units, which we set down in unit's  
place.

43 volumes, We then proceed to the next column,  
25 volumes, and add the 2 tens of the lower number  
— to the 4 tens of the upper number, mak-  
ing 6 tens, or 60, which we set down in  
ten's place, and the work is done.

Ans. 68 volumes.

It now appears that Harry's whole number of volumes is 6 tens and 8 units, or 68 volumes; that is,  $43 + 25 = 68$ .

14. A gentleman bought a carriage for 214 pounds, a horse for 30 pounds, and a saddle for 4 pounds; what was the whole amount?

Write the numbers as before directed, with units under units, tens under tens, &c.

OPERATION.

Carriage, 214 pounds, Add as before. The units will  
Horse, 30 pounds, be 8, the tens 4, and the hundreds  
Saddle, 4 pounds. 2, that is,  $214 + 30 + 4 = 248$ .

Answer, 248 pounds.

After the same manner are performed the following examples:

15. A man had 15 sheep in one pasture, 20 in another pasture, and 143 in another; how many sheep had he in the three pastures?  $15 + 20 + 143 =$  how many?

16. A man has three farms, one containing 500 acres, another 213 acres, and another 76 acres; how many acres in the three farms?  $500 + 213 + 76 =$  how many?

17. Bought a farm for 625 pounds, and afterward sold it so as to gain 150 pounds; what did I sell the farm for?  $625 + 150 =$  how many?

Hitherto the amount of any one column, when added up, has not exceeded 9; consequently has been expressed by a single figure. But it will frequently happen that the amount of a single column will exceed 9, requiring two or more figures to express it.

18. There are three bags of money. The first contains  
B 2

976 pounds, the second 653 pounds, the third 524 pounds ; what is the amount contained in all the bags 1

**OPERATION.**

*First bag,* 876

*Second bag,* 653

*Third bag,* 524

*Amount.* 2053

Writing down the numbers as already directed, we begin with the right hand, or unit column, and find the amount to be 13, that is, 3 units and 1 ten. Setting down the 3 units, or right hand figure, in unit's place, directly under the column, we reserve

the 1 ten, or left hand figure, to be added with the other tens, in the next column, saying, 1, which we reserved, to 2 makes 3, and 5 are 8, and 7 are 15, which is 5 units of its own order, and 1 unit of the next higher order, that is, 5 tens and 1 hundred. Setting down the 5 tens, or right hand figure, directly under the column of tens, we reserve the left hand figure, or 1 hundred, to be added in the column of hundreds, saying 1 to 5 is 6, and 6 are 12, and 8 are 20, which being the last column, we set down the whole number, writing the 0, or right hand figure, directly under the column, and carrying forward the 2, or left hand figure, to the next place, or place of thousands. Wherefore, we find the whole amount of money contained in the three bags to be 2053 pounds—the answer.

**PROOF.** We may reverse the order, and, beginning at the top, add the figures downward. If the two results are alike, the work is supposed to be right.

From the examples and illustrations now given, we derive the following

**RULE.**

I. Write the numbers to be added, one under another, placing units under units, tens under tens, &c. and draw a line underneath.

II. Begin at the right hand or unit column, and add together all the figures contained in that column : if the amount does not exceed 9, write it under the column ; but if the amount exceed 9, so that it shall require two or more figures to express it, write down the unit figure only under the column ; the figure or figures to the left hand of units, being tens, are so many units of the next higher order, which, being reserved, must be carried forward, and added to the first figure in the next column.

III. Add each succeeding column in the same manner, and set down the whole amount at the last column.

19. A  
1817 pou  
pounds, a  
of hay pu

20. A  
pounds, I

21. A  
850 bush  
els of bar  
amount 1

22. St  
sterling ;  
sion-Hou  
pounds ;  
Monume  
sums !

23. If  
ing coun  
Dorches  
Stanstea  
32,871 ;

what w  
ties at t

24. F  
from Ea  
more ;  
the bui  
kings fr  
363 ye  
christia  
to the

286  
310



EXAMPLES FOR PRACTICE.

19. A man bought four loads of hay; one load weighed 1817 pounds, another weighed 1950 pounds, another 2156 pounds, and another 2210 pounds; what was the amount of hay purchased?

20. A person owes A 100 pounds, B 522 pounds, C 785 pounds, D 92 pounds; what is the amount of his debts?

21. A farmer raised in one year 1200 bushels of wheat, 850 bushels of Indian corn, 1000 bushels of oats, 1086 bushels of barley, and 74 bushels of peas; what was the whole amount? *Ans.* 4210.

22. St. Paul's Cathedral, in London, cost 800,000 pounds sterling; the Royal Exchange 80,000 pounds; the Mansion-House 40,000 pounds; Black Friars Bridge 152,840 pounds; Westminster Bridge 389,000 pounds, and the Monument 13,000 pounds; what is the amount of these sums? *Ans.* 1,474,840 pounds.

23. If at the census in 1831, the population of the following counties was as follows:—Lower Canada: Gaspé, 4,171; Dorchester, 11,946; Nicolet, 12,504; Sherbrooke, 6,814; Stanstead, 8,272; Upper Canada: Gore, 23,552; Home, 32,871; Niagara, 21,974; London, 26,180; Ottawa, 4,456; what was the whole number of inhabitants in these counties at that time? *Ans.* 152,740.

24. From the creation to the departure of the Israelites from Egypt was 2513 years; to the siege of Troy, 307 years more; to the building of Solomon's Temple, 180 years; to the building of Rome, 251 years; to the expulsion of the kings from Rome, 244 years; to the destruction of Carthage, 363 years; to the death of Julius Cesar, 102 years; to the christian era, 44 years; required the time from the creation to the christian era. *Ans.* 4004 years.

25.

2863705421061  
3107429315638  
6253034792  
247135  
8673

---

26.

4367583021463  
1752349713620  
6081275306217  
5652174630128  
8703263472013

---

27.

5364207631023  
 2812345672948  
 6057042087094  
 3162835906718  
 7604286537892

---

28.

902375468213  
 283496732670  
 930634216732  
 236547802436  
 8050607080900

---

29. What is the amount of 46723, 6742, and 986 pounds?

30. A man has three orchards; in the first there are 140 trees that bear apples, and 64 trees that bear peaches; in the second, 234 trees bear apples, and 73 bear cherries; in the third, 47 trees bear plums, 36 bear pears, and 25 bear cherries; how many trees in all the orchards?

## SUPPLEMENT

### TO NUMERATION AND ADDITION.

#### QUESTIONS.

1. What is a single or individual thing called? 2. What is notation? 3. What are the methods of notation now in use? 4. How many are the Arabic characters or figures? 5. What is numeration? 6. What is a fundamental law in notation? 7. What is addition? 8. What is the rule for addition? 9. What is the result, or number sought, called? 10. What is the sign of addition? 11. — of equality? 12. How is addition proved?

#### EXERCISES.

1. Washington was born in the year of our Lord 1732; he was 67 years old when he died; in what year of our Lord did he die?

2. The invasion of Greece by Xerxes took place 481 years before Christ; how long ago is that this current year 1833?

3. There are two numbers, the less number is 8671, the difference between the numbers is 597; what is the greater number.

4. A man borrowed a sum of money, and paid in part 684 pounds; the sum left unpaid was 876 pounds, what was the sum borrowed?

28.

5 4 6 8 2 1 3  
3 7 3 2 6 7 0  
2 1 6 7 3 2  
8 0 2 4 3 6  
7 0 8 0 9 0

986 pounds;  
there are 140  
peaches; in  
cherries; in  
and 25 bear

ON.

What is nota-  
use? 4. How  
numeration?  
is addition?  
lt, or number  
11. — of

Lord 1732;  
ear of our

481 years  
ear 1833?  
8671, the  
he greater

d in part  
ds, what

5. There are four numbers, the first 317, the second 812, the third 1350, and the fourth as much as the other three; what is the sum of them all?

6. A gentleman left his daughter 16 thousand, 16 hundred and 16 pounds; he left his son 1800 more than his daughter; what was his son's portion, and what was the amount of the whole estate?  
*Ans.* { Son's portion, 19,416.  
          { Whole estate, 37,032.

7. A man, at his death, left his estate to his four children, who, after paying debts to the amount of 1476 pounds, received 4768 pounds each; how much was the whole estate?  
*Ans.* 20548.

8. A man bought four hogs, each weighing 375 pounds; how much did they all weigh?  
*Ans.* 1500.

9. The fore quarters of an ox weigh one hundred and eight pounds each, the hind quarters weigh one hundred and twenty-four pounds each, the hide seventy-six pounds, and the tallow sixty pounds; what is the whole weight of the ox?  
*Ans.* 600.

10. A man, being asked his age, said he was thirty-four years old when his eldest son was born, who was then fifteen years of age; what was the age of the father?

11. A man sold two cows for five pounds each, twenty bushels of corn for three pounds, and one hundred pounds of tallow for two pounds; what was his due?

## SUBTRACTION

### OF SIMPLE NUMBERS.

1. Charles, having 11 pence, bought a book, for which he gave 5 pence; how many pence had he left?

2. John had 12 apples; he gave 5 of them to his brother; how many had he left?

3. Peter played at marbles; he had 23 when he began, but when he had done he had only 12; how many did he lose?

4. A man bought an article for 17 shillings and sold it again for 22 shillings; how many shillings did he gain?

5. Charles is 9 years old, and Andrew is 13; what is the difference in their ages?

6. A man borrowed 50 pounds, and paid all but 18; how many pounds did he pay? that is, take 18 from 50, and how many would there be left?

7. John bought several articles for 19 shillings; he gave for 4 books 8 shillings; what did the other articles cost him?

8. Peter bought a trunk for 17 shillings, and sold it for 22 shillings; how many shillings did he gain by the bargain?

9. Peter sold a waggon for 22 shillings, which was 5 shillings more than he gave for it; how many shillings did he give for the waggon?

10. A boy, being asked how old he was, said that he was 25 years younger than his father, whose age was 33 years; how old was the boy?

The taking of a less number from a greater (as in the foregoing examples) is called *Subtraction*. The greater number is called the *minuend*, the less number the *subtrahend*, and what is left after subtraction is called the *difference*, or *remainder*.

11. If the minuend be 8, and the subtrahend 3, what is the difference or remainder? *Ans.* 5.

12. If the subtrahend be 4, and the minuend 16, what is the remainder?

13. Samuel bought a book for 11 pence; he paid down 4 pence; how many pence more must he pay?

**SIGN.** A short horizontal line, —, is the sign of subtraction. It is usually read *minus*, which is a Latin word signifying *less*. It shows that the number *after* it is to be taken from the number *before* it. Thus,  $8 - 3 = 5$ , is read 8 minus or less 3 is equal to 5; or 3 from 8 leaves 5. The latter expression is to be used by the pupil in committing the following

2

3

4

5

6

7

8

9

10

3

4

5

7 - 3

8 - 5

9 - 4

12 - 3

13 - 4

¶ 7.

examples

done in

operation

ore it is

aming

14. A

hem by

Here

e take

undred

he less

o place

From 2

Take 1

SUBTRACTION TABLE.

2-2=0	6-3=3	5-5=0	7-7=0
3-2=1	7-3=4	6-5=1	8-7=1
4-2=2	8-3=5	7-5=2	9-7=2
5-2=3	9-3=6	8-5=3	10-7=3
6-2=4	10-3=7	9-5=4	8-8=0
7-2=5	4-4=0	10-5=5	9-8=1
8-2=6	5-4=1	6-6=0	10-8=2
9-2=7	6-4=2	7-6=1	9-9=0
10-2=8	7-4=3	8-6=2	10-9=1
3-3=0	8-4=4	9-6=3	
4-3=1	9-4=5	10-6=4	
5-3=2	10-4=6		

7-3= how many ?

18-7= how many ?

8-5= how many ?

28-7= how many ?

9-4= how many ?

22-13= how many ?

12-3= how many ?

33-5= how many ?

13-4= how many ?

41-15= how many ?

77. When the numbers are *small*, as in the foregoing examples, the taking of a less number from a greater is readily done in the *mind*; but when the numbers are *large*, the operation is most easily performed part at a time, and therefore it is necessary to *write* the numbers down before performing the operation.

14. A farmer having a flock of 237 sheep, lost 114 of them by disease; how many had he left ?

Here we have 4 units to be taken from 7 units, 1 ten to be taken from 3 tens, and 1 hundred to be taken from 2 hundreds. It will therefore be most convenient to write the less number under the greater, observing, as in addition, to place units under units, tens under tens, &c. thus :

OPERATION.

From 237 the *minuend*,  
Take 114 the *subtrahend*,

123 the *remainder*.

We now begin with the units, saying, 4 (units) from 7, (units,) and there remain 3, (units,) which we set down directly under the column in unit's place.



Then, proceeding to the next column we say 1 (ten) from 3, (tens,) and there remain 2 (tens,) which we set down in *ten's* place. Proceeding to the next column, we say, 1 (hundred) from 2, (hundreds,) and there remains 1, (hundred,) which we set down in *hundred's* place, and the work is done. It now appears, that the number of sheep left was 123 : that is,  $237 - 114 = 123$ .

After the same manner are performed the following examples :

15. There are two farms ; one is valued at 973 pounds, and the other at 421 pounds ; what is the difference in the value of the two farms ?

16. A man's property is worth 2170 pounds, but he has debts to the amount of 1110 pounds ; what will remain after paying his debts.

17. James, having 15 shillings bought a book for which he gave 7 shillings ; how many shillings had he left ?

OPERATION.

15 shillings.

7 shillings.

—

8 shillings left.

A difficulty presents itself here ; for we cannot take 7 from 5 ; but we can take 7 from 15, and there will remain 8.

18. A man bought several articles for 85 pounds, and other articles for 27 pounds ; what did the former cost him more than the latter ?

OPERATION. The same difficulty meets us here as in *First articles*, 85 the last example ; we cannot take 7 from *Other articles*, 27 5 ; but in the last example the larger number consisted of 1 ten and 5 units, which

*Difference*, 58 together make 15 ; we therefore took 7 from 15. Here we have 8 tens and 5 units. We can now, in the mind, suppose 1 ten taken from the 8 tens, which would leave 7 tens, and this 1 ten we can suppose joined to the 5 units, making 15. We can now take 7 from 15, as before, and there will remain 8, which we set down. The taking of 1 ten out of 8 tens, and joining it with the 5 units, is called *borrowing ten*. Proceeding to the next higher order, or tens, we must consider the upper figure 8, from which we borrowed, 1 less, calling it seven ; then, taking 2 (tens) from 7 (tens) there will remain five (tens,) which we set down, making the difference 58. Or, instead of mak-

¶ 7, 8  
ing th  
lower  
same ;  
19.  
how m  
many  
20.  
21.  
¶ 8  
the re  
22.  
many  
23.  
had h  
24.  
for 16  
75 —  
25.  
many  
26.  
many  
Her  
forego  
To  
subtra  
equal  
To  
amou  
it, an  
Thus  
7—7=  
Fr  
the f  
  
I.  
placi  
line  
II.  
in th  
rema  
III.  
figur

1 (ten) from  
set down in  
, we say, 1  
ins 1, (hun-  
nd the work  
leep left was

llowing ex

pounds, and  
in the value

, but he has  
ll remain af

ok for which  
e left ?

here; for we  
ve can take  
men 8.

pounds, and  
ner cost him

as here as in  
take 7 from  
larger num-  
units, which  
efore took 7

We can now,  
tens, which

ose joined to  
from 15, as

down. The

n the 5 units,

t higher or-  
ure 8, from

then, taking  
tens,) which  
tead of mak-

ing the *upper* figure 1 *less*, calling it 7, we may make the *lower* figure 1 *more*, calling it 3, and the result will be the same; for 3 from 8 leaves 5, the same as 2 from 7.

19. A man borrowed 713 pounds, and paid 471 pounds; how many pounds did he then owe?  $713 - 471 =$  how many? *Ans.* 242 pounds.

20.  $1612 - 465 =$  how many? *Ans.* 1147.

21.  $43751 - 6782 =$  how many? *Ans.* 36969.

¶ 8. The pupil will readily perceive, that subtraction is the *reverse* of addition.

22. A man bought 40 sheep, and sold 18 of them; how many had he left?  $40 - 18 =$  how many? *Ans.* 22 sheep.

23. A man sold 18 sheep, and had 22 left; how many had he at first?  $18 + 22 =$  how many?

24. A man bought some articles for 75 pounds, and others for 16 pounds; what was the difference of the costs?  $75 - 16 =$  how many? Reversed,  $59 + 16 =$  how many?

25.  $114 - 103 =$  how many? Reversed,  $11 + 103 =$  how many?

26.  $143 - 76 =$  how many? Reversed,  $67 + 76 =$  how many?

Hence, subtraction may be proved by *addition*, as in the foregoing examples, and addition by *subtraction*.

*To prove subtraction*, we may add the remainder to the *subtrahend*, and, if the work is right, the amount will be equal to the *minuend*.

*To prove addition*, we may *subtract*, successively, from the amount, the *several numbers* which were added to produce it, and, if the work is right, there will be no remainder. Thus  $7 + 8 + 6 = 21$ ; *proof*,  $21 - 6 = 15$ , and  $15 - 8 = 7$ , and  $7 - 7 = 0$ .

From the remarks and illustrations now given, we deduce the following

### RULE.

I. Write down the numbers, the less under the greater, placing units under units, tens under tens, &c. and draw a line under them.

II. Beginning with units, take successively each figure in the *lower* number from the figure over it, and write the remainder directly below.

III. When the figure in the lower number *exceeds* the figure over it, suppose 10 to be added to the *upper* figure;

but in this case we must add 1 to the *lower* figure in the next column *before* subtracting. This is called *borrowing* 10.

### EXAMPLES FOR PRACTICE.

27. If a farm and the buildings on it be valued at 3000 pounds, and the buildings alone be valued at 1500 pounds, what is the value of the land?

28. The population of the two Canadas, at the last census, was 789,000, at the census before, the population was 648,000; what was the difference in the two enumerations?

29. What is the difference between 7,648,203 and 928,671?

30. How much must you add to 358,642 to make 1,487,945?

31. A man bought an estate for 3798 pounds, and sold it again for 4137 pounds; did he gain or lose by it? and how much?

32. From 364,710,825,193 take 27,940,386,574.

33. From 831,025,405,270 take 651,308,604,782.

34. From 127,368,047,216,842 take 978,654,327,352.

### SUPPLEMENT

### TO SUBTRACTION.

#### QUESTIONS.

1. What is *subtraction*? 2. What is the *greater* number called? 3. — the *less* number? 4. What is the *result* or *answer* called? 5. What is the *sign* of subtraction? 6. What is the *rule*? 7. What is understood by *borrowing ten*? 8. Of what is subtraction the *reverse*? 9. How is subtraction proved? 10. How is addition proved by subtraction?

#### EXERCISES.

1. How long from the discovery of America by Columbus, in 1492, to the period of the cession by France of all her possessions in North America to Great Britain, in 1763.

2. Supposing a man to have been born in the year 1773, how old was he in 1827?

3. Supposing a man to have been 80 years old in the year 1826, in what year was he born?

4. There are two numbers, whose difference is 8764; the greater number is 15687; I demand the less?

figure in the  
borrowing 10.

ued at 3000  
500 pounds,

the last cen-  
sulation was  
umerations 1  
48,203 and

2 to make

, and sold it  
! and how

,574.

4,783.

,927,352.

umber called?  
number called?  
e 7. What  
action the re-  
dition proved

by Colum-  
ance of all  
n, in 1763.  
year 1773,

old in the

8764; the

5. What number is that which, taken from 3794, leaves 865?

6. What number is that to which if you add 789, it will become 6350?

7. In a certain city, there were 123,706 inhabitants; in another 43,940; how many more inhabitants were there in one than in the other?

8. A man possessing an estate of twelve thousand pounds, gave two thousand five hundred pounds to each of his two daughters, and the remainder to his son; what was his son's share?

9. From seventeen million take fifty-six thousand, and what will remain?

10. What number, together with these three, viz. 1801, 2561, and 3120, will make ten thousand?

11. A man bought a horse for 35 pounds, and a chaise for 47 pounds; how much more did he give for the chaise than for the horse?

12. A man borrows 7 ten dollar bills and three one dollar bills, and pays at one time 4 ten dollar bills and 5 one dollar bills; how many ten dollar bills and one dollar bills must he afterwards pay to cancel the debt?

*Ans.* 3 ten doll. bills and 8 one doll.

13. The greater of two numbers is 24, and the less is 16; what is their difference?

14. The greater of two numbers is 24, and their difference 8; what is the less number?

15. The sum of two numbers is 40, the less is 16; what is the greater?

16. A tree, 68 feet high, was broken off by the wind; the top part, which fell, was 49 feet long; how high was the stump which was left?

17. Elizabeth became Queen of England in 1558; how many years since?

18. A man carried his produce to market; he sold his pork for 14 pounds, his cheese for 11 pounds, and his butter for 9 pounds; he received, in pay, salt to the value of 6 pounds, 3 pounds worth of sugar, 2 pounds worth of molasses, and the rest in money; how much money did he receive?

*Ans.* 23 pounds.

19. A boy bought several sleds for 13 shillings, and gave 6 shillings to have them repaired; he sold them for 18 shill-

ings ; did he gain or lose by the bargain ? and how much ?

20. One man travels 67 miles in a day, another man follows at the rate of 42 miles a day ; if they both start from the same place at the same time, how far will they be apart at the close of the first day ? — of the second ? — of the third ? — of the fourth ?

21. One man starts from York Monday morning, and travels at the rate of 40 miles a day ; another starts from the same place Tuesday morning, and follows at the rate of 70 miles a day ; how far are they apart Tuesday night ?

*Ans.* 10 miles.

22. A man owing 379 pounds, paid at one time 47 pounds, at another time 84 pounds, at another time, 27 pounds, and at another 143 pounds ; how much did he then owe ?

*Ans.* 82 pounds.

23. A man has property to the amount of 34764 pounds, but there are demands against him to the amount of 14297 pounds ; how many pounds will be left after the payment of his debts ?

24. Four men bought a lot of land for 482 pounds ; the first man paid 274 pounds, the second man 194 pounds less than the first, and the third man 20 pounds less than the second ; how much did the second, third, and the fourth man pay ?

*Ans.* { The second paid 80.  
The third paid 60.  
The fourth paid 68.

25. A man, having 10,000 pounds, gave away 9 pounds ; how many had he left ?

*Ans.* 9991.

## MULTIPLICATION

### OF SIMPLE NUMBERS.

¶ 8. 1. If one orange cost 2 pence, how many pence must I give for 2 oranges ? — how many pence for 3 oranges ? — for 4 oranges ?

2. One bushel of apples cost 3 shillings ; how many shillings must I give for 2 bushels ? — for 3 bushels ?

3. One gallon contains 4 quarts ; how many quarts in 2 gallons ? — in 3 gallons ? — in 4 gallons ?

4. Three men bought a horse ; each man paid 6 pounds

how much ?  
 other man fol-  
 both start from  
 they be apart  
 ond ! — of

morning, and  
 or starts from  
 at the rate of  
 day night ?  
 ns. 10 miles.  
 e 47 pounds,  
 pounds, and  
 owe ?

a. 82 pounds.  
 1764 pounds,  
 ant of 14297  
 the payment

pounds ; the  
 pounds less  
 less than the  
 d the fourth  
 ond paid 80.  
 rd. paid 60.  
 rth paid 68.  
 y 9 pounds ;  
 Ans. 9991.

many pence  
 pence for 3

how many  
 bushels ?  
 quarts in 2

6 pounds

for his share ; how many pounds did the horse cost them ?

5. A man has 4 farms worth 95 pounds each ; how many pounds are they all worth ?

6. In one pound there are 20 shillings ; how many shillings in 5 pounds ?

7. How much will 4 pair of shoes cost at 9 shillings a pair ?

8. How much will 3 pounds of tea cost at 5 shillings a pound ?

9. There are 24 hours in 1 day ; how many hours in 2 days ? — in 3 days ? — in 4 days ? — in 7 days ?

10. Six boys met a beggar and gave him 9 pence each ; how many pence did the beggar receive ?

When questions occur, (as in the above examples,) where the same number is to be added to itself several times, the operation may be much facilitated by a rule, called *Multiplication*, in which the number to be repeated is called the *multiplicand*, and the number which shows how many times the multiplicand is to be repeated is called the *multiplier*. The multiplicand and multiplier, when spoken of *collectively*, are called the *factors*, (producers,) and the answer is called the *product*.

11. There is an orchard in which there are 5 rows of trees, and 27 trees in each row ; how many trees in the orchard ?

In the first row, 27 trees.  
 " second " 27 "  
 " third " 27 "  
 " fourth " 27 "  
 " fifth " 27 "

In this example, it is evident that the whole number of trees will be equal to the amount of five 27's added together.

In adding, we find that 7 taken five times amounts to 35. We write

down the five units, and reserve the 3 tens ; the amount of 2 taken five times is 10, and the 3, which we reserved, makes 13, which, written to the left of units, makes the whole number of trees 135.

If we have learned that 7 taken 5 times amounts to 35, and that 2 taken 5 times amounts to 10, it is plain we need write the number 27 but *once*, and then, setting the multiplier under it, we may say, 5 times 7 are 35, writing down



the 5, and reserving the 3 (tens) as in addition. Again 5

<i>Multiplicand,</i>	27 trees in each row.	times 2 (tens) are
<i>Multiplier,</i>	5 rows.	10 (tens,) and 3,
		(tens,) which we
		reserved, make 13,
<i>Product,</i>	135 trees, <i>Ans.</i>	(tens,) as before.

¶ 10. 12. There are on a board 3 rows of spots, and 4 spots in each row ; how many spots on the board ?

⊗ ⊗ ⊗ ⊗      A slight inspection of the figure will  
 ⊗ ⊗ ⊗ ⊗      show that the number of spots may be  
 ⊗ ⊗ ⊗ ⊗      found either by taking 4 three times, (3  
                                  times 4 are 12,) or by taking 3 four times,  
                                  (4 times 3 are 12 ;) for we may say there  
 are three rows of 4 spots each, or 4 rows of 3 spots each ;  
 therefore, we may use either of the given numbers for a  
 multiplier, as best suits our convenience. We generally  
 write the numbers as in subtraction, the larger number up-  
 permost, with units under units, tens under tens, &c. Thus,

<i>Multiplicand,</i>	4 spots.	<i>Note.</i>	4 and 3 are the factors,
<i>Multiplier,</i>	3 rows.		which produce the product 12.

*Product,*      12 *Ans.*

Hence,—*Multiplication is a short way of performing many additions ; in other words,—It is the method of repeating any number any given number of times.*

**SIGN.** Two short lines crossing each other in the form of the letter X, are the sign of multiplication. Thus,  $3 \times 4 = 12$ , signifies that 3 times 4 are equal to 12, or 4 times 3 are 12.

*Note.* Before any progress can be made in this rule, the following table must be committed perfectly to memory.

¶ 10.

times 0
2 × 1
2 × 2
2 × 3
2 × 4
2 × 5
2 × 6
2 × 7
2 × 8
2 × 9
2 × 10
2 × 11
2 × 12
× 0
× 1
× 2
× 3
× 4
× 5
× 6
× 7
× 8
× 9
× 10
× 11
× 12
× 0
× 1
× 2
× 3
× 4
× 5
× 6
× 7
× 8
× 9

# MULTIPLICATION TABLE.

1 times 0 are 0	4 × 10 = 40	7 × 7 = 49	10 × 4 = 40
2 × 1 = 2	4 × 11 = 44	7 × 8 = 56	10 × 5 = 50
2 × 2 = 4	4 × 12 = 48	7 × 9 = 63	10 × 6 = 60
2 × 3 = 6	5 × 0 = 0	7 × 10 = 70	10 × 7 = 70
2 × 4 = 8	5 × 1 = 5	7 × 11 = 77	10 × 8 = 80
2 × 5 = 10	5 × 2 = 10	7 × 12 = 84	10 × 9 = 90
2 × 6 = 12	5 × 3 = 15	8 × 0 = 0	10 × 10 = 100
2 × 7 = 14	5 × 4 = 20	8 × 1 = 8	10 × 11 = 110
2 × 8 = 16	5 × 5 = 25	8 × 2 = 16	10 × 12 = 120
2 × 9 = 18	5 × 6 = 30	8 × 3 = 24	11 × 0 = 0
2 × 10 = 20	5 × 7 = 35	8 × 4 = 32	11 × 1 = 11
2 × 11 = 22	5 × 8 = 40	8 × 5 = 40	11 × 2 = 22
2 × 12 = 24	5 × 9 = 45	8 × 6 = 48	11 × 3 = 33
3 × 0 = 0	5 × 10 = 50	8 × 7 = 56	11 × 4 = 44
3 × 1 = 3	5 × 11 = 55	8 × 8 = 64	11 × 5 = 55
3 × 2 = 6	5 × 12 = 60	8 × 9 = 72	11 × 6 = 66
3 × 3 = 9	6 × 0 = 0	8 × 10 = 80	11 × 7 = 77
3 × 4 = 12	6 × 1 = 6	8 × 11 = 88	11 × 8 = 88
3 × 5 = 15	6 × 2 = 12	8 × 12 = 96	11 × 9 = 99
3 × 6 = 18	6 × 3 = 18	9 × 0 = 0	11 × 10 = 110
3 × 7 = 21	6 × 4 = 24	9 × 1 = 9	11 × 11 = 121
3 × 8 = 24	6 × 5 = 30	9 × 2 = 18	11 × 12 = 132
3 × 9 = 27	6 × 6 = 36	9 × 3 = 27	12 × 0 = 0
3 × 10 = 30	6 × 7 = 42	9 × 4 = 36	12 × 1 = 12
3 × 11 = 33	6 × 8 = 48	9 × 5 = 45	12 × 2 = 24
3 × 12 = 36	6 × 9 = 54	9 × 6 = 54	12 × 3 = 36
4 × 0 = 0	6 × 10 = 60	9 × 7 = 63	12 × 4 = 48
4 × 1 = 4	6 × 11 = 66	9 × 8 = 72	12 × 5 = 60
4 × 2 = 8	6 × 12 = 72	9 × 9 = 81	12 × 6 = 72
4 × 3 = 12	7 × 0 = 0	9 × 10 = 90	12 × 7 = 84
4 × 4 = 16	7 × 1 = 7	9 × 11 = 99	12 × 8 = 96
4 × 5 = 20	7 × 2 = 14	9 × 12 = 108	12 × 9 = 108
4 × 6 = 24	7 × 3 = 21	10 × 0 = 0	12 × 10 = 120
4 × 7 = 28	7 × 4 = 28	10 × 1 = 10	12 × 11 = 132
4 × 8 = 32	7 × 5 = 35	10 × 2 = 20	12 × 12 = 144
4 × 9 = 36	7 × 6 = 42	10 × 3 = 30	

Again 5 (tens) are 5, and 3, which we make 13, as before.

dots, and 4 1! figure will ts may be e times, (3 four times, say there dots each; nbers for a generally umber up- &c. Thus, he factors, duct 12.

ning many eating any

the form us, 3 × 4 4 times 3

rule, the nory.

- $9 \times 2 =$  how many ?       $4 \times 3 \times 2 = 24$ .  
 $4 \times 6 =$  how many ?       $3 \times 2 \times 5 =$  how many ?  
 $8 \times 9 =$  how many ?       $7 \times 1 \times 2 =$  how many ?  
 $3 \times 7 =$  how many ?       $8 \times 3 \times 2 =$  how many ?  
 $5 \times 5 =$  how many ?       $3 \times 2 \times 4 \times 5 =$  how many ?

13. What will 84 barrels of flour cost at 2 pounds a barrel ? *Ans.* 168 pounds.

14. A merchant bought 12 dozen hats at the rate of 12 pounds per dozen ; what did they cost ? *Ans.* 144 pounds.

15. How many inches are there in 253 feet, every foot being 12 inches ?

**OPERATION.** The product of 12, with each of the significant figures or digits, having been committed to memory from the multiplication table, it is just as easy to multiply by 12 as by a single figure. Thus, 12 times 3 are 36, &c.

*Ans.* 3036  
 16. What will 476 barrels of fish cost at 3 pounds a barrel ? *Ans.* 1428 pounds.

17. A piece of very valuable land, containing 33 acres, was sold for 246 pounds an acre ; what did the whole come to ?

As 12 is the largest number, the product of which, with the nine digits, is found in the multiplication table, therefore, when the multiplier exceeds 12, we multiply by each figure in the multiplier separately. Thus :

**OPERATION.**

246 pounds, the price of one acre.

33 number of acres.

738 pounds, the price of three acres.

738 pounds, the price of thirty acres,

*Ans.* 8118 pounds, the price of 33 acres.

price of 3 acres. We then multiply by the 3 tens, writing the first figure of the product (8) in *ten's* place, that is, directly under the figure by which we multiply. It now appears that the product by the 3 tens consists of the same figures as the product by the 3 units ; but there is this difference—the figures in the product by the 3 tens are all removed one place further to the left hand, by which their value is increased *tenfold*, which is as it should be, because the price

The multiplier consists of 3 tens and 3 units.

First, multiplying by the 3 units gives us 738 pounds the

of 30 acres, the product of these lowing

I. W. multipl and dra

II. V. right ha contain ing the

III. figure remem duct d Having the mu sum of

18. there

19. how n

20. chest

21. pound

22. an ho

rate 1

year.

23. cont

were

24.

25.

26.

many ?  
many ?  
many ?  
how many ?

of 30 acres is evidently ten times as much as the price of 3 acres, that is, 7380 pounds; and it is plain, that these two products, added together, give the price of 33 acres.

These examples will be sufficient to establish the following

### RULE.

I. Write down the multiplicand, under which write the multiplier, placing units under units, tens under tens, &c., and draw a line underneath.

II. When the multiplier does *not* exceed 12, begin at the right hand of the multiplicand, and multiply each figure contained in it by the multiplier, setting down, and carrying the same as in addition.

III. When the multiplier *exceeds* 12, multiply by each figure separately, first by the *units*, then by the *tens*, &c., remembering always to place the first figure of each product directly under the figure by which you multiply. Having gone through in this manner with each figure in the multiplier, add their several products together, and the sum of them will be the product required.

### EXAMPLES FOR PRACTICE.

18. There are 320 rods in a mile; how many rods are there in 57 miles?

19. Suppose it to be 706 miles from Halifax to Quebec; how many rods is it?

20. What will 784 chests of tea cost, at 17 pounds a chest?

21. If 1851 men receive 758 pounds apiece; how many pounds will they all receive? *Ans.* 1403058 pounds.

22. There are 24 hours in a day; if a ship sail 7 miles in an hour, how many miles will she sail in 1 day, at that rate? how many miles in 36 days? how many miles in 1 year, or 365 days? *Ans.* 61320 miles in 1 year.

23. A merchant bought 13 pieces of cloth, each piece containing 28 yards, at 2 pounds a yard; how many yards, were there, and what was the whole cost?

*Ans.* The whole cost was 728 pounds.

24. Multiply 37864 by 235. *Product,* 8808040.

25. " 29831 " 952. " 28399112.

26. " 93956 " 8704. " 817793024.

## CONTRACTIONS IN MULTIPLICATION.

1. *When the multiplier is a composite number.*

¶ 11. Any number, which may be produced by the multiplication of two or more numbers, is called a *composite number*. Thus, 15, which arises from the multiplication of 5 and 3, ( $5 \times 3 = 15$ ;) is a composite number, and the numbers 5 and 3, which, multiplied together, produce it, are called *component parts* or *factors* of that number. So, also, 24 is a composite number; its *component parts*, or *factors*, may be 2 and 12, ( $2 \times 12 = 24$ ;) or they may be 4 and 6, ( $4 \times 6 = 24$ ;) or they may be 2, 3, and 4. ( $2 \times 3 \times 4 = 24$ .)

1. What will 15 pieces of cloth cost, at 4 pounds a piece?

4	15 pieces are equal to $5 \times 3$ pieces. The cost
5	of 5 pieces would be $5 \times 4 = 20$ pounds; and be-
—	cause 15 pieces contain 3 times 5 pieces, so the
20	cost of 15 pieces will evidently be 3 times the
3	cost of 5 pieces, that is, 20 pounds $\times 3 = 60$ pounds.
—	<i>Ans.</i> 60 pounds.

60

Wherefore, *If the multiplier be a composite number*, we may, if we please, *multiply the multiplicand first by one of the component parts; that product by the other, and so on, if the component parts be more than two; and, having in this way multiplied by each of the component parts, the last product will be the product required.*

2. What will 136 tons of potashes come to, at 24 pounds per ton?

$6 \times 4 = 24$ . It follows, therefore, that 6 and 4 are component parts or factors of 24. Hence,

136 tons.

6 one of the component parts, or factors.

816 pounds, the price of 6 tons.

4 the other component part, or factor.

*Ans.* 3264 pounds, the price of 136 tons.

3. Supposing 342 men to be employed in a certain piece of work, for which they are to receive 28 pounds, each, how much will they all receive?

$7 \times 4 = 28$

*Ans.* 9576 pounds.

4. Multiply 367 by 48.	Product, 17616.
5. " 853 " 56.	" 47768.
6. " 1088 " 72.	" 78192.

II. When the multiplier is 10, 100, 1000, &c.

¶ 12. It will be recollected, (¶ 3.) that any figure, on being removed one place towards the *left* hand, has its value increased *tenfold*; hence, to multiply any number by 10, it is only necessary to *write a cypher on the right hand of it*. Thus, 10 times 25 are 250; for the 5, which was *units* before, is now made *tens*, and the 2, which was *tens*, before, is now made *hundreds*. So, also, if any figure be removed *two* places towards the left hand, its value is increased 100 times, &c. Hence,

When the multiplier is 10, 100, 1000, or 1 with any number of ciphers annexed, annex as many ciphers to the multiplicand as there are ciphers in the multiplier, and the multiplicand, so increased, will be the product required. Thus,

Multiply 46 by	10,	the product is	460.
" 83 "	100, "	" "	8300.
" 95 "	1000, "	" "	95000.

# EXAMPLES FOR PRACTICE.

1. What will 76 loads of corn cost, at 10 pounds a load?
2. If 100 men receive 32 pounds each, how many pounds will they all receive?
3. What will 1000 pieces of broadcloth cost, estimating each piece at 78 pounds?
4. Multiply 5682 by 10000.
5. " 82134 " 100000.

¶ 13. On the principle suggested in the last ¶, it follows,

When there are ciphers on the *right hand* of the multiplicand, multiplier, either or both, we may, at first, neglect these ciphers, multiplying by the *significant figures* only; after which we must annex as many ciphers to the product as there are ciphers on the right hand of the multiplicand and multiplier, counted together.



## EXAMPLES FOR PRACTICE.

1. If 1300 men receive 460 pounds apiece, how many pounds will they all receive ?

OPERATION.

$$\begin{array}{r} 460 \\ 1300 \\ \hline 138 \\ 46 \end{array}$$

*Ans, 598000 pounds.*

The ciphers in the multiplicand and multiplier, counted together, are *three*. Disregarding these, we write the *significant* figures of the multiplier under the *significant* figures of the multiplicand, and multiply ; after which we annex three ciphers to the right hand of the

product, which gives the true answer.

2. The number of distinct buildings in New England, appropriated to the spinning, weaving, and printing of cotton goods, was estimated, in 1826, at 400, running, on an average, 700 spindles each ; what was the whole number of spindles ?

3. Multiply 257 by 6300.

4. " 8600 " 17.

5. " 9340 " 460.

6. " 5200 " 410.

7. " 378 " 204.

OPERATION.

$$\begin{array}{r} 378 \\ 204 \\ \hline 1512 \\ 000 \\ 756 \\ \hline 77112 \end{array}$$

In the operation it will be seen, that multiplying by ciphers produces nothing. Therefore,

III. *When there are ciphers between the significant figures of the multiplier, we may omit the ciphers, multiplying by the significant figures only, placing the first figure of each product directly under the figure by which we multiply.*

## EXAMPLES FOR PRACTICE.

8. Multiply 154326 by 3007.

9.  
10.  
11.

1. W  
ed call  
answer  
multip  
does it  
for mu  
less th  
dure?  
stood  
may y  
To mu  
When  
plier,  
phers  
to be

1.  
so m  
each  
taken  
2.  
be 1  
and  
inha  
3.  
man

OPERATION.

154326

3007

1080282

462978

*Product, 464058282*

- |             |         |        |
|-------------|---------|--------|
| 9. Multiply | 543 by  | 206.   |
| 10. "       | 1620 "  | 2103.  |
| 11. "       | 36243 " | 32004. |

SUPPLEMENT

TO MULTIPLICATION.

QUESTIONS.

1. What is multiplication? 2. What is the number *to be multiplied* called? 3. — to multiply *by* called? 4. What is the *result* or *answer* called? 5. Taken *collectively*, what are the multiplicand and multiplier called? 6. What is the *sign* of multiplication? 7. What does it show? 8. In what *order* must the given numbers be placed for multiplication? 9. How do you proceed when the multiplier is *less* than 12? 10. When it *exceeds* 12, what is the method of procedure? 11. What is a *composite* number? 12. What is to be understood by the *component parts*, or *factors*, of any number? 13. How may you proceed when the multiplier is a *composite number*? 14. To multiply by 10, 100, 1000, &c., what suffices? 15. Why? 16. When there are *ciphers on the right hand* of the multiplicand, multiplier, either or both, how may we proceed? 17. When there are *ciphers between* the significant figures of the multiplier, how are they to be treated?

EXERCISES.

1. An army of 10700 men having plundered a city, took so much money, that, when it was shared among them, each man received 46 pounds; what was the sum of money taken?

2. Supposing the number of houses in a certain town to be 145, each house, on an average, containing two families, and each family 6 members, what would be the number of inhabitants in that town? *Ans. 1740.*

3. If 46 men can do a piece of work in 60 days, how many men will it take to do it in one day?

D

4. Two men depart from the same place, and travel in opposite directions, one at the rate of 27 miles a day, the other 31 miles a day ; how far apart will they be at the end of 6 days ? *Ans.* 348 miles.

5. What number is that, the factors of which are 4, 7, 6, and 20 ? *Ans.* 3360.

6. If 18 men can do a piece of work in 90 days, how long will it take one man to do the same ?

7. What sum of money must be divided between 27 men, so that each man may receive 115 pounds ?

8. There is a certain number, the factors of which are 89 and 265 ; what is that number ?

9. What is that number, of which 9, 12, and 14 are factors ?

10. If a carriage wheel turn round 346 times in running 1 mile, how many times will it turn round in the distance from Quebec to Montreal it being 180 miles. *Ans.* 62280.

11. In one minute are 60 seconds ; how many seconds, in 4 minutes ? — in 5 minutes ? — in 20 minutes ? — in 40 minutes ?

12. In one hour are 60 minutes ; how many seconds : an hour ? — in two hours ? How many seconds from nine o'clock in the morning till noon ?

13. In one pound are 4 dollars ; how many dollars in 3 pounds ? — in 300 pounds ? — in 467 pounds ?

14. Two men, A and B, start from the same place at the same time, and travel the same way ; A travels 52 miles a day, and B 44 miles a day ; how far apart will they be at the end of 10 days ?

15. If the interest of 100 pounds, for one year, be six pounds, how many pounds will be the interest for 2 years ? — for 4 years ? — for 10 years ? for 35 years ? — for 84 years ?

16. If the interest of one hundred pounds, for one year, be six pounds, what is the interest for two hundred pounds the same time ? — 7 hundred pounds ? — 8 hundred pounds ? — 95 hundred pounds ?

17. A farmer sold 468 pounds of pork, at 3 pence a pound, and 48 pounds of cheese, at 4 pence a pound ; how many pence must he receive in pay ?

18. A boy bought 10 oranges ; he kept 7 of them, and sold

the other

19. 7

6, 9, 8,

20. 1

in 8 ho

quarts i

pints in

† 14

how ma

2. Ja

how ma

3. Jo

who rec

them to

4. If

at 2 pe

5. H

at 2 po

6. If

is one y

7. A

chilling

8. H

— in

9. H

10. I

would

11. I

and the

Rem

parts,

parts,

parts,

one pa

12.

each of

and travel  
s a day, the  
be at the  
348 miles.  
are 4, 7, 6,  
*Ans.* 3360.  
s, how long

the others for 5 pence a piece; how many pence did he receive?

19. The component parts of a certain number are 4, 5, 7, 6, 9, 8, and 3; what is the number?

20. In 1 hogshead are 63 gallons; how many gallons in 8 hogsheads? In 1 gallon are 4 quarts; how many quarts in 8 hogsheads? In 1 quart are 2 pints; how many pints in 8 hogsheads?

pen 27 men,

which are 89

14 are fac-

in running  
he distance

ns. 62280.  
seconds, in  
utes? —

r seconds  
conds from

dollars in  
unds?

place at the  
52 miles a  
they be at

ear, be six  
or 2 years?  
ears? —

one year,  
red pounds  
8 hundred

3 pence a  
und; how

n, and sold

## ***DIVISION*** OF SIMPLE NUMBERS.

¶ 14. 1. James divided 12 apples among four boys; how many did he give each boy?

2. James would divide 12 apples among three boys; how many must he give each boy?

3. John had 15 apples, and gave them to his playmates, who received 3 apples each; how many boys did he give them to?

4. If you had 20 pence, how many cakes could you buy at 2 pence a piece?

5. How many yards of cloth could you buy for 30 pounds, at 2 pounds a yard?

6. If you pay 250 shillings for 10 yards of cloth, what is one yard worth?

7. A man works 6 days for 42 shillings; how many shillings is that for one day?

8. How many quarts in 4 pints? — in 6 pints? — in 10 pints?

9. How many times is 8 contained in 88?

10. If a man can travel 4 miles an hour, how many hours would it take him to travel 24 miles?

11. In an orchard there are 28 trees standing in rows, and there are 3 trees in a row; how many rows are there?

*Remark.* When any one thing is divided into two equal parts, one of those parts is called a *half*; if into 3 equal parts, one of those parts is called a *third*; if into four equal parts, one part is called a *quarter* or a *fourth*; if into five, one part is called a *fifth*, and so on.

12. A boy had two apples, and gave one half an apple to each of his companions; how many were his companions?

13. A boy divided four apples among his companions, by giving them one third of an apple each; among how many did he divide his apples?

14. How many quarters in 3 oranges?

15. How many oranges would it take to give 12 boys one quarter of an orange each?

16. How much is one half of 12 apples?

17. How much is one third of 12?

18. How much is one fourth of 12?

19. A man had 30 sheep, and sold one fifth of them; how many of them did he sell?

20. A man purchased sheep for 13 shillings apiece, and paid for them all 117 shillings; what was their number?

21. How many oranges, at 3 pence each, may be bought for 12 pence?

It is plain, that as many times as 3 pence can be taken from 12 pence, so many oranges may be bought; the object, therefore, is to find how many times 3 is contained in 12.

12 pence.  
*First orange,* 3 pence.

—  
9

*Second orange,* 3 pence.

—  
6

*Third orange,* 3 pence.

—  
3

*Fourth orange,* 3 pence.

—  
0

We see in this example, that we may take 3 from 12 four times, after which there is no remainder; consequently, *subtraction* alone is sufficient for the operation; but we may come to the same result by a process, in most cases much shorter, called *Division*.

¶ 15. It is plain, that the cost of one orange, (3 pence,) multiplied by the number of oranges, (4,) is equal to the cost of all the oranges, (12 pence;) 12 is, therefore, a *product*, and 3 one of its factors; and to find how many times 3 is contained in 12 is to find the *other* factor, which, multiplied into 3, will produce 12. This factor we find, by trial, to be 4, ( $4 \times 3 = 12$ ;) consequently, is contained in 12 4 times. *Ans. 4 oranges.*

22. A man would divide 12 oranges equally among 3 children; how many oranges would each child have?

Here the object is to divide the 12 oranges into 3 equal parts, and to ascertain the number of oranges in each of

¶ 15.

those p  
ample, a  
by 3, w  
to be 4.

As, t  
many a  
way of  
and ma  
one num  
number  
cess of  
given p

The  
and an  
ber giv  
one of  
sought,  
how m

SIGN  
between  
is to be  
is read,  
pression  
dots, th  
visor u  
before.

The  
ing ta  
time, 2

2  
4  
6  
8  
10  
12  
14  
16  
18  
20  
22  
24  
26  
28  
30  
32  
34  
36  
38  
40  
42  
44  
46  
48  
50  
52  
54  
56  
58  
60  
62  
64  
66  
68  
70  
72  
74  
76  
78  
80  
82  
84  
86  
88  
90  
92  
94  
96  
98  
100

those parts. The operation is evidently as in the last example, and consists in finding a number, which, multiplied by 3, will produce 12. This number we have already found to be 4. *Ans.* 4 oranges apiece.

As, therefore, *multiplication* is a short way of performing many *additions* of the same number; so *division* is a short way of performing many *subtractions* of the same number; and may be defined, *The method of finding how many times one number is contained in another; and also of dividing a number into any number of equal parts.* In all cases, the process of division consists in finding one of the factors of a given product, when the other factor is known.

The number given to be divided, is called the *dividend*, and answers to the *product* in multiplication. The number given to divide by is called the *divisor*, and answers to one of the factors in multiplication. The *result*, or *answer* sought, is called the *quotient*, (from the Latin word *quoties*, how many ?) and answers to the other factor.

**SIGN.** The sign for division is a short horizontal line between two dots,  $\div$ . It shows that the number *before* it is to be divided by the number *after* it. Thus  $27 \div 9 = 3$  is read, 27 divided by 9 is equal to 3; or to shorten the expression, 27 by 9 is 3; or 9 in 27 3 times. In place of the dots, the *dividend* is often written over the line, and the *divisor* under it, to express division; thus,  $\overset{27}{\underset{9}{\div}} = 3$ , read as before.

The reading used by the pupil in committing the following table may be 2 by 2 is 1, 4 by 2, &c. or 2 in 2 one time, 2 in 4 two times, &c.

### DIVISION TABLE.

$\frac{2}{2} = 1$	$\frac{3}{3} = 1$	$\frac{4}{4} = 1$	$\frac{5}{5} = 1$	$\frac{6}{6} = 1$	$\frac{7}{7} = 1$
$\frac{4}{2} = 2$	$\frac{6}{3} = 2$	$\frac{8}{4} = 2$	$\frac{10}{5} = 2$	$\frac{12}{6} = 2$	$\frac{14}{7} = 2$
$\frac{6}{3} = 3$	$\frac{9}{3} = 3$	$\frac{12}{4} = 3$	$\frac{15}{5} = 3$	$\frac{18}{6} = 3$	$\frac{21}{7} = 3$
$\frac{8}{4} = 4$	$\frac{12}{3} = 4$	$\frac{16}{4} = 4$	$\frac{20}{5} = 4$	$\frac{24}{6} = 4$	$\frac{28}{7} = 4$
$\frac{10}{5} = 5$	$\frac{15}{3} = 5$	$\frac{20}{4} = 5$	$\frac{25}{5} = 5$	$\frac{30}{6} = 5$	$\frac{35}{7} = 5$
$\frac{12}{6} = 6$	$\frac{18}{3} = 6$	$\frac{24}{4} = 6$	$\frac{30}{5} = 6$	$\frac{36}{6} = 6$	$\frac{42}{7} = 6$
$\frac{14}{7} = 7$	$\frac{21}{3} = 7$	$\frac{28}{4} = 7$	$\frac{35}{5} = 7$	$\frac{42}{6} = 7$	$\frac{49}{7} = 7$
$\frac{16}{8} = 8$	$\frac{24}{3} = 8$	$\frac{32}{4} = 8$	$\frac{40}{5} = 8$	$\frac{48}{6} = 8$	$\frac{56}{7} = 8$
$\frac{18}{9} = 9$	$\frac{27}{3} = 9$	$\frac{36}{4} = 9$	$\frac{45}{5} = 9$	$\frac{54}{6} = 9$	$\frac{63}{7} = 9$



## DIVISION TABLE—CONTINUED.

$\frac{8}{8} = 1$	$\frac{8}{8} = 1$	$\frac{16}{16} = 1$	$\frac{11}{11} = 1$	$\frac{13}{13} = 1$
$\frac{16}{8} = 2$	$\frac{16}{8} = 2$	$\frac{32}{16} = 2$	$\frac{22}{11} = 2$	$\frac{26}{13} = 2$
$\frac{24}{8} = 3$	$\frac{24}{8} = 3$	$\frac{48}{16} = 3$	$\frac{33}{11} = 3$	$\frac{39}{13} = 3$
$\frac{32}{8} = 4$	$\frac{32}{8} = 4$	$\frac{64}{16} = 4$	$\frac{44}{11} = 4$	$\frac{52}{13} = 4$
$\frac{40}{8} = 5$	$\frac{40}{8} = 5$	$\frac{80}{16} = 5$	$\frac{55}{11} = 5$	$\frac{65}{13} = 5$
$\frac{48}{8} = 6$	$\frac{48}{8} = 6$	$\frac{96}{16} = 6$	$\frac{66}{11} = 6$	$\frac{78}{13} = 6$
$\frac{56}{8} = 7$	$\frac{56}{8} = 7$	$\frac{112}{16} = 7$	$\frac{77}{11} = 7$	$\frac{91}{13} = 7$
$\frac{64}{8} = 8$	$\frac{64}{8} = 8$	$\frac{128}{16} = 8$	$\frac{88}{11} = 8$	$\frac{104}{13} = 8$
$\frac{72}{8} = 9$	$\frac{72}{8} = 9$	$\frac{144}{16} = 9$	$\frac{99}{11} = 9$	$\frac{117}{13} = 9$

$28 \div 7$ , or  $\frac{28}{7} =$  how many ?  $49 \div 7$ , or  $\frac{49}{7} =$  how many ?  
 $42 \div 6$ , or  $\frac{42}{6} =$  how many ?  $32 \div 4$ , or  $\frac{32}{4} =$  how many ?  
 $54 \div 9$ , or  $\frac{54}{9} =$  how many ?  $99 \div 11$ , or  $\frac{99}{11} =$  how many ?  
 $32 \div 8$ , or  $\frac{32}{8} =$  how many ?  $84 \div 12$ , or  $\frac{84}{12} =$  how many ?  
 $33 \div 11$ , or  $\frac{33}{11} =$  how many ?  $108 \div 12$ , or  $\frac{108}{12} =$  how many ?

¶ 16. 23. How many yards of cloth, at 4 shillings a yard, can be bought for 856 shillings ?

Here the number to be divided is 856, which therefore is the *dividend*; 4 is the number to divide by, and therefore the *divisor*. It is not evident how many times 4 is contained in so large a number as 856. This difficulty will be readily overcome, if we decompose this number, thus :

$$856 = 800 + 40 + 16.$$

Beginning with the hundreds, we readily perceive that 4 is contained in 8 2 times; consequently, in 800 it is contained 200 times. Proceeding to the tens, 4 is contained in 4 1 time, and consequently in 40 it is contained 10 times. Lastly, in 16 it is contained 4 times. We now have  $200 + 10 + 4 = 214$  for the quotient, or the number of times 4 is contained in 856.

*Ans.* 214 yards.

We may arrive to the same result without decomposing the dividend, except as it is done in the mind, taking it by parts, in the following manner :

*Dividend.*

Divisor, 4 ) 856

Quotient, 214

For the sake of convenience, we write down the dividend with the divisor on the left, and draw a line between them; we also draw a line underneath.

Then, beginning on the left hand, we seek how often the

divisor (4)  
 finding it  
 which fall  
 Proceeding  
 which we  
 But after  
 ten left.  
 Then, 4  
 work is  
 This m  
 Division.  
 on partly  
 the divis

From  
rule for

1. Fi  
 first figu  
 ly unde  
 next fig

II. F  
 dividen  
 all the

PRO  
 tient a  
 have a  
 merely  
 Hen  
 other.

To  
 tient,  
 as the  
 tient,  
 as the  
 To  
 factor  
 other

24  
 men

divisor (4) is contained in 8, (hundreds,) the left hand figure ; finding it to be 2 times, we write 2 directly under the 8, which falling in the place of hundreds, is in reality 200. Proceeding to tens, 4 is contained in 5 (tens) 1 time, which we set down in *ten's* place, directly under the 5 (tens.) But after taking 4 times ten out of the 5 tens, there is 1 ten left. This 1 ten we join to the 6 units, making 16. Then, 4 into 16 goes 4 times, which we set down, and the work is done.

This manner of performing the operation is called *Short Division*. The computation, it may be perceived, is carried on *partly* in the mind, which it is always easy to do when the divisor does not exceed 12.

### RULE.

*From the illustration of this example, we derive this general rule for dividing, when the divisor does not exceed 12 :*

I. Find how many times the divisor is contained in the first figure, or figures, of the dividend, and, setting it directly under the dividend, carry the remainder, if any, to the next figure as so many tens.

II. Find how many times the divisor is contained in *this* dividend, and set it down as before, continuing so to do till all the figures in the dividend are divided.

**PROOF.** We have seen, (¶ 15,) that the divisor and quotient are factors, whose product is the dividend, and we have also seen, that dividing the dividend by *one* factor is merely a process for finding the *other*.

Hence *division* and *multiplication* mutually prove each other.

*To prove division*, we may *multiply* the divisor by the quotient, and, if the work be right, the product will be the same as the dividend ; or we may divide *the dividend* by the *quotient*, and, if the work is right, the *result* will be the same as the divisor.

*To prove multiplication*, we may divide *the product* by *one factor*, and, if the work be right, the *quotient* will be the *other factor*.

### EXAMPLES FOR PRACTICE.

24. A man would divide 13,462,725 pounds among 5 men ; how many pounds would each receive ?

## OPERATION.

*Dividend.*  
 Divisor, 5)13,462,725  
 Quotient, 2,692,545

## PROOF.

*Quotient.*  
 2,692,545  
 5 divisor.

---

 13,462,725

its proof, it is plain, as before stated, that division is the reverse of multiplication, and that the two rules mutually prove each other.

25. How many yards of cloth can be bought for 4,354,560 shillings, at 2 shillings a yard? — at 3 shillings? — at 4 shillings? — at 5 shillings? — at 6 shillings? — at 7? — at 8? — at 9? — at 10?

*Note.* Let the pupil be required to prove the foregoing, and all of the following examples.

26. Divide 1005903360 by 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, and 12.

27. If 2 pints make a quart, how many quarts in 8 pints? — in 12 pints? — in 20 pints? — in 24 pints? — in 248 pints? — in 3674 pints? — in 47632 pints?

28. Four quarts make a gallon; how many gallons in 8 quarts? — in 12 quarts? — in 20 quarts? — in 36 quarts? — in 368 quarts? — in 4896 quarts? in 5436144 quarts?

29. A man gave 86 apples to 5 boys; how many apples would each boy receive?

*Dividend.*  
 Divisor, 5)86  
 Quotient, 17—1 *Remainder.*

each boy's share would be 17 apples; but there is 1 apple left.

¶ 17. 5)86

---

 17½

In order to divide *all* the apples equally among the boys, it is plain, we must divide this *one* remaining apple into 5

equal parts.  
 Then each  
 part of a

The 1  
 is, whole  
 ing parts  
 (that is

Fract  
 one dire  
 showing  
 lower.  
 called t  
 called t

*Note.*  
 and a f

In th  
 after ca  
 bers, is  
 dividen  
 this re  
 5 equal  
 of par  
 that it  
 the di

The  
 right  
 under

*Proof*

5 equ  
 apple  
 5 tim

30  
 ornu

equal parts, and give one of these parts to each of the boys. Then each boy's share would be 17 apples, and one fifth part of another apple; which is written thus,  $17\frac{1}{5}$  apples.

*Ans.*  $17\frac{1}{5}$  apples each.

The 17, expressing *whole* apples, are called *integers*, (that is, *whole* numbers.) The  $\frac{1}{5}$  (one fifth) of an apple, expressing *part* of a broken or divided apple, is called a *fraction*, (that is a *broken* number.)

Fractions as we here see, are written with two numbers, one directly over the other, with a short line between them, showing that the *upper* number is to be divided by the *lower*. The upper number, or *dividend*, is, in fractions, called the *numerator*, and the lower number, or *divisor*, is called the *denominator*.

*Note.* A number like  $17\frac{1}{5}$ , composed of integers (17) and a fraction, ( $\frac{1}{5}$ ), is called a *mixed number*.

In the preceding example, the one apple, which was left after carrying the division as far as could be by *whole* numbers, is called the *remainder*, and is evidently a part of the *dividend* yet undivided. In order to complete the division, this remainder, as we before remarked, must be divided into 5 equal parts; but the *divisor* itself expresses the number of parts. If, now, we examine the fraction, we shall see, that it consists of the remainder (1) for its *numerator*, and the divisor (5) for its *denominator*.

Therefore, if there be a *remainder*, set it down at the right hand of the quotient for the *numerator* of a fraction, under which write the divisor for its *denominator*.

*Proof the last example.*

$$\begin{array}{r} 17\frac{1}{5} \\ 5 \\ \hline 86 \end{array}$$

In proving this example, we find it necessary to multiply our fraction by 5; but this is easily done, if we consider, that the fraction  $\frac{1}{5}$  expresses one part of an apple divided into

5 equal parts; hence, 5 times  $\frac{1}{5}$  is  $\frac{5}{5}=1$ , that is, one *whole* apple, which we reserve to be added to the *units*, saying, 5 times 7 are 35, and one we reserved makes 36, &c.

30. Eight men drew a bounty of 453 pounds from government, how many pounds did each receive?

*Dividend.*

Divisor, 8 ) 453

Quotient,  $56\frac{5}{8}$

Here, after carrying the division as far as possible by *whole* numbers, we have a remainder of 5 pounds, which, written as above directed, gives for the answer 56 pounds and  $\frac{5}{8}$  (five eighths) of another pound, to each man.

¶ 18. Here we may notice, that the eighth part of 5 pounds is the same as 5 times the eighth part of 1 pound, that is, the eighth part of 5 pounds is  $\frac{5}{8}$  of a pound. Hence,  $\frac{5}{8}$  expresses the quotient of 5 divided by 8.

*Proof.*  $\frac{5}{8}$  is 5 parts; and 8 times 5 is 40, that is,  $40 = 5$ , which, reserved and added to the product of 8 times 6, makes 53, &c. Hence, to multiply a fraction, we may multiply the *numerator*, and divide the product by the *denominator*.

$56\frac{5}{8}$   
 $\times 8$   


---

453

Or, in proving division, we may multiply the *whole* number in the quotient *only*, and to the product *add* the remainder; and this, till the pupil shall be more particularly taught in fractions, will be more easy in practice. Thus,  $56 \times 8 = 448$ , and  $448 + 5$ , the remainder,  $= 453$ , as before.

31. There are 7 days in a week; how many weeks in 365 days? *Ans.* 52 $\frac{1}{4}$  weeks.

32. When flour is worth 2 pounds a barrel, how many barrels may be bought for 25 pounds? how many for 51 pounds? — for 487 pounds? — for 7631 pounds?

33. Divide 640 pounds among 4 men.

$640 \div 4$ , or  $64^0 = 160$  pounds, *Ans.*

34.  $678 \div 6$  or  $67\frac{8}{6} =$  how many? *Ans.* 113.

35.  $50\frac{4}{5}^0 =$  how many?

36.  $72\frac{3}{4}^4 =$  how many?

37.  $34\frac{6}{9}^4 =$  how many? *Ans.* 384 $\frac{2}{3}$ .

38.  $27\frac{6}{11}^4 =$  how many?

39.  $40\frac{2}{3}^0 =$  how many?

40.  $20\frac{14}{12}^0 =$  how many?

¶ 19. 41. Divide 4370 pounds equally among 21 men.

When, as in this example, the divisor exceeds 12, it is evident that the computation cannot be readily carried on in the mind, as in the foregoing examples. Wherefore, it is more convenient to write down the computation *at length*, in the following manner:

¶ 19.

OPE  
Divisor, 21 )

parts, we  
finding it  
of the div  
43 being  
This, how  
by tens a  
dividend,  
be in the

It is p  
to be tak  
divisor (2  
making 4  
dividend,  
bring dov

We th  
tens;) 1  
quotient,  
170. W  
and, find  
multiply  
168, unc  
der to be  
of the q  
for the a

This i  
Division

From

I. P  
them b  
dividen  
II.

OPERATION.

Divisor, Dividend, Quotient.

$$21 \overline{) 4370} \left( 208 \frac{2}{1} \right.$$

42

170

168

2 Remainder.

We may write the divisor and dividend as in short division, but, instead of writing the quotient *under* the dividend, it will be found more convenient to set it to the *right hand*.

Taking the dividend by parts, we seek how often we can have 21 in 43 (hundreds;) finding it to be 2 times, we set down 2 on the right hand of the dividend for the highest figure in the quotient. The 43 being *hundreds*, it follows, that the 2 must be hundreds. This, however, we need not regard, for it is to be followed by *tens* and *units*, obtained from the tens and units of the dividend, and will therefore, at the end of the operation, be in the place of hundreds, as it should be.

It is plain that 2 (hundred) times 21 pounds ought now to be taken out of the dividend; therefore, we multiply the divisor (21) by the quotient figure 2 (hundred) now found, making 42, (hundred,) which, written under the 43 in the dividend, we subtract, and to the remainder, 1, (hundred,) bring down the 7, (tens,) making 17 tens.

We then seek how often the divisor is contained in 17, (tens;) finding that it will not go, we write a cipher in the quotient, and bring down the next figure, making the whole 170. We then seek how often 21 can be contained in 170, and, finding it to be 8 times, we write 8 in the quotient, and, multiplying the divisor by this number, we set the product, 168, under the 170; then subtracting, we find the remainder to be 2, which, written as a fraction on the right hand of the quotient, as already explained, gives  $208 \frac{2}{1}$  pounds, for the *answer*.

This manner of performing the operation is called *Long Division*. It consists in writing down the *whole* computation.

From the above example, we derive the following

RULE.

I. Place the divisor on the left of the dividend, separate them by a line, and draw another line on the right of the dividend to separate it from the quotient.

II. Take as many figures, on the left of the dividend, as



contain the divisor once or more ; seek how many times they contain it, and place the answer on the right hand of the dividend for the first figure in the quotient.

III. Multiply the divisor by this quotient figure, and write the product under that part of the dividend taken.

IV. Subtract the product from the figures above, and to the remainder bring down the next figure in the dividend, and divide the number it makes up, as before. So continue to do, till all the figures in the dividend shall have been brought down and divided.

*Note 1.* Having brought down a figure to the remainder, if the number it makes up be *less* than the divisor, write a cipher in the quotient, and bring down the next figure.

*Note 2.* If the product of the divisor, by any quotient figure, be *greater* than the part of the dividend taken, it is an evidence that the quotient figure is *too large*, and must be diminished. If the remainder at any time be *greater* than the divisor, or equal to it, the quotient figure is *too small*, and must be increased.

#### EXAMPLES FOR PRACTICE.

1. How many hogshheads of molasses, at 7 pounds a hogsh-head, may be bought for 6318 pounds ?

*Ans.* 902 $\frac{1}{2}$  hogshheads.

2. If a man's income be 1248 pounds a year, how much is that per week, there being 52 weeks in a year ?

*Ans.* 24 pounds per week.

3. What will be the quotient of 153598, divided by 29 ?

*Ans.* 5296 $\frac{1}{2}$ .

4. How many times is 63 contained in 30131 ?

*Ans.* 478 $\frac{1}{3}$  times ; that is, 478 times, and  $\frac{1}{3}$  of another time.

5. What will be the several quotients of 7652, divided by 16, 23, 34, 86, and 92 ?

6. If a farm, containing 256 acres, be worth 1850 pounds, what is that per acre ?

7. What will be the quotient of 974932, divided by 365 ?

*Ans.* 2671 $\frac{1}{5}$ .

8. Divide 3228242 pounds equally among 563 men ; how many pounds must each man receive ? *Ans.* 5734 pounds.

9. If 57624 be divided into 216,586, and 976 equal parts, what will be the magnitude of one of each of these equal parts ?

*Ans.* T  
parts will  
10. Ho

11. T  
said to tr  
there bei  
12. 12  
13. 40  
14. 98

I.  
† 20.  
much w  
15 ya  
the cost  
are 3 ti  
but one

Henc  
if we p  
parts, a  
other ;

2. If  
will it  
It wi  
O  
24=6

3. I  
4. I  
5. I  
6. I

† 2  
what

times they  
and of the

figure, and  
d taken.

ve, and to  
e dividend,  
So continue  
have been

remainder,  
or, write a  
figure.

by quotient  
taken, it is  
, and must  
be greater  
figure is too

nds a hogs-

hogsheads.  
how much  
r?

per week.  
led by 29?  
s. 5296 $\frac{1}{4}$ .

of another

divided by

0 pounds,

l by 365?

2671 $\frac{17}{355}$ .

men; how

4 pounds.

76 equal

a of these

*Ans.* The magnitude of one of the last of these equal parts will be 59 $\frac{4}{5}$ .

10. How many times does 1030603615 contain 3215?

*Ans.* 320561 times.

11. The earth in its annual revolution round the sun, is said to travel 596088000 miles; what is that per hour, there being 8766 hours in a year?

12.  $1234567890 =$  how many?

13.  $40793920 =$  how many?

14.  $987649031 =$  how many?

## CONTRACTIONS IN DIVISION.

I. *When the DIVISOR is a COMPOSITE NUMBER.*

¶ 20. 1. Bought 15 yards of cloth for 30 pounds; how much was that per yard?

15 yards are  $3 \times 5$  yards. If there had been but 5 yards, the cost of one yard would be  $3\text{ }^s = 6$  pounds; but as there are 3 times 5 yards, the cost of one yard will evidently be but one third part of 6 pounds; that is,  $2\text{ }^s = 2$  pounds, *Ans.*

Hence, when the divisor is a composite number, we may, if we please, divide the dividend by one of the component parts, and the quotient, arising from that division, by the other; the last quotient will be the answer.

2. If a man can travel 24 miles in a day, how many days will it take him to travel 264 miles?

It will evidently take him as many days as 264 contains 24.

OPERATION.

$24 = 6 \times 4.$        $6)264$

—  
4)44

11 days.

or,

$24)264(11 \text{ days, } \textit{Ans.}$

24

—  
24

—  
24

3. Divide 576 by 48 =  $(8 \times 6.)$

4. Divide 1260 by 63 =  $(7 \times 9.)$

5. Divide 2430 by 81.

6. Divide 448 by 56.

II. *To divide by 10, 100, 1000, &c.*

¶ 21. 1. A note of 2478 pounds is owned by 10 men; what is each man's share?

E

Each man's share will be equal to the number of *tens* contained in the whole sum, and, if one of the figures be cut off at the right hand, all the figures to the left may be considered so many *tens*; therefore, each man's share will be 247  $\frac{1}{10}$  pounds.

It is evident, also, that if 2 figures had been cut off from the right, all the remaining figures would have been so many *hundreds*; if 3 figures, so many *thousands*, &c. Hence, we derive this general RULE for dividing by 10, 100, 1000, &c.: Cut off from the right of the dividend so many figures as there are ciphers in the divisor; the figures to the left of the point will express the *quotient*, and those to the right, the remainder.

2. How many 100 in 42400 ? *Ans.* 424.

424|00 Here the divisor is 100; we therefore cut off 2 figures on the right hand, and all the figures to the left (424) express the number of hundreds.

3. How many 100 in 34567 ? *Ans.* 345  $\frac{67}{100}$ .

4. How many hundreds in 4567840 hundreds ?

5. How many hundreds in 345600 hundreds ?

6. How many 100 in 42604 hundreds ? *Ans.* 426  $\frac{4}{100}$ .

7. How many thousands in 4000 ? — in 25000 ?

8. How many thousands in 6487 thousands ? *Ans.* 6  $\frac{487}{1000}$ .

9. How many thousands in 42863 thousands ? — in 368456 thousands ? — in 96842378 thousands ?

10. How many tens in 40 ? in 400 ? in 20 ? in 468 ? in 4784 ? in 34640 ?

III. When there are CIPHERS on the right hand of the divisor.

T 22. 1. Divide 480 pounds among 40 men ?

OPERATION.

4|0)480

12 pounds, *Ans.*

In this example, our divisor, (40,) is a composite number, ( $10 \times 4 = 40$ ;) we may therefore, divide by *one* component part, (10,) and that quotient by the *other*, (4;) but to divide by 10, we have seen, is but to cut off the right hand figure, leaving the figures to the left of the point for the quotient, which we divide by 4, and the work is done. It is evident, that, if our divisor had been 400, we should have cut off 2 figures, and have divided in the same manner; if 4000, 3 figures, &c. Hence, this general RULE: When there are ciphers at the right hand

of the dividend;  
the remainder  
cut off from

2. Divi

Divisor

3. Divi

4. How  
pounds, &

5. Div

6. Div

7. Div  
50, 600,

1. What  
is it? 3.  
be divided  
What is t  
&c.? 6.  
sign of di  
expressing  
formed?  
proved?  
fractions,  
When the  
how is it  
is the up  
How do  
the denomi  
division?  
how may  
how may  
at the rig

of the divisor, cut them off, and also as many places in the dividend; divide the remaining figures in the dividend by the remaining figures in the divisor; then annex the figures cut off from the dividend, to the remainder.

2. Divide 748346 by 8000.

*Dividend.*

*Divisor*, 8|000748|346.

*Quotient*, 93.—4346 *Remainder*.

*Ans.* 934346

3. Divide 46720367 by 4200000.

*Dividend.*

42|00000467|20367(11.429367 *Quotient.*

42

47

42

520367 *Remainder.*

4. How many pieces of cloth can be bought for 346500 pounds, at 20 pounds per piece?

5. Divide 76428400 by 900000.

6. Divide 345006000 by 84000.

7. Divide 4680000 by 20, 200, 2000, 20000, 3000, 4000, 50, 600, 70000, and 80.

## SUPPLEMENT TO DIVISION.

### QUESTIONS.

1. What is division? 2. In what does the *process* of division consist? 3. Division is the *reverse* of what? 4. What is the *number to be divided* called, and to what does it answer in multiplication? 5. What is the *number to divide by* called, and to what does it answer, &c.? 6. What is the *result* or *answer* called, &c.? 7. What is the *sign* of division, and what does it show? 8. What is the *other way* of expressing division? 9. What is *short division*, and how is it performed? 10. How is division *proved*? 11. How is *multiplication* proved? 12. What are *integers*, or whole numbers? 13. What are *fractions*, or broken numbers? 14. What is a *mixed number*? 15. When there is any thing *left* after division, what is it called, and how is it to be written? 16. How are *fractions written*? 17. What is the upper number called? 18. — the lower number? 19. How do you multiply a fraction? 20. To what do the numerator and the denominator of a fraction answer in division? 21. What is *long division*? 22. Rule? 23. When the divisor is a composite number, how may we proceed? 24. When the divisor is 10, 100, or 1000, &c. how may the operation be contracted? 25. When there are ciphers at the right hand of the divisor how may we proceed?

## EXERCISES.

1. An army of 1500 men, having plundered a city, took 2625000 pounds; what was each man's share?

2. A certain number of men were concerned in the payment of 18950 pounds, and each man paid 25 pounds; what was the number of men?

3. If 7412 eggs be packed in 34 baskets, how many in a basket?

4. What number must I multiply by 135 that the product may be 505710?

5. Light moves with such amazing rapidity, as to pass from the sun to the earth in about the space of 8 minutes. Admitting the distance, as usually computed to be 95000000 miles, at what rate per minute does it travel?

6. If the product of two numbers be 704, and the multiplier be 11, what is the multiplicand? *Ans. 64.*

7. If the product be 704, and the multiplicand 64, what is the multiplier? *Ans. 11.*

8. The divisor is 18, and the dividend 144; what is the quotient?

9. The quotient of two numbers is 8, and the dividend 144; what is the divisor?

10. A man wishes to travel 585 miles in 13 days; how many miles must he travel each day?

11. If a man travels 45 miles a day, in how many days will he travel 585 miles?

12. A man sold 140 cows for 560 pounds; how much was that for each cow?

13. A man, selling his cows for 4 pounds each, received for all 560 pounds; how many cows did he sell?

14. If 12 inches make a foot, how many feet are there in 364812 inches?

15. If 364812 inches are 30401 feet, how many inches make 1 foot?

17. If you would divide 48750 pounds among 50 men, how many pounds would you give to each one?

16. If you distribute 48750 pounds among a number of men, in such a manner as to give to each one 975 pounds, how many men receive a share?

18. A man has 17484 pounds of tea in 186 chests; how many pounds in each chest?

a city, took

in the pay-  
ments; what

many in a

the product

as to pass  
8 minutes.  
95000000

the multi-

*Ans.* 64.

64, what

*Ans.* 11.

what is the

the dividend

days; how

many days

how much

, received

re there in

ny inches

50 men,

number of

5 pounds,

sts; how

19. A man would put up 17484 pounds of tea into chests containing 94 pounds each; how many chests must he have?

20. In a certain town there are 1740 inhabitants, and 12 persons in each house; how many houses are there? — in each house are 2 families, how many persons in each family?

21. If 2760 men can dig a certain canal in one day, how many days would it take 46 men to do the same? How many men would it take to do the work in 15 days? — in 5 days? — in 20 days? — 40 days? — in 120 days?

22. If a carriage wheel turns round 62280 times in running from Quebec to Montreal, a distance of 180 miles, how many times does it turn in running 1 mile? *Ans.* 346.

23. Sixty seconds make 1 minute; how many minutes in 3600 seconds? — in 86400 seconds? — in 604800 seconds? — in 2419200 seconds?

24. Sixty minutes make one hour; how many hours in 1440 minutes? — in 10080 minutes? — in 40320 minutes? — in 525960 minutes?

25. Twenty-four hours make a day; how many days in 168 hours? — in 672 hours? — in 3766 hours?

26. How many times can I subtract forty-eight from four hundred and eighty?

27. How many times 3478 is equal to 47854?

28. A bushel of grain is 32 quarts; how many quarts must I dip out of a chest of grain to make one half ( $\frac{1}{2}$ ) of a bushel? — for one fourth ( $\frac{1}{4}$ ) of a bushel? — for one eighth ( $\frac{1}{8}$ ) of a bushel? *Ans.* to the last 4 quarts.

29. How many is  $\frac{1}{2}$  of 20? —  $\frac{1}{2}$  of 48? —  $\frac{1}{2}$  of 247? —  $\frac{1}{2}$  of 847? —  $\frac{1}{2}$  of 345878? —  $\frac{1}{2}$  of 204030648? *Ans.* to the last, 102015324.

30. How many walnuts are one third part ( $\frac{1}{3}$ ) of 3 walnuts? —  $\frac{1}{3}$  of 6 walnuts? —  $\frac{1}{3}$  of 12 walnuts? —  $\frac{1}{3}$  of 30? —  $\frac{1}{3}$  of 45? —  $\frac{1}{3}$  of 300? — of 478? —  $\frac{1}{3}$  of 3456320? *Ans.* to the last, 11521066.

31. What is  $\frac{1}{4}$  of 4? —  $\frac{1}{4}$  of 20? —  $\frac{1}{4}$  of 320? —  $\frac{1}{4}$  of 7843? *Ans.* to the last, 1960.

### MISCELLANEOUS QUESTIONS.

*Involving the principles of the preceding rules.*

*Note.* The preceding rules, viz. Numeration, Addition.

Subtraction, Multiplication, and Division, are called the *Fundamental Rules of Arithmetic*, because they are the foundation of all other rules.

1. A man bought a chaise for 57 pounds, and a horse for 34 pounds; what did they both cost?

2. If a horse and chaise cost 91 pounds, and the chaise cost 57 pounds, what is the cost of the horse? If the horse cost 24 pounds, what is the cost of the chaise?

3. If the sum of 2 numbers be 487, and the greater number be 348, what is the less number? If the less number be 139, what is the greater number?

4. If the minuend be 7842, and the subtrahend 3481, what is the remainder? If the remainder be 4361, and the minuend be 7842, what is the subtrahend?

¶ 23. When the minuend and the subtrahend are given, how do you find the remainder?

When the minuend and remainder are given, how do you find the subtrahend?

When the subtrahend and the remainder are given, how do you find the minuend?

When you have the *sum* of two numbers, and *one* of them given, how do you find the other?

When you have the *greater* of two numbers, and their *difference* given, how do you find the less number?

When you have the *less* of two numbers, and their *difference* given, how do you find the *greater* number?

5. The sum of two numbers is 48, and *one* of the numbers is 19; what is the other?

6. The *greater* of two numbers is 29, and their *difference* 19; what is the less number?

7. The *less* of two numbers is 19, and their *difference* is 10; what is the *greater*?

8. A man bought 5 pieces of cloth at 44 pounds a piece; 974 dozen of shoes, at 3 pounds a dozen; 600 pieces of calico, at 6 pounds a piece; what is the amount?

9. A man sold six cows at 5 pounds each, and a yoke of oxen, for 19 pounds; in pay, he received a chaise, worth 31 pounds, and the rest in money; how much money did he receive?

10. What will be the cost of 15 pounds of butter, at 7 pence per pound?

11.  
shilling

¶ 2  
any co  
number

the pri  
the wh

Wh  
&c. is  
bushel

Wh  
also th  
ber of

12.  
cost o

13.  
pence

Wh  
Wh

find t  
Wh  
find t

Wh  
find t  
14.  
15.  
16.  
17.

he sa  
How

pence  
18

ten c  
ty-fo  
four  
appl

L  
vers

19  
and  
and  
is th



11. How many bushels of wheat can you buy for 4870 shillings, at 8 shillings per bushel?

¶ 24. When the price of *one* pound, *one* bushel, &c. of any commodity is given, how do you find the cost of any number of pounds, or bushels, &c. of that commodity? If the price of the 1 pound, &c. be in shillings, in what will the whole cost be? If in pence, what?

When the cost of any given number of pounds, or bushels, &c. is given, how do you find the price of *one* pound or bushel, &c. In what kind of money will the answer be?

When the cost of a number of pounds, &c. is given, and also the price of *one* pound, &c. how do you find the number of pounds, &c.

12. When rye is 4 shillings per bushel, what will be the cost of 212 bushels?

13. If 18 pounds of tea cost 173 pounds, (that is 41520 pence) what is the price of one pound?

When the factors are given, how do you find the product?

When the product and one factor are given, how do you find the other factor?

When the divisor and quotient are given, how do you find the dividend?

When the dividend and quotient are given, how do you find the divisor?

14. What is the product of 754 and 25?

15. What number multiplied by 25, will produce 18850?

16. What number, multiplied by 754, will produce 18850?

17. If a man save 5 pence a day, how many pence would he save in a year, (365 days,)? — how many in 45 years? How many cows could he buy with the money, at 742 pence each?

18. A boy bought a number of apples; he gave away ten of them to his companions, and afterwards bought thirty-four more, and divided half of what he then had among four companions, who received 8 apples each; how many apples did the boy first buy?

Let the pupil take the last number of apples, 8, and reverse the process.


*Ans.* 40 apples.

19. There is a certain number, to which, if 4 be added, and 7 be subtracted, and the difference be multiplied by 8, and the product divided by 3, the quotient will be 64; what is that number?

*Ans.* 27.

20. A board has 8 rows of 8 squares each; how many squares on the board?

25. 21. There is a spot of ground 5 rods long, and 3 rods wide; how many square rods does it contain?



rows is equal to the number of rods, on the other side; therefore,  $5 \times 3 = 15$ , the number of squares. *Ans.* 15 square rods.

A figure, like A, B, C, D, having its opposite sides equal and parallel, is called a *parallelogram* or *oblong*.

22. There is an oblong field, 40 rods long, and 24 rods wide; how many square rods does it contain?

23. How many square inches in a board 12 inches long, and 12 inches broad? *Ans.* 144.

24. How many square feet in a board 14 feet long and 2 feet wide?

25. A certain township is six miles square; how many square miles does it contain? *Ans.* 36.

26. A man bought a lot of land for 2246 pounds; he sold one half of it for 1175 pounds at the rate of 3 pounds per acre; how many acres did he buy? and what did it cost him per acre?

27. A boy bought a sled for 56 pence, and sold it again for 8 quarts of walnuts; he sold one half of the nuts at 8 pence a quart, and gave the rest for a penknife, which he sold for 18 pence; how many pence did he lose by his bargains?

28. In a certain school-house, there are 5 rows of desks; in each row are six seats, and each seat will accommodate 2 pupils; there are also two rows, of 3 seats each, of the same size as the others, and one long seat where 8 pupils may sit; how many scholars will this house accommodate? *Ans.* 80.

29. How many square feet of boards will it take for the

*Note.* A square rod is a square (like one of those in the annexed figure) measuring a rod on each side. By an inspection of the figure, it will be seen, that there are as many squares in a row as rods on one side, and that the number of

floor of  
12 square

30. T  
many y  
cover th  
yards?

31. A  
is its br

32. F  
can be b

33. F  
must be

34. 4  
35. 4

36. A  
men; l

is  $\frac{1}{2}$  of

37. 1  
— of

38. 1  
— of

39. 1  
60?

246876  
40.

41.  
feet in

— in  
42.

cost?  
— 1

43.  
can be

44.  
rate of

in one  
days?

this la  
years

45.  
35 pe

46.

floor of a room 16 feet long, and 15 feet wide, if we allow 12 square feet for waste ?

30. There is a room 6 yards long and 5 yards wide ; how many yards of carpeting, a yard wide, will be sufficient to cover the floors, if the hearth and fireplace occupy 3 square yards ?

31. A board, 14 feet long, contains 28 square feet ; what is its breadth ?

32. How many pounds of pork, worth 4 pence a pound, can be bought for 144 pence ?

33. How many pounds of butter, at 9 pence per pound, must be paid for 25 pounds of tea, at 38 pence per pound ?

34.  $4+5+6+1+8=$  how many ?

35.  $4+3+10-2-4+6-7=$  how many ?

36. A man divides 30 bushels of potatoes among 3 poor men ; how many bushels does each man receive ? What is  $\frac{1}{3}$  of thirty ? How many are  $\frac{2}{3}$  (two thirds) of 30 ?

37. How many are one third ( $\frac{1}{3}$ ) of 3 ? — of 6 ? — of 9 ? — of 282 ? — of 45674312 ?

38. How many are two thirds ( $\frac{2}{3}$ ) of 3 ? — of 6 ? — of 9 ? — of 282 ? — of 45674312 ?

39. How many are  $\frac{1}{4}$  of 40 ? —  $\frac{3}{4}$  of 40 ? —  $\frac{1}{2}$  of 60 ? —  $\frac{3}{4}$  of 60 ? —  $\frac{1}{4}$  of 80 ? — of 124 ? — of 246876 ? —  $\frac{3}{4}$  of 246876 ?

40. How many is  $\frac{1}{2}$  of 80 ? —  $\frac{3}{4}$  of 80 ? —  $\frac{3}{4}$  of 100 ?

41. An inch is one twelfth part ( $\frac{1}{12}$ ) of a foot ; how many feet in 12 inches ? — in 24 inches ? — in 36 inches ? — in 12243648 inches ?

42. If 4 pounds of tea cost 128 pence, what does 1 pound cost ? — 2 pounds ? — 3 pounds ? — 5 pounds ? — 100 pounds ?

43. When oranges are worth 4 pence apiece, how many can be bought for 1464 pence ?

44. The earth, in moving round the sun, travels at the rate of 68000 miles an hour ; how many miles does it travel in one day, (24 hours ?) how many miles in one year, (365 days ?) and how many days would it take a man to travel this last distance, at the rate of 40 miles a day ? how many years ?

*Ans. to the last, 40800.*

45. How many pence can a man earn in 20 weeks, at 35 pence per day, Sundays excepted ?

46. A man married at the age of 23 ; he lived with his

wife 14 years; she then died, leaving him a daughter, 12 years of age; 8 years after the daughter was married to a man 5 years older than herself, who was 40 years of age when the father died; how old was the father at his death?

*Ans.* 60.

47. There is a field 20 rods long, and 8 rods wide; how many square rods does it contain? *Ans.* 160 rods.

48. What is the width of a field, which is 20 rods long, and contains 160 square rods.

49. What is the length of a field, 8 rods wide, and containing 160 square rods?

50. What is the width of a piece of land, 25 rods long, and containing 400 square rods?

## COMPOUND NUMBERS.

¶ 26. A number expressing things of the same kind is called a *simple number*; thus, 100 men, 56 years, 75 cents, are each of them simple numbers; but when a number expresses things of different kinds, it is called a *compound number*; thus, 46 pounds 7 shillings and 6 pence, is a compound number; so 4 years 6 months and 3 days, 43 dollars 25 cents and 3 mills, are compound numbers.

*Note.* Different kinds, or names, are usually called *different denominations*.

## REDUCTION.

¶ 27. In this Province as in England, money is reckoned in pounds, shillings, pence and farthings. In the United States, money is reckoned in dollars, cents and mills. These are called denominations of money. Time is reckoned in years, months, weeks, days, hours, minutes, and seconds, called denominations of time. Distance is reckoned in miles, rods, feet, and inches, called denominations of measure, &c.

The relative value of these denominations is exhibited in tables, which the pupil must commit to memory.

¶ 27

TI  
curre  
its ce  
Scoti  
denom  
pence  
nomin  
ney,  
perny  
fourth  
be see  
ratio d  
exhibi  
for pa

12

20

*Note.*

penny;  
farthing

It wi  
ver coin  
Canada,

DENO

A Guin  
A Hal  
A thir  
A Joh  
A half  
A Moie  
An Ea  
A half

# HALIFAX CURRENCY.

The present currency of Lower Canada, is called Halifax currency, having been introduced into this Province, after its cession to Great Britain, by France, in 1763, from Nova Scotia. The denominations are the same in name as the denominations of English money, i. e. pounds, shillings, pence, and farthings; and the ratios of the different denominations to each other are the same as in English money, i. e. the shilling is one twentieth of the pound, the penny one twelfth of the shilling, and the farthing one fourth of the penny. In value they are different, as will be seen in the ¶ upon reduction of currencies; where the ratio of each to the other, and of both to Federal Money is exhibited, with the method of ascertaining them in practice, for particular sums.

2 farthings (qrs.)	make	1 half-penny,	marked	½d.
4 "	"	"	1 penny,	" d.
12 pence	"	"	1 shilling,	" s.
20 shillings	"	"	1 pound,	" £.

*Note.* Farthings are often written as the fraction of a penny; thus, 1 farthing is written ¼d., 2 farthings, ½d., 3 farthings, ¾d.

It will be proper to insert here a list of the gold and silver coins, made current by law in the Province of Lower Canada, with their several weights and values annexed.

## GOLD COINS.

*English, American, and Portuguese.*

DENOMINATION.	WEIGHT.		VALUE.		
	poles.	grs.	£.	s.	d.
A Guinea,	5	6	1	3	4
A Half Guinea,	2	15	0	11	8
A third Guinea,	1	18	0	7	9
A Johannes,	18	0	4	0	0
A half Johannes,	9	0	2	0	0
A Moidore,	6	18	1	10	0
An Eagle,	11	6	2	10	0
A half Eagle,	5	15	1	5	0

## GOLD COINS—CONTINUED.

*French and Spanish.*

DENOMINATION.	WEIGHT.		VALUE.		
	<i>pnls.</i>	<i>grs.</i>	£	s.	d.
A Doubloon, - -	17	0	3	14	6
A half Doubloon, -	8	12	1	17	3
A Louis d'Or, coined before 1793.	5	4	1	2	8
A Pistole, coined before 1793.	4	4	0	18	3
The 40 Francs piece, coined } before 1792.	8	6	1	16	2
The 40 Francs piece, coined } since 1792.	8	6	1	16	2
The 20 Francs piece, coined } before 1792.	4	3	0	18	1
The 20 Francs piece, coined } since 1792.	4	3	0	18	1

## SILVER COINS.

A Crown, - - - - -	0	5	6
An English shilling, - - - - -	0	1	1
A Dollar, - - - - -	0	5	0
A Pistareen, - - - - -	0	0	10
A half Pistareen, - - - - -	0	0	5
The American Dollar, - - - - -	0	5	0
A French piece of 4 livres 10 sols Tournois,	0	4	2
A French piece of 6 livres, - - -	0	5	6
The piece of 5 Francs Tournois, coined } since 1792,	0	4	8
The French piece of 36 sols Tournois,	0	1	8
The French piece of 24 sols Tournois,	0	1	1

The statute provides, that "all the higher and lower denominations of the above gold and silver coins, shall also pass current, and be deemed a legal tender in payment of all debts, and demands whatsoever, in this Province, in the same proportions, respectively."

*Two pence farthing* is allowed for every grain over, or under the legal weight, on English, American, and Portuguese gold; and *two pence and one fifth*, for every grain over or under the legal weight, on French and Spanish gold.

Payments in gold above *twenty pounds* shall be made in bulk, where one of the parties making or receiving the

VALUE.

	s.	d.
3	14	6
1	17	3
1	2	8
0	18	3
1	16	2
1	16	2
0	18	1
0	18	1
5	6	
1	1	
5	0	
0	10	
0	5	
5	0	
4	2	
5	6	
4	8	
1	8	
1	1	

and lower de-  
s, shall also  
payment of  
vince, in the  
ain over, or  
, and Portu-  
y grain over  
ish gold.  
be made in  
ceiving the

same shall require it; English, American, and Portuguese gold at *eighty-nine shillings per ounce Troy*; French, and Spanish gold at *eighty-seven shillings and eight pence half-penny per ounce Troy*:—and a deduction shall be made of  $\frac{1}{4}$  grain Troy, for each piece of gold coin so weighed, as a compensation to the receiver for the loss that may accrue in afterwards paying away the same by the single piece.

In copper coin no person is obliged to receive more than one shilling at one payment.

How many farthings in 1 penny? — in 2 pence? — in 3 pence? — in 4 pence? — in 8 pence? — in 12 pence? — in 1 shilling? — in 2 shillings?

How many pence in 2 shillings? — in 3 s. ? — in 4 s. ? — in 6 s. ? — in 8 s. ? — in 10 s. ? — in 12 s. ? — in 2s. 3d. ? — in 2s. 4d. ? — in 4s. 3d. ?

How many shillings in 1 pound? — in 2£. ? — in 3£. ? — in 4£. ? — in 4£. 6s. ? — in 6£. 8s. ? — in 3£. 10s. ? — in 2£. 15s. ?

How many pence in 4 farthings? — in 8 farthings? — in 12 farthings? — in 24 farthings? — in 32 farthings? — in 36 farthings? — in 48 qrs. ? How many shillings in 48 qrs. ? — in 96 qrs. ?

How many shillings in 24 pence? — in 36d. ? — in 48d. ? — in 72d. ? — in 96d. ? — in 120d. ? — in 26d. ? — in 27d. ? — in 28d. ? — in 30d. ? — in 42d. ? — in 51d. ?

How many pounds in 20 shillings? — in 40s. ? — in 60s. ? — in 80s. ? — in 86s. ? — in 128s. ? — in 70s. ? — in 55s. ?

The changing of *one* kind, or denomination, into *another* kind, or denomination, without altering their value, is called *Reduction*. (¶ 27.) Thus, when we change shillings into pounds, or pounds into shillings, we are said to *reduce* them. From the foregoing examples, it is evident, that, when we reduce a denomination of *greater* value into a denomination of *less* value, the reduction is performed by *multiplication*; and it is then called *Reduction Descending*. But when we reduce a denomination of *less* value into one of *greater* value, the reduction is performed by *division*; it is then called *Reduction Ascending*. Thus, to reduce pounds



to shillings, it is plain, we must *multiply* by 20. And again, to reduce shillings to pounds, we must *divide* by 20. It follows, therefore, that *reduction descending and ascending reciprocally prove each other.*

1. In 17£. 13s. 6½d. how many farthings?

OPERATION.

£. s. d. qrs.

17 13 6 3

20s.

353s. in 17£. 13s.

12d.

4242d.

4q.

16971qrs. the Ans.

In the above example, because 20 shillings make 1 pound, therefore we multiply 17£. by 20, increasing the product by the addition of the given shillings, (13,) which, it is evident, must always be done in like cases; then, because 12 pence make 1 shilling, we multiply the shillings (353) by 12, adding in the given pence, (6.) Lastly, because 4 farthings make 1 penny, we multiply the pence (4242) by 4, adding in the given farthings, (3.) We then find, that in 17£. 13s. 6½d., are contained 16971 farthings.

2. In 16971 farthings, how many pounds?

OPERATION.

Farthings in a penny, 4)16971 3qrs.

Pence in a shilling, 12)4242 6d.

Shillings in a pound, 20)353 13s.

17£.

Ans. 17£. 13s. 6½d.

Farthings will be reduced to pence, if we divide them by 4, because every 4 farthings make 1 penny. Therefore, 16971 farthings divided by 4, the quotient is 4242 pence, and a remainder of 3, which is farthings, of the same name as the dividend. We then divide the pence (4242) by 12, reducing them to shillings; and the shillings (353) by 20, reducing them to pounds. The last quotient, 17£., with the several remainders, 13s. 6d. 3qrs. constitute the answer.

*Note.* In dividing 353s. by 20, we cut off the cipher, &c., as taught † 22.

† 28. The process in the foregoing examples, if carefully examined, will be found to be as follows, viz.

*To reduce high denominations to lower.*—Multiply the highest denomination by that num-

*To reduce low denominations to higher.*—Divide the lowest denomination given by that

ber w  
less to  
(incre  
numbe  
less d  
in the  
succee  
til you  
denom

In t  
are de  
and fa  
obviou  
where  
which  
all cor

3. 1

4.

5.

6.

7.

8.

9.

10.

to fart

12.

3s. 4d

14.

each,

16.

many

No

we m

lings

ber which it takes of the next number which it takes of the less to make 1 of this higher, same to make 1 of the next (increasing the product by the higher." Proceed in the same number given, if any, of that manner with each succeeding less denomination.) Proceed denomination, until you have in the same manner with each brought it to the denomina- succeeding denomination, un- tion required. til you have brought it to the denomination required.

In the two examples, from which the above general rules are deduced, the denominations are pounds, shillings, pence and farthings, considered as in Halifax Currency; but it is obvious that these rules can be applied to all currencies where the denominations are the same; or to currencies in which the denominations are different; and in general to all compound numbers.

### EXAMPLES FOR PRACTICE.

- |  |   |
|--|---|
| 3. Reduce 20£. 14s. 2d. to pence.                      | <i>Ans. 4970.</i>                         |
| 4. " 24£. to farthings.                                | <i>Ans. 23040.</i>                        |
| 5. " 66£. 6s. 6d. to pence.                            | <i>Ans. 15912.</i>                        |
| 6. " 158£. to farthings.                               | <i>Ans. 151680.</i>                       |
| 7. " 1234£. 15s. 7d. to farthings.                     | <i>Ans. 1185388.</i>                      |
| 8. " 337587 farthings to pounds, &c.                   | <i>Ans. 351£. 13s. 0d, 3q.</i>            |
| 9. " 1185388 farthings to pounds, &c.                  | <i>Ans. 1234£. 15s. 7d.</i>               |
| 10. Reduce 32£. 15s. 8d. to farthings.                 | 11. Reduce 31472 farthings to pounds.     |
| 12. In 29 guineas, at 1£. 3s. 4d. each, how many qrs.? | 13. In 38976 farthings, how many guineas? |
| 14. Reduce \$163, at 6s. each, to pence?               | 15. Reduce 11736 pence to dollars.        |
| 16. In 15 guineas, how many pounds?                    | 17. Reduce 21£. to guineas.               |

*Note.* We cannot reduce guineas *directly* to pounds, but we may reduce the guineas to *shillings*, and then the shillings to pounds.

## OLD CURRENCY.

12 deniers make 1 sou.  
20 sous " 1 livre, or franc.

The livre is 10d. Halifax currency.

In 32 livres 10 sous, how many sous ?

In 97 livres 11 sous, how many sous ?

In 650 sous, how many livres ?

In 1951 sous, how many livres ?

In 10 livres 6 sous 9 deniers, how many deniers ?

How many pounds currency in 96 livres ?

## FEDERAL MONEY.

† 29. Federal money is the coin of the United States. The kinds, or denominations, are eagles, dollars, dimes, cents, and mills.

## TABLE.

10 mills - - - are equal to - 1 cent.  
10 cents, (=100 mills,) - - - =1 dime.  
10 dimes, (=100 cents=1000 mills,) - =1 dollar.  
10 dollars, (=100 dimes=1000 cents=10000 mills) =1 eagle.\*

Sign. This character, \$, placed before a number, shows it to express *federal money*.

As 10 mills make a cent, 10 cents a dime, 10 dimes a dollar, &c. it is plain, that the relative value of mills, cents, dimes, dollars and eagles corresponds to the orders of units, tens, hundreds, &c. in simple numbers. Hence, they may be read either in the *lowest* denomination, or *partly* in a *higher*, and partly in the *lowest* denomination. Thus :

3 eagles.  
4 dollars.  
6 dimes.  
2 cents.  
2 mills.

\$34652 may be read, 34652 mills; or 3465 cents and 2 mills; or, reckoning the eagles *tens* of dollars, and the

\* The eagle is a gold coin, the dollar and dime are silver coins, the cent is a copper coin. The mill is only imaginary, there being no coin of that denomination. There are half eagles, half dollars, half dimes, and half cents, real coins.

Writ  
and 2 m  
mills;  
mill; 2  
cents and  
five cen

† The ch  
the comma  
comma us

dimes *ten* of cents, which is the usual practice, the whole may be read, 34 dollars 65 cents and 2 mills.

For ease in calculating, a point (·) called a *separatrix*,† is placed between the dollars and cents, showing that all the figures at the left hand express dollars, while the *two first figures* at the right hand express cents, and the *third*, mills. Thus, the above example is written \$34·652; that is, 34 dollars 65 cents 2 mills, as above. As 100 cents make a dollar, the *cents* may be any number from 1 to 99, often requiring *two* figures to express them; for this reason, *two* places are appropriated to cents, at the right hand of the point, and if the number of cents be less than *ten*, requiring but *one* figure to express them, the *ten's* place must be filled with a cipher. Thus, 2 dollars and 6 cents are written 2·06. 10 mills make a cent, and consequently the *mills* never exceed 9, and are always expressed by a *single* figure. Only *one* place, therefore, is appropriated to mills, that is, the place immediately following cents, or the *third* place from the point. When there are no cents to be written, it is evident that we must write *two* ciphers to fill up the places of cents. Thus, 2 dollars and 7 mills are written 2·007. Six cents are written, 06, and 7 mills are written '007.

*Note.* Sometimes 5 mills =  $\frac{1}{2}$  a cent is expressed fractionally: thus, '125 (twelve cents and five mills) is expressed 12 $\frac{1}{2}$ , (twelve and a half cents.)

17 dollars and 8 mills are written, 17·008

4 dollars 5 cents, - - - - - 4·05

75 cents, - - - - - '75

24 dollars, - - - - - 24·

9 cents, - - - - - '09

4 mills, - - - - - '004

6 dollars 1 cent and 3 mills, - 6·013

Write down 470 dollars 2 cents; 342 dollars 40 cents and 2 mills; 100 dollars, 1 cent and 4 mills; 1 mill; 2 mills; 3 mills; 4 mills;  $\frac{1}{2}$  cent, or 5 mills; 1 cent and 1 mill; 2 cents and 3 mills; six cents and one mill; sixty cents and one mill; four dollars and one cent; three cents; five cents; nine cents.

† The character used for the *separatrix*, in the "Scholars' Arithmetic," was the comma; the comma *inverted* is here adopted, to distinguish it from the comma used in punctuation

## REDUCTION OF FEDERAL MONEY.

¶ 29. How many mills in one cent? — in 2 cents? — in 3 cents? — in 4 cents? — in 6 cents? — in 9 cents? — in 10 cents? — in 30 cents? — in 78 cents? — in 100 cents, (=1 dollar)? — in 2 dollars? — in 3 dollars? — in 4 dollars? — in 484 cents? — in 563 cents? — in 1 cent and 2 mills? — in 4 cents and 5 mills?

How many cents in 2 dollars? — in 4 dollars? — in 8 dollars? — in 3 dollars and 15 cents? — in 5 dollars and 20 cents? — in 8 dollars and 20 cents? — in 4 dollars and 6 cents?

How many dollars in 400 cents? — in 600 cents? — in 380 cents? — in 40765 cents? How many cents in 1000 mills? How many dollars in 1000 mills? — in 3000 mills? — in 8000 mills? — in 4378 mills? — in 846732 mills?

As there are 10 mills in one cent, it is plain that cents are changed or reduced to mills by multiplying them by 10, that is, by merely annexing a cipher, (¶ 12.) 100 cents make a dollar; therefore dollars are changed to cents by annexing 2 ciphers, and to mills by annexing 3 ciphers. Thus, 16 dollars = 1600 cents = 16000 mills. Again, to change mills back to dollars, we have only to cut off the *three right hand figures*, (¶ 21;) and to change cents to dollars, cut off the *two right hand figures*, when all the figures to the *left* will be dollars, and the figures to the *right*, cents and mills.

Reduce 34 dollars to cents.

*Ans.* 3400.

Reduce 240 dollars and 14 cents to cents.

*Ans.* 24014 cents.

Reduce \$748'143 to mills.

*Ans.* 748143 mills.

Reduce 748143 mills to dollars.

*Ans.* \$748'143.

Reduce 3467489 mills to dollars.

*Ans.* 3467'489.

Reduce 48742 cents to dollars.

*Ans.* \$487'42.

Reduce 1234678 mills to dollars.

Reduce 3469876 cents to dollars.

Reduce \$4887'467 to mills.

Reduce 984 mills to dollars.

*Ans.* \$ '984.

Reduce 7 mills to dollars.

*Ans.* \$ '007.

Reduce \$ '014 to mills.

Reduce 17846 cents to dollars.

Re  
Re  
Re  
500 c  
Re

¶ 3.  
its pa  
weigh  
drugs  
weigh

24  
20  
12

1. I  
silver  
5 oz.  
3. I  
pwts.  
5. I  
9 grs.  
grains

App  
sicians  
pounds

\*The  
is unders  
said to b  
the stan  
Silver  
standard  
melted t  
† The  
Troy, ar

Reduce 984321 cents to mills.

Reduce 9617½ cents to dollars.

*Ans.* \$96·17½.

Reduce 206½ cents, 503 cents, 106 cents, 921½ cents, 500 cents, 726½ cents, to dollars.

Reduce 86753 mills, 96000 mills 6042 mills, to dollars.

### TROY WEIGHT.

§31. It is established by law, that the pound Troy, with its parts, multiples, and proportions, shall be the standard weight for weighing gold\* and silver in coin or bullion, drugs, and precious stones. The denominations of Troy weight are pounds, ounces, pennyweights and grains.

#### TABLE.

24 grains (grs.) make 1 pennyweight, marked pwt.  
20 pennyweights - - - 1 ounce, - - - - - oz.  
12 ounces - - - - - 1 pound, - - - - - lb.

1. How many grains in a silver tankard weighing 3 lb. 5 oz. ?

2. In 19680 grains how many pounds, &c.

3. Reduce 210 lb. 8 oz. 12 pwt. to pennyweights.

4. In 50572 pwt. how many pounds ?

5. In 7 lb. 11 oz. 3 pwt. 9 grs. of silver, how many grains ?

6. Reduce 45681 grains to pounds.

### APOTHECARIES' WEIGHT.

Apothecaries' weight† is used by apothecaries and physicians, in compounding medicines. The denominations are pounds, ounces, drams, scruples, and grains.

#### TABLE.

20 grains, (grs.) make 1 scruple, marked ℞.  
3 scruples - - - 1 dram, - - - 3.  
8 drams - - - 1 ounce, - - - ̄3.  
12 ounces - - - 1 pound, - - - lb,

\*The fineness of gold is tried by fire, and is reckoned in CARATS, by which is understood the 24th part of any quantity; if it lose nothing in the trial, it is said to be 24 carats fine; if it lose 2 carats, it is then 22 carats fine, which is the standard for gold.

Silver which abides the fire without loss is said to be 12 ounces fine. The standard for silver coin is 11 oz. 2 pwt. of fine silver, and 18 pwt. of copper melted together.

† The pound and ounce apothecaries' weight and the pound and ounce Troy, are the same, only differently divided, and subdivided.

7. In 9 lb. 83. 1 3. 2 9. | 8. Reduce 55790 grs. to  
19 grs., how many grains. | pounds.

### AVOIRDUPOIS WEIGHT.\*

It is established by law that the pound Avoirdupois with its parts, &c. shall be considered as the standard for weighing every thing commonly sold by weight, except those articles, in weighing which, Troy weight is used. The denominations are tons, hundreds, quarters, pounds ounces, and drams.

#### TABLE.

16 drams, (drs.)	make	1 ounce,	-	marked	-	oz.
16 ounces	- - - -	1 pound,	- - - -	-	-	lb.
28 pounds	- - - -	1 quarter,	- - - -	-	-	qr.
4 quarters	- - - -	1 hundred weight,	- - - -	-	-	cwt.
20 hundred weight	- -	1 ton,	- - - -	-	-	T.

*Note 1.* In this kind of weight, the words *gross* and *net* are used. Gross is the weight of the goods, together with the box, bale, bag, cask, &c, which contains them. Net weight is the weight of the goods only, after deducting the weight of the box, bale, bag, or cask, &c., and all other allowances.

*Note 2.* A hundred weight, it will be perceived, is 112 lb. Merchants at the present time, in the principal sea ports of the United States, buy and sell by the 100 pounds.

9. A merchant would put 109 cwt. 0 qrs. 12 lb. of raisins into boxes, containing 26 lb. each; how many boxes will it require?

11. In 12 tons, 15 cwt. 1 qr. 19 lb. 6 oz. 12 dr. how many drams?

13. In 28 lb. avoirdupois, how many pounds Troy?

10. In 470 boxes of raisins, containing 26 lb. each, how many cwt.?

12. In 7323500 drams, how many tons?

14. In 34 lb. 0 oz. 6 pwt. 16 grs. Troy, how many pounds avoirdupois?

\* 175 oz. Troy = 192 oz. avoirdupois, and 175 lb. Troy = 144 lb. avoirdupois, 1 lb. Troy = 5760 grains, and 1 lb. avoirdupois = 7000 grains Troy.

Cloth  
sold by  
English  
measur  
the Eng  
be used  
quarter

4 nails,  
4 quart  
3 quart  
5 quart  
6 quart

16.  
how m  
18.  
many  
Note.

Long  
things,  
The de  
rods, y

3 ba  
12 in  
3 fe  
5 1/2 y  
40 ro  
8 fu  
3 m  
60 g

360 d



# CLOTH MEASURE.

Cloth measure is used in selling cloths and other goods, sold by the yard, or ell. It is established by law that the English yard with its parts, &c. shall be the standard for measuring all kinds of cloth or stuffs made of wool flax, &c. the English ell, when there is a special contract for it may be used with its parts. The denominations are ells, yards, quarters, and nails.

## TABLE.

4 nails, (na.) or 9 inches,	make 1 quarter,	marked	qr.
4 quarters, or 36 inches,	- 1 yard,	- - -	yd.
3 quarters, - - - - -	- 1 ell Flemish,	- -	E. Fl.
5 quarters, - - - - -	- 1 ell English,	- -	E. E.
6 quarters, - - - - -	- 1 ell French,	- -	E. Fr.

16. In 573 yds. 1 qr. 1 na. how many nails?  
17. In 9173 nails, how many yards?

18. In 151 ells Eng. how many yards?  
19. In 1883 yards, how many ells English?

Notes. Consult ¶ 28. ex. 16.

# LONG MEASURE.

Long measure is used in measuring distances, or other things, where *length* is considered without regard to *breadth*. The denominations are degrees, leagues, miles, furlongs, rods, yards, feet, inches, and barley-corns.

## TABLE.

3 barley-corns, (bar.)	make 1 inch,	- marked	- in.
12 inches,	- - - - -	1 foot,	- - - - - ft.
3 feet,	- - - - -	1 yard,	- - - - - yd.
5½ yards, or 16½ feet,	- - - - -	1 rod, perch, or poie,	r. p.
40 rods, or 220 yards,	- - - - -	1 furlong,	- - - - - fur.
8 furlongs, or 320 rods,	- - - - -	1 mile,	- - - - - M.
3 miles,	- - - - -	1 league,	- - - - - L.
60 geographical, or 69½ statute miles,	- - - - -	1 degree,	- - - - - deg. or °
360 degrees,	- - - - -	{ a great circle, or circumference of the earth.	

It is established by law, that the *Paris* foot with its parts, &c. shall be the standard measure of length, for measuring land, wood, timber, stone, masons', carpenters', and joiners' work. The English foot may be used when there is a special contract for it.

## TABLE.

12 lines make 1 inch.

12 inches - - 1 foot.

6 feet - - - 1 toise.

1 French foot =  $1\frac{17}{60}$  English feet.

3 toises make 1 rod.

10 rods - - - 1 arpent.

84 arpens - - 1 league.

20. How many barley-corns will reach round the globe, it being 360 degrees?

*Note.* To multiply by 2 is to take the multiplicand 2 times; to multiply by 1 is to take the multiplicand 1 time; to multiply by  $\frac{1}{2}$  is to take the multiplicand half a time, that is, the half of it. Therefore, to reduce 360 degrees to statute miles, we multiply first by the whole number, 69, and to the product add half the multiplicand. Thus:

$$\begin{array}{r} 360 \\ \times 69\frac{1}{2} \\ \hline \end{array}$$
69 $\frac{1}{2}$ 

3240

2160

180 half the multiplicand.

25020 statute miles in 360°.

22. How many inches from Quebec to Three Rivers, supposing it to be 90 miles?

24. How many times will a wheel 16 feet and 6 inches in circumference, turn round in the distance from Quebec to St Annes, supposing it to be 60 miles?

21. In 4755801600 barley-corns, how many degrees?

*Note.* The barley-corns being divided by 3, and that quotient by 12, we have 132106500 feet, which are to be reduced to rods. We cannot easily divide by 16 $\frac{1}{2}$  on account of the fraction  $\frac{1}{2}$ ; but 16 $\frac{1}{2}$  feet = 33 half feet, in 1 rod; and 132106500 feet = 264211200 half feet, which, divided by 33, gives 8006400 rods.

Hence, when the divisor is encumbered with a fraction,  $\frac{1}{2}$  or  $\frac{1}{4}$ , &c., we may reduce the divisor to halves or fourths &c., and reduce the dividend to the same; then the quotient will be the true answer.

23. In 30539520 inches, how many miles?

25. If a wheel 16 feet 6 in. in circumference, turn round 19200 times in going from Quebec to St. Annes, what is the distance?

26. In pens, how many to

Square thing, w inations inches.

it require a yard in make 3 make 2 3 feet w will clear

3 feet

3 feet	
3 feet	
3 feet	
3 feet	

144 square

12

es i

9 square

in

30 $\frac{1}{2}$  square

sq

40 square

4

640

*Note.*

in length

inches in

625 square

its parts,  
measuring  
and join-  
there is a

26. In 28 leagues, 43 arpens, how many feet? how many rods? how many arpens? how many toises? how many rods?

rod.  
arpent.  
league.

### LAND OR SQUARE MEASURE.

Square measure is used in measuring land, and any other thing, where *length* and *breadth* are considered. The denominations are miles, acres, roods, perches, yards, feet and inches.

0 barley-  
egrees?

-corns be-  
and that  
we have  
which are to

We can-  
by 16½ on  
ion ½; but  
feet, in 1  
00 feet =  
et, which,  
s 8006400

e divisor is  
a fraction,  
may reduce  
s or fourths  
e dividend  
n the quo-  
ue answer.

20 inches,

3 feet 6 in.  
urn round  
going from  
es, what is

¶ 32. 3 feet in length make a yard in long measure; but it requires 3 feet in length, and 3 feet in breadth, to make a yard in square measure; 3 feet in length and 1 foot wide, make 3 square feet; 3 feet in length and 2 feet wide, make 2 times 3, that is, 6 square feet; 3 feet in length and 3 feet wide make 3 times 3, that is 9 square feet. This will clearly appear from the annexed figure.

3 feet=1 yard.



It is plain, also, that a square foot, that is, a square 12 inches in length, and 12 inches in breadth, must contain  $12 \times 12 = 144$  square inches.

### TABLE.

144 square inches	= $12 \times 12$ ; that is, 12 inches in length and 12 inches in breadth,	} make 1 square foot.
9 square feet	= $3 \times 3$ ; that is, 3 feet in length and 3 feet in breadth	
60½ square yards	= $5½ \times 5½$ , or $272½$ square feet = $16½ \times 16½$	} 1 square rod, perch or pole
40 square rods,		
4	roods, or 160 square rods,	1 acre.
640	acres,	1 square mile.

*Note.* Gunter's chain, used in measuring land is 4 rods in length. It consists of 100 links, each link being 7½ inches in length; 25 links make 1 rod, long measure, and 625 square links make 1 square rod.

## FRENCH SQUARE MEASURE.

144	square inches	make	1 square foot.
36	- feet	- -	1 toise.
9	- toises	- -	1 rod.
100	- rods	- -	1 arpent.
7056	- arpens	-	1 league.
62500	french feet	=	71289 English feet.

Reduce 16 leagues to feet, — to toises, — to rods.

Reduce 98764321 feet to toises, — to rods, — to arpens, — to leagues.

28. In 17 acres 3 roods 12 poles, how many square feet? how many acres?

Note. In reducing rods to feet, the multiplier will be square feet to be divided by 272 $\frac{1}{4}$ . To multiply by  $\frac{1}{4}$ , is to take a fourth part of the multiplier; that is, to the lowest denomination contained in the same as shown † 28, it; then reduce the dividend to fourths, that is, to the same denomination, as shown † 31, ex. 20.

30. Reduce 64 square miles to square feet. 31. In 1,784,217,600 sq. feet, how many square miles?

32. There is a town 6 miles square; how many square miles in that town? how many acres? 33. Reduce 23040 acres to square miles.

## SOLID OR CUBIC MEASURE.

Solid or cubic measure is used in measuring things that have length, breadth, and *thickness*; such as timber, wood, stone, bales of goods, &c. The denominations are cords, tons, yards, feet, and inches.

† 33. It has been shown, that a square yard contains  $3 \times 3 = 9$  square feet. A cubic yard is 3 feet long, 3 feet wide, and 3 feet thick. Were it 3 feet long, 3 feet wide and one foot thick, it would contain 9 cubic feet; if 2 feet thick, it would contain  $2 \times 9 = 18$  cubic feet; and, as it is

3 feet



1728 s

tha

12

27 so

40 fe

fe

128 s

is

width,

Note.

16 solid

1 foot i

1 cord.

10

32. I

timber

34.

how m

36.

wood t

38.


how m

many s

40.

inches

3 feet thick, it does contain  $3 \times 9 = 27$  cubic feet. This will clearly appear from the annexed figure.



It is plain, also, that a cubic foot, that is, a solid 12 inches in length, 12 inches in breadth, and 12 inches in thickness, will contain  $12 \times 12 \times 12 = 1728$  solid or cubic inches.

### TABLE.

1728 solid inches, $= 12 \times 12 \times 12$ , that is, 12 inches in length, 12 in breadth, 12 in thickness,	}	make 1 solid foot.
27 solid feet, $= 3 \times 3 \times 3$		1 solid yard.
40 feet of round timber, or 50 feet of hewn timber,	}	1 ton or load.
128 solid feet, $= 8 \times 4 \times 4$ , that is 8 feet in length, 4 feet in width, and 4 feet in height,		1 cord of wood.

*Note.* What is called a *cord foot*, in measuring wood, is 16 solid feet; that is, 4 feet in length, 4 feet in width, and 1 foot in height, and 8 such feet, that is, 8 *cord feet* make 1 cord.

### FRENCH SOLID MEASURE.

1728 solid inches make 1 solid foot.

216 - - feet - 1 toise.

1000 French feet = 1218.186432 English feet.

32. Reduce 9 tons of round timber to cubic inches.

34. In 37 cord feet of wood, how many solid feet?

36. Reduce 64 cord feet of wood to cords.

38. In 16 cords of wood, how many cord feet? how many solid feet?

40. In 12 toises how many inches?

33. In 622080 cubic inches how many tons of round timber?

35. In 592 solid feet of wood, how many cord feet?

37. In 8 cords of wood, how many cord feet?

39. In 2048 solid feet of wood, how many cord feet? how many cords?

41. In 834692773 inches how many feet? how many toises?

## WINE MEASURE.

It is established by law that the wine gallon with its parts, &c. shall be the standard liquid measure, for measuring wine, cider, beer, and all other liquids commonly sold by gauge, or measure of capacity. The denominations are tuns, pipes, hogsheads, barrels, gallons, quarts, pints, and gills.

## TABLE.

4 gills (gi.)	-	make	- -	1 pint, marked	pt.
2 pints	-	-	-	1 quart, -	qt.
4 quarts	-	-	-	1 gallon, -	gal.
31½ gallons	-	-	-	1 barrel, -	bar.
63 gallons	-	-	-	1 hogshead, -	hhd.
2 hogsheads	-	-	-	1 pipe, -	P.
2 pipes, or four hogsheads	-	-	-	1 tun, -	T.

*Note.* A gallon, wine measure, contains 231 cubic inches.

42. Reduce 12 pipes of wine to pints.	43. In 12096 pints of wine, how many pipes?
44. In 9 P. 1 hhd. 22 gals. 3 qts. how many gills?	45. Reduce 39032 gills to pipes?
46. In a tun of cider, how many gallons?	47. Reduce 252 gallons to tuns.

## ALE OR BEER MEASURE.

Ale or beer measure is used in measuring ale, beer, and milk. The denominations are hogsheads, barrels, gallons, quarts, and pints.

## TABLE.

2 pints (pts.)	make	1 quart, marked	qt.
4 quarts	-	1 gallon, -	gal.
36 gallons	-	1 barrel, -	bar.
54 gallons	-	1 hogshead, -	hhd.

*Note.* A gallon, beer measure, contains 282 cubic inches.

48 Reduce 47 bar. 18 gal. of ale to pints?	49. In 13680 pints of ale, how many barrels?
50. In 29 hhds. of beer, how many pints?	51. Reduce 12528 pints to hogsheads?

Dry  
grain.  
chald.

2 pin  
8 qu  
4 pe

36 bu  
Not  
A V  
deep,  
It is  
parts,  
Dry

20

16

3

2

2

12

Th  
for th  
to be

52

how

54

coal

T  
days

60

60

# DRY MEASURE.

Dry measure is used in measuring all dry goods, such as grain, fruit, roots, salt, coal, &c. The denominations are chaldrons, bushels, pecks, quarts, and pints.

## TABLE.

2 pints (pts.)	make	-	1 quart,	-	marked	-	qt.
8 quarts,	-	-	-	1 peck,	-	-	pk.
4 pecks	-	-	-	1 bushel,	-	-	bu.
36 bushels	-	-	-	1 chaldron,	-	-	ch.

*Note.* A gallon dry measure, contains 268 $\frac{1}{2}$  cubic inches.

A Winchester bushel is 18 $\frac{1}{2}$  inches in diameter, 8 inches deep, and contains 2150 $\frac{1}{2}$  cubic inches.

It is established by law, that the Canadian Minot, with its parts, multiples, and proportions, shall be the standard in Dry Measure.

1 pot = 116'94580 English cubic feet.

20 pots make 1 minot.

## OLD MEASURE.

16 litrons	-	make	-	1	-	-	-	boisseau.
3 boisseaux	-	-	-	1	-	-	-	minot.
2 minots	-	-	-	1	-	-	-	nine.
2 mines	-	-	-	1	-	-	-	setier.
12 setiers	-	-	-	1	-	-	-	muid.

40 French cubic inches = 1 litron.

The standard measure for the sale and purchase of coal, for this Province, is the chaldron of 36 minots, each minot to be heaped up.

52. In 75 bushels of wheat how many pints?

53. In 4800 pints, how many bushels?

54. Reduce 42 chaldrons of coal to pecks.

55. In 6048 pecks, how many chaldrons?

## TIME.

The denominations of time are years, months, weeks, days, hours, minutes, and seconds.

## TABLE.

60 seconds (s.)	-	make	-	1 minute,	marked	m.
60 minutes	-	-	-	1 hour,	-	h.



24 hours	- - - - -	1 day,	- - - - -	d.
7 days	- - - - -	1 week,	- - - - -	w.
4 weeks	- - - - -	1 month,	- - - - -	mo.
13 months, 1 day and 6 hours,	} 1 common, or }	Julian year,	}	yr.
or 365 days and 6 hours,				

† 34. The year is also divided into 12 calendar months, which in the order of their succession are numbered as follows, viz.

January,	1st month, has 31 days.
February,	2d, - - - 28
March,	3d, - - - 31
April,	4th, - - - 30
May,	5th, - - - 31
June,	6th, - - - 30
July,	7th, - - - 31
August,	8th, - - - 31
September,	9th, - - - 30
October,	10th, - - - 31
November,	11th, - - - 30
December,	12th, - - - 31

*Note.* When any year can be divided by 4 without a remainder, it is called leap year, in which February has 29 days.

The number of days in each month may be easily fixed in the mind by committing to memory the following lines :

Thirty days hath September,  
April, June and November,  
February twenty-eight alone ;  
All the rest have thirty-one.

The first seven letters in the alphabet, A, B, C, D, E, F, G, are used to mark the several days of the week, and they are disposed in such a manner, for every year, that the letter A shall stand for the 1st day of January, B for the 2d, &c. In pursuance of this order, the letter which shall stand for *Sunday*, in any year, is called the *Dominical* letter for that year. The Dominical letter being known, the day of the week on which each month comes in may be readily calculated from the following couplet :

At Dover Dwells George Brown, Esquire,  
Good Carlos Finch And David Fryer.

These words correspond to the 12 months of the year, and the *first letter* in each word marks the day of the week on

which  
day  
to find  
year  
answer  
when

No  
one for  
the y

56  
be 18  
how  
you,  
hours

58.  
the 1  
14th  
sively  
60.

the co  
betwe  
land,  
settle  
which  
1783

Cir  
tude ;  
other  
cles, a

60 sec  
60 mi  
30 de  
12 sig

Not  
into 3

62.  
second

- d.  
- w.  
- mo.  
- yr.

ar months,  
ered as fol-

n any year  
by 4 with-  
t, it is call-  
in which  
29 days.

asily fixed  
ing lines :

o, E, F, G,  
d they are  
e letter A  
d, &c. In  
d for Sun-  
that year.  
the week  
calculated

year, and  
week on

which each corresponding month comes in ; whence any other day may be easily found. For example, let it be required to find on what day of the week the 4th of July falls, in the year 1827, the Dominical letter for which year is G. Good answers to July ; consequently, July comes in on a Sunday ; wherefore the 4th of July falls on Wednesday.

*Note.* There are *two* Dominical letters in *leap* years, one for January and February, and *another* for the rest of the year.

56. Supposing your age to be 15y. 10d. 11h. 37m. 45s., how many seconds old are you, allowing 365 days 6 hours to the year ?

58. How many minutes from the 1st day of January to the 14th day of August, inclusively ?

60. How many minutes from the commencement of the war between America and England, April 19th, 1775, to the settlement of a general peace, which took place Jan. 20th, 1783 ?

57. Reduce 475047465 seconds to years.

59. Reduce 325440 minutes to days.

61. In 4079160 minutes, how many years ?

### CIRCULAR MEASURE, OR MOTION.

Circular measure is used in reckoning latitude and longitude ; also in computing the revolution of the earth and other planets round the sun. The denominations are circles, signs, degrees, minutes, and seconds.

#### TABLE.

60 seconds (")	make	1 minute,	marked
60 minutes	-	1 degree,	-
30 degrees	-	1 sign,	-
12 signs, or 360 degrees,		1 circle of the zodiac.	

*Note.* Every circle, whether great or small, is divisible into 360 equal parts, called degrees.

62. Reduce 9s. 13° 25' to seconds.

63. In 1020300" how many degrees ?

The following are *denominations* of things not included in the tables :—

12 particular things	make	1 dozen.
12 dozen	-	1 gross.
12 gross, or 144 dozen,	-	1 great gross.

Also,

20 particular things	make	1 score.
6 points make 1 line,	{	used in measuring the length of
12 lines - 1 inch,		the rods of clock pendulums.
4 inches - 1 hand,	{	used in measuring the height of
		horses.
6 feet - 1 fathom,		used in measuring depths at sea.
112 pounds	make	1 quintal of fish.
24 sheets of paper	make	1 quire.
20 quires	-	1 ream.

## SUPPLEMENT TO REDUCTION.

### QUESTIONS.

1. What is reduction? 2. Of how many varieties is reduction? 3. What is understood by *different denominations*, as of money, weight, measure, &c.? 4. How are high denominations brought into lower? 5. How are low denominations brought into higher? 6. What are the denominations of Halifax currency? 7. What name is given to the currency of this Province? 8. And why? 9. Are the ratios of the different denominations to each other the same as in English money? 10. Will the rule for reduction of one denomination to another in Halifax currency, apply to all currencies in which the denominations are of the same name? 11. What is the use of Troy weight, and what are the denominations? 12. — avoirdupoise weight? — the denominations? 13. What distinction do you make between *gross* and *net* weight? 14. What distinctions do you make between long, square, and cubic measure? 15. What are the denominations in long measure? 16. — in square measure? 17. — in cubic measure? 18. How do you multiply by  $\frac{1}{4}$ ? 19. When the divisor contains a fraction, how do you proceed? 20. How is the superficial contents of a square figure found? 21. How is the solid contents of any body found in cubic measure? 22. How many solid or cubic feet of wood make a cord? 23. What is understood by a *cord foot*? 24. How many such feet make a cord? 25. What are the denominations of dry measure? 26. — of wine measure? 27. — of time? 28. — of circular measure? 29. For what is circular measure used? 30. How many rods in length is Gunter's chain? of how many links does it consist? how many links make a rod? 31. How many rods in a mile? 32. How many square rods in an acre? 33. How many pounds make 1 cwt.?

## EXERCISES.

1. In 154 dollars, at 5s. each, how many pounds, &c.

*Ans.* 38*l.* 10*s.*

2. In 36 guineas, at 1*l.* 3*s.* 4*d.* each, how many crowns, at 5*s.* 6*d.* ?

*Ans.* 131 crowns and 2*s.* 10*d.* over.

3. How many rings, each weighing 5pwt. 7grs., may be made of 3lb. 5oz. 16pwt. 2grs. of gold ?

*Ans.* 158.

4. Suppose a bridge to be 212 rods in length, how many times will a chaise wheel, 18 feet 6 inches in circumference, turn round in passing over it ?

*Ans.* 189  $\frac{11}{12}$  times.

5. In 470 boxes sugar, each 26lb., how many cwt. ?

6. In 10lb. of silver, how many spoons, each weighing 1oz. 10pwt. ?

7. How many shingles, each covering a space 4 inches one way and 6 inches the other, would it take to cover 1 square foot ? How many to cover a roof 40 feet long, and 24 wide ? (See ¶ 25.)

*Ans.* to the last, 5760 shingles.

8. How many cords of wood in a pile 26 feet long 4 feet wide, and 6 feet high ?

*Ans.* 4 cords, and 7 cord feet.

9. There is a room 18 feet in length, 16 feet in width, and 8 feet in height ; how many rolls of paper, 2 feet wide, and containing 11 yards in each roll, will it take to cover the walls ?

*Ans.* 8  $\frac{1}{2}$ .

10. How many cord feet in a load of wood 6  $\frac{1}{2}$  feet long, 2 feet wide, and 5 feet high ?

*Ans.* 4  $\frac{1}{6}$  cord feet.

11. If a ship sail 7 miles an hour, how far will she sail, at that rate, in 3w. 4d. 16h ?

12. A merchant sold 12 hhds. of brandy, at 83 a gallon ; how much did each hogshead come to, and to how much in currency did the whole amount ?

13. How much cloth, at 7*s.* a yard, may be bought for 29*l.* 1*s.* ?

14. A goldsmith sold a tankard for 10*l.* 8*s.* at the rate of 5*s.* 4*d.* per ounce ; how much did it weigh ?

15. An ingot of gold weighs 2lb. 8oz. 16pwt. ; how much is it worth at 3*d.* per pwt. ?

16. At 11 pence a pound, what will 1 T. 2cwt. 3qrs. 16lb. of lead come to ?

17. Reduce 14445 ells Flemish to ells English.

18. There is a house, the roof of which is 44  $\frac{1}{2}$  feet in length, and 20 feet in width, on each of the two sides ; if 3 shingles in width cover one foot in length, how many

shingles will it take to lay one course on this roof? if 3 courses make one foot, how many courses will there be on one side of the roof? how many shingles will it take to cover one side? — to cover both sides?

*Ans.* 16020 shingles.

19. How many steps, of 30 inches each, must a man take in travelling  $54\frac{1}{2}$  miles?

20. How many seconds of time would a person redeem in 40 years, by rising each morning  $\frac{1}{2}$  hour earlier than he now does?

21. If a man lay up  $\frac{1}{2}$  dollar each day, Sundays excepted, how many pounds would he lay up in 45 years?

22. If 9 candles are made from 1 pound of tallow, how many dozen can be made from 24 pounds and 10 ounces?

23. If one pound of wool make 60 knots of yarn, how many skeins, of ten knots each, may be spun from 4 pounds 6 ounces of wool?

## ADDITION

### OF COMPOUND NUMBERS.

¶ 35. 1. A boy bought a knife for 9 pence, and a comb for 3 pence; how much did he give for both? *Ans.* 1 shilling.

2. A boy gave 2s. 6 d. for a slate, and 4s. 6 d. for a book; how much did he give for both?

3. Bought one book for 1s. 6 d., another for 2s. 3 d., another for 7 d.; how much did they all cost? *Ans.* 4s. 4 d.

4. How many gallons are 2 qts. + 3 qts. + 1 qt.?

5. How many gallons are 3 qts. + 2 qts. + 1 qt. + 3 qts. + 2 qts.?

6. How many shillings are 2d. + 3d. + 5d. + 6d. + 7d.?

7. How many pence are 1qr. + 2qrs. + 3qrs. + 2qrs. + 1qr.?

8. How many pounds are 4s. + 10s. + 15s. + 1s.?

9. How many minutes are 30 sec. + 45sec. + 20sec.?

10. How many hours are 40 min. + 25 min. + 6min.?

11. How many days are 4h. + 8h. + 10h. + 20h.?

12. How many yards in length are 1f. + 2f. + 1f.?

¶ 35.

13.

lin. 1

14.

1 ft. 8

15.

16.

wine;

qts.; h

17.

oxen fo

he pay

When

to write

nominat

those of

OPF

*Ans.* 4

Next,

the 1s. v

=1£.

the 1£.

be 41£.

*Note.*

a higher

takes of

In additi

thus, be

tween sh

The a

RULE f

I. W

same de

II. A

denomin

13. How many feet are 4in. + 8in. + 10in. + 3in. + 1in. ?

14. How much is the amount of 1yd. 2ft. 6in. + 2yds. 1 ft. 8 in. ?

15. What is the amount of 2s. 6d. + 4s. 3d. + 7s. 8d. ?

16. A man has 2 bottles, which he wishes to fill with wine; one will contain 2 gal. 3 qts. 1 pt. and the other 3 qts.; how much wine can be put in them ?

17. A man bought a horse for 15£. 14s. 6d., a pair of oxen for 20£. 2s. 8d., and a cow for 5£. 6s. 4d.; what did he pay for all ?

When the numbers are large, it will be most convenient to write them down, placing those of the same kind, or denomination, directly under each other, and, beginning with those of the least value, to add up each kind separately.

OPERATION.

£.	s.	d.
15	14	6
20	2	8
5	6	4
<hr/>		
Ans. 41	3	6

In this example, adding up the column of pence, we find the amount to be 18 pence, which being = 1s. 6d., it is plain that we may write down the 6d. under the column of pence, and reserve the 1s. to be added in with the *other* shillings.

Next, adding up the column of shillings, together with the 1s. which we reserved, we find the amount to be 23s. = 1£. 3s. Setting the 3s. under its *own* column, we add the 1£. with the *other* pounds, and, finding the amount to be 41£. we write it down, and the work is done.

Ans. 41£. 3s. 6d.

*Note.* It will be recollected, that, to reduce a lower into a higher denomination, we divide by the number which it takes of the lower to make one of the higher denomination. In addition, this is usually called *carrying* for that number: thus, between pence and shillings, we carry for 12, and between shillings and pounds, for 20, &c.

The above process may be given in the form of a general *RULE* for the *Addition of Compound Numbers*.

I. Write the numbers to be added so that those of the same denomination may stand directly under each other.

II. Add together the numbers in the column of the lowest denomination, and carry for that number which it takes of

the same to make 1 of the next *higher* denomination. Proceed in this manner with all the denominations, till you come to the last, whose amount is written as in simple numbers.

*Proof.* The same as in addition of simple numbers.

## EXAMPLES FOR PRACTICE.

## HALIFAX CURRENCY.

£	s.	d.	qr.	£	s.	d.	£	s.	d.
46	11	3	2	72	9	6½	183	19	4
16	7	4	4	18	0	10½	8	17	10
538	19	7	1	36	16	5½	15	4	

£.	s.	d.	£.	s.	d.	£.	s.	d.
14	0	7½	37	15	8	61	3	2½
8	15	3	14	12	9½	7	16	8
62	4	7	17	14	9	29	13	10½
4	17	8	23	10	9½	12	16	2
23	0	4½	8	6	0	0	7	5½
6	6	7	14	0	5½	24	13	0
91	0	10½	54	2	7½	5	0	10½

No examples in Federal Money are here introduced, although the general rule for the addition of all compound numbers is precisely applicable to the addition of Federal Money, since that consists of *different denominations*. In Federal Money the denominations increase and decrease in a *decimal ratio*. The pupil is therefore referred to the rules for the *Addition, Subtraction, Multiplication and Division of Decimals*, which are the same absolutely with the rules for the addition, subtraction, multiplication and division of Federal money.

## TROY WEIGHT.

lb.	oz.	pwt.	gr.	oz.	pwt.	gr.	oz.	pwt.	gr.
36	7	10	11	6	14	9			18
42	6	9	13	8	6	16		13	16
81	7	16	15	3	11	10	3	7	4

Boug  
cup, we  
2 pwt.

T. c  
14  
25  
7

A ma  
23 cwt.  
19 cwt.  
how ma

yds. 9  
36  
41  
65

Ther  
viz. 36  
3 na. ;

De  
5  
21  
67



n. Pro.  
till you  
ple num.

Bought a silver tankard, weighing 2lb. 3 oz., a silver cup, weighing 3 oz. 10 pwt. and a silver thimble, weighing 2 pwt. 13 grs. ; what was the weight of the whole ?

bers.

AVOIRDUPOIS WEIGHT.

T.	cwt.	qr.	lb.	oz.	dr.	cwt.	qr.	lb.	oz.	dr.
14	11	1	16	5	10	16	3	18	6	14
25	0	2	11	8	15	2	16	8	12	
7	18	0	25	11	9			22	11	10

s. d.  
19 4  
17 10  
15 4

A man bought 5 loads of hay, weighing as follows, viz. 23 cwt. (=1 T. 3 cwt.) 2 qrs. 17 lb. ; 21 cwt. 1 qr. 16 lb. ; 19 cwt. 0 qr. 24 lb. ; 24 cwt. 3 qrs. ; 11 cwt. 0 qr. 1 lb. ; how many tons in the whole ?

CLOTH MEASURE.

yds.	qr.	na.	E. Fl.	qr.	na.	E. E.	qr.	na.
36	1	2	41	1	2	75	4	2
41	2	3	18	2	3	31	1	0
65	3	1	57	0	2	28	3	1

roduced, al  
compound  
of Federa.  
ations. In  
decrease in  
to the rules  
Division of  
the rules for  
sion of Fed.

There are four pieces of cloth, which measure as follows, viz. 36 yds. 2 qrs. 1 na. ; 18 yds. 1 qr. 2 na. ; 46 yds. 3 qrs. 3 na. ; 12 yds. 0 qr. 2 na. ; how many yards in the whole ?

LONG MEASURE.

Deg.	mi.	fur.	r.	ft.	in.	bar.	Mi.	fur.	pol.
59	46	6	29	15	10	2	3	7	
216	39	1	36	14	6	1			
678	53	7	24	9	8	1	8	6	27

et. gr.  
18  
13 16  
7 4

## LAND OR SQUARE MEASURE.

<i>Pol.</i>	<i>ft.</i>	<i>in.</i>	<i>A. rood.</i>	<i>pol.</i>	<i>ft.</i>	<i>in.</i>
36	179	137	56	3	37	245
19	248	119	29	1	28	93
12	96	75	416	2	31	128

There are 3 fields, which measure as follows, viz. 17 A. 3r. 16 p. : 28 A. 5r. 18p. ; 11A. 0r. 25p. ; how much land in the three fields ?

## SOLID OR CUBIC MEASURE.

<i>Ton.</i>	<i>ft.</i>	<i>in.</i>	<i>yds.</i>	<i>ft.</i>	<i>in.</i>	<i>cords.</i>	<i>ft.</i>
29	36	1229	75	22	1412	37	119
12	19	64	9	26	195	4	110
8	11	917	3	19	1091	48	127

## WINE MEASURE.

<i>Hhds.</i>	<i>gal.</i>	<i>qts.</i>	<i>pts.</i>	<i>Tun.</i>	<i>hhd.</i>	<i>gal.</i>	<i>qts.</i>
51	53	1	1	37	2	37	2
27	39	3	0	19	1	50	1
9	13	0	1	28	2	0	0

A merchant bought two casks of brandy, containing as follows, viz. 70 gal. 3 qts. ; 67 gal. 1qt. ; how many hogsheads, of 63 gal. each, in the whole ?

## DRY MEASURE.

<i>Bus.</i>	<i>p.</i>	<i>qt.</i>	<i>pt.</i>	<i>Ch.</i>	<i>bus.</i>	<i>p.</i>	<i>qt.</i>
36	2	5	1	48	27	3	5
19	3	7	0	6	29	1	7

TIME.

Y.	mo.	w.	d.	h.	m.	s.
57	11	3	6	23	55	11
84	9	2	0	16	42	18
32	6	0	5	5	18	5

Y.	m.	w.	d.
40	3	1	5
16	7	0	4
27	5	2	0

z. 17 A.  
uch land

ds. ft.  
7 119  
4 110  
3 127

qts.  
2  
1  
0

taining as  
any hogs

# SUBTRACTION

## OF COMPOUND NUMBERS.

† 36. 1. A boy bought a knife for 9 pence, and sold it for 1s. 4d.; how much did he gain by the bargain?

2. A boy bought a slate for 2s. 6d., and a book for 3s. 6d.; how much more was the cost of the book than of the slate?

3. A boy owed his playmate 2s.; he paid him 1s. 6d.; how much did he then owe him?

4. Bought two books; the price of one was 4s. 6d., the price of the other 3s. 9d.; what was the difference of their costs?

5. A boy lent 5s. 3d.; he received in payment 2s. 6d.; how much was then due?

6. A man has a bottle of wine containing 2 gallons and 3 quarts; after turning out 3 quarts, how much remained?

7. How much is 4 gal. less 3 gal? 4 gal. — (less) 2qts? 4 gal. — 1qt? 4 gal. — 1 gal. 1qt? 4 gal. — 1 gal. 2qts? 4 gal. — 1 gal. 3qts? 4 gal. — 2 gal. 3qts? 4 gal. 1qt. — 1 gal. 3qts?

8. How much is 1ft. — (less) 6in? 1ft. — 8in? 6ft. 3 inches, — 1ft. 6in? 7ft. 8in. — 4ft. 2in? 7ft. 8in. — 5ft. 10in?

9. What is the difference between 4£ 6s. and 1£ 8s?

10. How much is 3£ — (less) 1s? 3£ — 2s? 3£ — 3s? 3£ — 15s? 3£ 4s. — 2£ 6s? 10£ 4s. — 5£ 8s?

11. A man bought a horse for 30£ 4s. 8d., and a cow for 5£ 14s. 6d.; what is the difference of their costs?

## OPERATION.

	£.	s.	d.
Minuend,	30	4	8
Subtrahend,	5	14	6
Ans.	24	10	2

As the two numbers are large, it will be convenient to write them down, the less under the greater, pence under pence, shillings under shillings, &c. We may now take 6d. from 8d., and

there will remain 2d. Proceeding to the shillings, we cannot take 14s. from 4s., but we may borrow, as in simple numbers, 1 from the pounds, = 20s., which joined to the 4s. makes 24s, from which taking 14s. leaves 10s, which we set down. We must now carry 1 to the 5£ making 6£ which taken from 30£ leaves 24£ and the work is done.

*Note.* The most convenient way in borrowing is, to subtract the subtrahend from the figure borrowed, and add the difference to the minuend. Thus, in the above example, 14 from 20 leaves 6, and 4 is 10.

The process in the foregoing example may be presented in the form of a *RULE for the Subtraction of Compound Numbers.*

I. Write down the sums or quantities, the less under the greater, placing those numbers which are of the same denomination directly under each other.

II. Beginning with the least denomination, take successively the lower number in each denomination from the upper, and write the remainder underneath, as in subtraction of simple numbers.

III. If the lower number of any denomination be greater than the upper, borrow as many units as make one of the next higher denomination, subtract the lower number therefrom, and to the remainder add the upper number, remembering always to add 1 to the next higher denomination for that which you borrowed.

*Proof.* Add the remainder and the subtrahend together, as in subtraction of simple numbers; if the work be right, the amount will be equal to the minuend.

## EXAMPLES FOR PRACTICE.

## HALIFAX CURRENCY.

£.	s.	d.	£.	s.	d.
79	17	8	103	3	2
35	12	4	71	12	5

1. A  
and rec  
ed due

2. A  
it, lost

3. A  
19£ 12s  
the oxe

4. A  
time, 13  
quantity

5. A  
when it  
cask; h

6. Th  
the lid a  
the tank

7. Fro  
8. Bo

7lb; so  
11lb. 5oz  
weight o

9. Bo  
2qrs; ty  
he other  
he black

10. F

11. A

£.	s.	d.
81	10	11 $\frac{1}{4}$
29	13	3

£.	s.	d.
245	12	0
27	9	4 $\frac{1}{4}$

520	11	3
109	17	4

631	14	7
6	19	8

# MISCELLANEOUS EXAMPLES.

1. A merchant sold goods to the amount of 136£ 7s. 6 $\frac{1}{2}$ d. and received in payment 50£ 10s. 4 $\frac{1}{2}$ d; how much remained due? *Ans.* 85£ 17s. 1 $\frac{1}{2}$ d.

2. A man bought a farm for 1256£ 10s. and, in selling it, lost 87£ 10s. 6d; how much did he sell it for? *Ans.* 1168£ 19s. 6d.

3. A man bought a horse for 27£ and a pair of oxen for 19£ 12s. 8 $\frac{1}{2}$ d; how much was the horse valued more than the oxen?

4. A merchant drew from a hogshead of molasses, at one time, 13gal. 3qts; at another time, 5gal. 2qts. 1pt; what quantity was there left? *Ans.* 43gal. 2qts. 1pt.

5. A pipe of brandy, containing 118gal. sprang a leak, when it was found only 97gal. 3qts. 1pt. remained in the cask; how much was the leakage?

6. There was a silver tankard which weighed 3lb. 4oz; the lid alone weighed 5oz. 7pwt. 13grs; how much did the tankard weigh without the lid?

7. From 15lb. 2oz. 5pwt. take 9oz. 8pwt. 10grs.

8. Bought a hogshead of sugar, weighing 9cwt. 2qrs. 17lb; sold at three several times as follows, viz. 2cwt. 1qr. 11lb. 5oz; 2qrs. 18lb. 10oz; 25lb. 6oz; what was the weight of sugar which remained unsold?

*Ans.* 6cwt. 1qr. 17lb. 11oz.

9. Bought a piece of black broadcloth, containing 36 yds. 2qrs; two pieces of blue, one containing 10 yds. 3qrs. 2na, the other, 18yds. 3qrs. 3na; how much more was there of the black than of the blue?

10. From 28 miles, 5fur. 16r. take 15 m. 6fur. 26r. 12ft.

11. A farmer has two mowing fields; one containing 13

acres 6 roods; the other, 14 acres 3 roods: he has two pastures also; one containing 26 A. 2 r. 27 p; the other, 45 A. 5 r. 33 p: how much more has he of pasture than of mowing?

12. From 64 A. 2r. 11p. 29ft. take 26 A. 5r. 34p. 132ft.

13. From a pile of wood, containing 21 cords, was sold, at one time, 8 cords 76 cubic feet; at another time, 5 cords 7 cord feet; what was the quantity of wood left?

14. How many days, hours and minutes of any year will be future time on the 4th day of July, 20 minutes past 3 o'clock, P. M? *Ans.* 180 days, 8 hours, 40 minutes.

15. On the same day, hour and minute of July, given in the above example, what will be the difference between the past and future time of that month?

16. A note, bearing date Dec. 28th, 1826, was paid Jan. 2d, 1827; how long was it at interest?

The distance of time from one date to that of another may be found by subtracting the first date from the last, observing to number the months according to their order. (¶ 34.)

OPERATION.

A. D. { 1827. 1st m. 2d day. *Note.* In casting interest, each month is reckoned 30 days.  
 { 1826. 12 — 28 —  
*Ans.* 0 0 4 days.

17. A note, bearing date Oct. 20th, 1823, was paid April 25th, 1825; how long was the note at interest?

18. What is the difference of time from Sept. 29, 1816, to April 2d, 1819? *Ans.* 2y. 6m. 3d.

19. London is  $51^{\circ} 32'$ , and Montreal  $45^{\circ} 30'$ , N. latitude; what is the difference of latitude between the two places? *Ans.*  $6^{\circ} 2'$ .

20. Montreal is  $73^{\circ} 20'$ , and the city of Washington is  $77^{\circ} 43'$  W. longitude; what is the difference of longitude between the two places? *Ans.*  $4^{\circ} 23'$ .

21. The island of Cuba lies between  $74^{\circ}$  and  $85^{\circ}$  W. longitude; how many degrees in longitude does it extend?

¶ 37. 1. When it is 12 o'clock at the most easterly extremity of the island of Cuba, what will be the hour at the most westerly extremity, the difference in longitude being  $11^{\circ}$ ?

*Note.* The circumference of the earth being  $360^{\circ}$ , and the earth performing one entire revolution in 24 hours, it

follows, that the motion of the earth, on its surface, from west to east, is

15° of motion in 1 hour of time; consequently,

1° of motion in 4 minutes of time, and

1' of motion in 4 seconds of time.

From these premises it follows, that, when there is a difference in longitude between two places, there will be a corresponding difference in the hour, or time of the day. The difference in longitude being 15°, the difference in time will be one hour, the place *easterly* having the time of the day 1 hour *earlier* than the place *westerly*, which must be particularly regarded.

If the difference in longitude be 1°, the difference in time will be 4 minutes, &c.

Hence,—If the difference in longitude, in degrees and minutes, between two places, be multiplied by 4, the product will be the difference in time, in minutes and seconds, which may be reduced to hours.

We are now prepared to answer the above question.

11°

4

44 minutes.

Hence, when it is 12 o'clock at the most easterly extremity of the island, it will be 16 minutes past 11 o'clock at the most western extremity.

2. Montreal being 73° 20' W. longitude and Washington 77° 43'; when it is 3 o'clock at the city of Washington, what is the hour at Montreal?

*Ans.* 17 minutes 32 seconds past 3 o'clock

3. Lower Canada being about 73°, and the Sandwich Islands about 155° W. longitude, when it is 28 minutes past 6 o'clock, A. M. at the Sandwich Islands, what will be the hour in Lower Canada?

*Ans.* 12 o'clock at noon, lacking 4 minutes

## MULTIPLICATION & DIVISION

### OF COMPOUND NUMBERS.

¶ 38. 1. A man bought 2 yards of cloth, at 1s. 6d. per yard; what was the cost?

H 2



2. If 2 yards of cloth cost 3 shillings, what is that per yard?

3. A man has three pieces of cloth, each measuring 10 yds. 3qrs; how many yards in whole?

4. If 3 equal pieces of cloth contain 32 yds. 1 qr., how much does each piece contain?

5. A man has five bottles, each containing 2 gal. 1 qt. 1 pt; how much wine do they all contain?

6. A man has 11 gal. 3qts. 1pt. of wine, which he would divide equally into 5 bottles; how much must he put into each bottle?

7. How many shillings are 3 times 8d? —  $3 \times 9d$ ? —  $3 \times 10d$ ? —  $4 \times 7d$ ? —  $7 \times 6d$ ? —  $10 \times 9d$ ? —  $2 \times 3qrs$ ? —  $5 \times 2qrs$ ?

8. How much is one third of 2 shillings? —  $\frac{1}{3}$  of 2s. 3d? —  $\frac{1}{3}$  of 2s. 6d? —  $\frac{1}{3}$  of 2s. 4d? —  $\frac{1}{3}$  of 3s. 6d? —  $\frac{1}{3}$  of 7s. 6d? —  $\frac{1}{3}$  of  $1\frac{1}{2}d$ ? —  $\frac{1}{3}$  of  $2\frac{1}{2}d$ ?

9. At 1£ 5s. 8 $\frac{1}{2}$ d. per yard, what will 6 yards of cloth cost? 10. If 6 yards of cloth cost 7£ 14s. 4 $\frac{1}{2}$ d, what is the price per yard?

Here, as the numbers are large, it will be most convenient to write them down before multiplying and dividing.

#### OPERATION.

£ s. d. qr.

1 5 8 3 price of 1 yard.  
Number of yards.

Ans. 7 14 4 2 cost of 6 yards.

#### OPERATION.

£ s. d. qr.

6) 7 14 4 2 cost of 6 yards.  
1 5 8 3 price of 1 yard.

Proceeding after the manner of short division, 6 is contained in 7£ 1 time, and 1£ over; we write down the 1£, and 4 to carry quotient, and reduce the remainder (1£) to shillings, (20s.) which, with the given shillings, (14s.) make 34s; and 4 to carry makes 34s = 6 in 34s. goes 5 times, and 4£ and 14s. over; 6 times 4s. over; 4s. reduced to pence 1£ are 6£, and 1 to carry = 48d, which, with the given pence, (4d.) make 52d; 6 down, and it is plain, that in 52d. goes 8 times, and 4d. the united products arising over; 4d. = 16qrs, which,

from t  
tions i  
ing fr  
numbe

11.  
by 7.  
13.  
of 5 p  
a pair  
15.  
each c  
6 qts,

17.  
cloth  
coats,  
3 na.

19.  
each c  
gills,

21.  
weigh  
weigh

23.  
hogsh  
9cwt.

25.  
weigh  
how

W  
of m  
comp  
compa

Th  
above  
produ

from the several denomina-  
tions is the real product aris-  
ing from the whole compound  
number.

with the given qrs. (2) = 18  
qrs; 6 in 18qrs. goes 3 times;  
and it is plain, that the united  
quotients arising from the  
several denominations, is the  
real quotient arising from the  
whole compound number.

11. Multiply 3*£* 4*s*. 6*d*.  
by 7.

12. Divide 22*£* 11*s*. 6*d*.  
by 7.

13. What will be the cost  
of 5 pairs of shoes at 10*s*. 6*d*.  
a pair?

14. A . . . . . for 5  
pairs of . . . . . is that a  
pair?

15. In 5 barrels of wheat,  
each containing 2 bus. 3 pks.  
6 qts, how many bushels?

16. If 14 bus. 2 pks. 6 qts. of  
wheat be equally divided into  
5 barrels, how many bushels  
will each contain?

17. How many yards of  
cloth will be required for 9  
coats, allowing 4 yds. 1qr.  
3 na. to each?

18. If 9 coats contain 39  
yds. 3qrs. 3na, what does 1  
coat contain?

19. In 7 bottles of wine,  
each containing 2qts. 1pt. 3  
gills, how many gallons?

20. If 5 gal. 1 gill of wine  
be divided equally into 7 bot-  
tles, how much will each con-  
tain?

21. What will be the  
weight of 8 silver cups, each  
weighing 5oz. 12pwt. 17grs?

22. If 8 silver cups weigh  
3lb. 9oz. 1pwt. 16grs, what  
is the weight of each?

23. How much sugar in 12  
hogsheads, each containing  
9cwt. 3qrs. 21lb?

24. If 119cwt. 1qr. of su-  
gar be divided into 12 hogs-  
heads, how much will each  
hogshead contain?


25. In 15 loads of hay, each  
weighing 1T. 3cwt. 2qrs,  
how many tons?


26. If 15 teams be loaded  
with 17T. 12cwt. 2qrs. of hay,  
how much is that to each team?

*When the multiplier, or divisor, exceeds 12, the operations  
of multiplying and dividing are not so easy, unless they be  
composite numbers; in that case, we may make use of the  
component parts, or factors, as was done in simple numbers.*

Thus 15, in the example 15 being a composite num-  
above, is a composite number, and 3 and 5 its compo-  
produced by the multiplica-  
nent parts, or factors, we may







6"



# Photographic Sciences Corporation

**23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503**

1.8 2.0 2.2 2.5  
3.6 3.2 2.8  
4.5 5.0 5.6

10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

tion of 3 and 5, ( $3 \times 5 = 15$ .) We may, therefore, multiply 1T. 3cwt. 2qrs. by one of these component parts, or factors, and the quotient thence arising by the other, or factors, and that product by the other, which will give the true answer, as already taught, true answer, as has been already taught, ( $\text{T } 20$ .)

## OPERATION.

T. cwt. qrs.

1 3 2

3 one of the factors.

3 10 2

5 the other factor.

17 12 2 the answer.

## OPERATION.

T. cwt. qrs.

One factor, 3) 17 12 2

The other factor, 5) 5 17 2

Ans. 1 3 2

27. What will 24 barrels of flour cost, at  $2\text{£ } 12\text{s. } 4\text{d.}$  a barrel?

29. What will 112lb. of sugar cost  $7\text{d.}$  per lb?

*Note.* 8, 7, and 2, are factors of 112.

31. How much brandy in 84 pipes, each containing 112 gal. 2qts. 1pt. 3g?

33. What will 139yds. of cloth cost, at  $3\text{£ } 6\text{s. } 5\text{d.}$  per yard?

139 is not a composite number. We may, however, decompose this number thus,  $139 = 100 + 30 + 9$ .

We may now multiply the price of 1 yard by 10, which will give the price of 10 yards, and this product again by 10, which will give the price of 100 yards.

28. Bought 24 barrels of flour for  $62\text{£ } 16\text{s.}$ ; how much was that per barrel?

30. If 1 cwt. of sugar cost  $3\text{£ } 7\text{s. } 8\text{d.}$ , what is that per lb?

32. Bought 84 pipes of brandy, containing 9468 gal. 1qt. 1pt; how much in a pipe?

34. Bought 139 yards of cloth for  $461\text{£ } 11\text{s. } 11\text{d.}$ ; what was that per yard?

When the divisor is such a number as cannot be produced by the multiplication of small numbers, the better way is to divide after the manner of long division, setting down the work of dividing and reducing in manner as follows:



wt. 2qrs. by  
ponent parts,  
the quotient  
y the other,  
e the true  
ady taught,

OPERATION.

T. cwt. qr.  
17 12 2

)5 17 2

. 1 3 2

barrels of  
how much  
el?  
sugar cost  
is that per

pipe of  
9468 gal.  
in a pipe?  
yards of  
11s. 11d;  
yard?  
is such a  
produced  
n of small  
way is to  
anner of  
ng down  
and re-  
follows :

We may then multiply the price of 10 yards by 3, which will give the price of 30 yards, and the price of 1 yard by 9, which will give the price of 9 yards, and these three products, added together, will evidently give the price of 139 yards; thus :

£. s. d.  
3 6 5 price of 1 yard.  
10

33 4 2 price of 10 yards.  
10

332 1 8 price of 100 yds.  
99 12 6 price of 30 yds.  
29 17 9 price of 9 yds.

461 11 11 price of 139 yds.

*Note.* In multiplying the price of 10 yards (33£ 4s. 2d.) by 3, to get the price of 30 yards, and in multiplying the price of 1 yard (3£ 6s. 5d.) by 9, to get the price of 9 yards, the multipliers, 3 and 9, need not be written down, but may be carried in the mind.

	£	s.	d.
139)	461	11	11 (3£
	417		
	44		
	20		
	891	(6s.	
	834		
	57		
	12		
	695	(5d.	
	695		

The divisor, 139, is contained in 461£ 3 times, (3£,) and a remainder of 44£, which must now be reduced to shillings, multiplying it by 20, and bringing in the given shillings, (11s,) making 891s, in which the divisor is contained 6 times, (6s,) and a remainder of 57s, which must be reduced to pence, multiplying it by 12, and bringing in the given pence, (11d,) together making 695d, in which the divisor is contained 5 times, (5d,) and 9, need not be written and no remainder.

The several quotients, 3£ 6s. 5d. evidently make the answer.

The processes in the foregoing examples may now be presented in the form of a

**RULE for the Multiplication of Compound Numbers.**

**RULE for the Division of Compound Numbers.**

1. When the multiplier does not exceed 12, multiply successively the numbers of each denomination, beginning with many times it is contained in

1. When the divisor does not exceed 12, in the manner of short division, find how many times it is contained in

the least, as in multiplication the highest denomination, un-  
of simple numbers, and carry der which write the quotient,  
as in addition of compound and if there be a remainder,  
numbers, setting down the reduce it to the next less de-  
whole product of the highest nomination, adding thereto the  
denomination.

number given, if any, of that  
denomination, and divide as  
before; so continue to do  
through all the denominations  
and the several quotients will  
be the answer.

II. If the multiplier exceed 12, and be a *composite* num-  
ber, we may multiply first by 12, and be a *composite*, we may di-  
vide first by one of the com-  
one of the component parts, ponent parts, that quotient by  
that product by another, and another, and so on, if the com-  
so on, if the component parts ponent parts be more than  
be more than two; the last two, the last quotient will be  
product will be the product re- the quotient required.

III. When the multiplier exceeds 12, and is not a compos-  
ite, multiply first by 10, posite number, divide after the  
and this product by 10, which manner of long division, set-  
will give the product for 100; ting down the work. Evi-  
and if the hundreds in the mul- ding and reducing.  
tiplier be more than one, mul-  
tiply the product of 100 by the  
number of hundreds; for the  
tens, multiply the product of  
10 by the number of tens; for  
the units, multiply the multi-  
plicand; and these several pro-  
ducts will be the product re-  
quired.

#### EXAMPLES FOR PRACTICE.

##### HALIFAX CURRENCY.

£	s.	d.	£	s.	d.
Multiply 81	6	5	93	4	11
by 11		17	21	48	
<hr/>			<hr/>		

Div

1. V

cloth

yard?

3. In

each o

1lb; he

5. H

wheat i

aining

7. W

accoo o

b 1.

9. If

feet of

many ro

1 days

1 39.

oth co

Note,

ence, o

£	s.	d.	£	s.	d.	
Multiply	98	8	10	64	11	2
by	78				93	

986	11	4	899	5	3
	73			145	

£	s.	d.	£	s.	d.	
Divide	77	11	9	143	9	3
"	140	2	3	1950	7	4
"	360	5	2	47	9	0
"	7856	8	9	562	8	3

MISCELLANEOUS EXAMPLES.

1. What will 350 yards of cloth cost, at 4s. 7½d. per yard?
2. Bought 359 yds. of cloth for 83£ 0s. 4½d; what was that a yard?
3. In 241 barrels of flour, each containing 1 cwt. 3 qrs. 1 lb; how many cwt?
4. If 441 cwt. 13 lb. of flour be contained in 241 barrels, how much in a barrel?
5. How many bushels of wheat in 135 bags, each containing 2 bu. 3 pks?
6. If 371 bu. 1 pk. of wheat be divided equally into 135 bags, how much will each bag contain?
7. What will 35 cwt. of tobacco cost, at 3s. 10½d. per lb?
8. At 759£ 10s. for 35 cwt. of tobacco, what is that per lb?
9. If 14 men build 42 rods of wall in one day, how many rods will they build in 12 days?
10. If 14 men build 92 rods of stone wall in 7½ days, how much is that per day?

† 39. 1. At 10s. per yard, what will 17849 yards of cloth cost?

Note. Operations in multiplication of pounds, shillings, pence, or of any compound numbers, may be facilitated by

taking *aliquot parts* of a *higher denomination*. Thus, in this last example, if the price had been 20s. i. e. 1s. per yard, it is clear, the price of the whole would have been equal to the whole number of yards in pounds, 17849; but the price is 10s. i. e. 1s. per yard, and so the price of the whole will be equal to  $\frac{1}{2}$  the number of yards, 17849 in pounds; 8924s. 10s.

When one quantity is contained in another exactly 2, 3, 4, 5, &c. times, it is called an *aliquot* or *even* part of that quantity; thus 6d. is an aliquot part of a shilling, because  $6d. \times 2 = 1$  shilling; so 3d. is an aliquot part of a shilling;  $3d. \times 4 = 1s.$  So 5s. is an aliquot part of a pound, for  $5s. \times 4 = 1s.$ ; and 8s. 4d. is an aliquot part of a pound, for  $8s. 4d. \times 6 = 1s.$ , &c.

From the illustration of the last example it appears, that, when the price per yard, pound, &c. is one of these aliquot parts of a shilling, or a pound, the cost may be found by dividing the *given number of yards, pounds, &c.* by that number which it takes of the price to make 1s. or 1s. If the price be 6d. we divide by 2; if 5s. we divide by 4; if 3s. 4d. by 6, &c. &c. This manner of calculating by aliquot parts, is called *Practice*.

2. What cost 34648 yards of cloth, at 10s. or 1s. per yard? — at 5s. = 1s. per yard? — at 4s. = 1s. per yard? — at 3s. 4d. = 1s. per yard? — at 2s. = 1s. per yard? *Ans. to last, 3464s. 16s.*

3. What cost 7430 pounds of sugar, at 6d. = 1s. per lb? — at 4d. = 1s. per lb? — at 3d. = 1s. per lb? — at 2d. = 1s. per lb? — at 1d. = 1s. per lb? *Ans. to the last, 1238s. 9d. = 46s. 8d.*

4. At 3s. 16s. per cwt, what will 2qrs. = 1cwt. cost? — what will 1qr. = 1cwt. cost? — what will 16lb. = 1cwt. cost? — what will 14lb. = 1cwt. cost? — what will 8lb. = 1cwt. cost? *Ans. to the last, 5s. 5d.*

5. What cost 340 yards of cloth, at 12s. 6d. per yard? 12s. 6d. = 10s. (= 1s.) and 2s. 6d. (= 1s.); therefore,

$$\frac{1}{2} \frac{1}{2} 340$$

$$170s. = \text{cost at 10s. per yard.}$$

$$42s. 10s. = \text{at 2s. 6d. per yard.}$$

$$\text{Ans. } 212s. 10s. = \text{at 12s. 6d. per yard.}$$

1.  
num  
num  
ditio  
what  
same  
Whe  
do y  
anoth  
west  
is th  
most  
being  
may  
oilita  
1s.?  
called

1.  
each  
ing 1  
tanks  
11oz.

No  
follow  
2.  
of gol  
weigh  
3.  
3cwt.  
4.  
many

us, in this  
per yard,  
n equal to  
t the price  
the whole  
n pounds ;

actly 2, 3,  
art of that  
ng, because  
a shilling ;  
nd, for 5s.  
pound, for

pears, that,  
hese aliquot  
be found by  
ke. by that  
s. or 1£. If  
ide by 4 ; if  
ating by ali-

s. or 1£ per  
4s = 1£ per  
at 2s = 10£  
3464£ 16s.  
= 1s. per lb ?  
er lb ? —

46£ 8s. 9d.  
= 1cwt. cost ?  
will 16lb. =  
? — what  
last, 5s. 5½d.  
s. per yard ?  
therefore,

yard.  
yard.  
yard.

Or,

$$10s. = \frac{1}{2} £ 340$$

$$2s. 6d = \frac{1}{4} \text{ of } 10s. ) 170£ \quad \text{at } 10s. \text{ per yard.}$$

$$42£ \quad 10s. \text{ at } 2s. 6d. \text{ per yard.}$$

$$Ans. 212£ \quad 10s. \text{ at } 12s. 6d. \text{ per yard.}$$

## SUPPLEMENT TO COMPOUND NUMBERS.

### QUESTIONS.

1. What distinction do you make between simple and compound numbers ? (¶ 26.) 2. What is the rule for addition of compound numbers ? 3. — for subtraction of, &c. ? 4. There are *three* conditions in the rule given for multiplication of compound numbers ; what are they, and the methods of procedure under each ? 5. The same questions in respect to the division of compound numbers ? 6. When the multiplier or divisor is encumbered with a fraction, how do you proceed ? 7. How is the distance of time from one date to another found ? 8. How many degrees does the earth revolve from west to east in 1 hour ? 9. In what time does it revolve 1° ? Where is the time or hour of the day earlier—at the place most easterly or most westerly ? 10. The difference in longitude between two places being known, how is the difference in time calculated ? 11. How may operations, in the multiplication of compound numbers, be facilitated ? 12. What are some of the aliquot parts of 1£ ? — of 1s. ? — of 1cwt. ? 13. What is this manner of operating usually called ?

### EXERCISES.

1. A gentleman is possessed of 1½ dozen of silver spoons, each weighing 3oz. 5pwt ; 2 doz. of tea spoons, each weighing 15pwt. 14gr ; 3 silver cans, each 9oz. 7pwt ; 2 silver tankards, each 21oz. 15pwt ; and 6 silver porringers, each 11oz. 18pwt ; what is the weight of the whole ?

$$Ans. 18lb. 4oz. 3pwt.$$

*Notes.* Let the pupil be required to reverse and prove the following examples :

2. An English guinea should weigh 5pwt. 6gr ; a piece of gold weighs 3pwt. 17gr ; how much is that short of the weight of a guinea ?

3. What is the weight of 6 chests of tea, each weighing 3cwt. 2qrs. 9lb ?

4. In 35 pieces of cloth, each measuring 27 yards, how many yards ?

5. How much brandy in 9 casks, each containing 45gal. 3qts. 1pt. 1

6. If 31cwt. 2qrs. 20lb. of sugar be distributed equally into 4 casks, how much will each contain ?

7. At  $4\frac{1}{2}$ d. per lb., what costs 1cwt. of rice ? — 2cwt.; — 3cwt. 1

*Note.* The pupil will recollect, that 8, 7 and 2 are factors of 112, and may be used in place of that number.

8. If 800cwt. of cocoa cost 18£ 13s. 4d. what is that per cwt. 1 what is it per lb. 1

9. What will  $9\frac{1}{2}$ cwt. of copper cost at 5s. 9d. per lb. ?

10. If  $6\frac{1}{2}$ cwt. of chocolate cost 72£ 16s. what is that per lb. 1

11. What cost 456 bushels of potatoes, at 2s. 6d. per bushel. 1

*Note.* 2s. 6d. is  $\frac{1}{4}$  of 1£. (See ¶ 39.)

12. What cost 86 yards of broadcloth, at 15s. per yard ?

*Note.* Consult ¶ 39, ex. 5.

13. What cost 7846 pounds of tea, at 7s. 6d. per lb. ? — at 14s. per lb. ? — 13s. 4d. ?

14. At 894.25 per cwt. what will be the cost of 2qrs. of tea ? — of 3 qrs. ? — of 14 lbs. ? — of 21 lbs. ? — of 16 lbs. ? — of 24 lbs. ?

*Note.* Consult ¶ 39, ex. 4 and 5.

15. What will be the cost of 2 pks. and 4qts. of wheat, at 8s. 6d. per bushel. 1

16. Supposing a meteor to appear so high in the heavens as to be visible at Montreal,  $73^{\circ} 20'$ , at the city of Washington,  $77^{\circ} 43'$ , and at the Sandwich Islands,  $155^{\circ}$  W. longitude, and that its appearance at the city of Washington be at 7 minutes past 9 o'clock in the evening ; what will be the hour and minute of its appearance at Montreal and at the Sandwich Islands ?

## FRACTIONS.

¶ 40. We have seen, (¶ 17,) that numbers expressing whole things are called *integers*, or *whole numbers* ; but that, in division, it is often necessary to divide or break a whole thing into *parts*, and that these parts are called *fractions*, or *broken numbers*.

It w  
or unit  
ed thir  
sixths,  
the nu  
if the  
tenths,  
thus, 1  
taught,  
the nar  
the line

The  
a unit  
these p

1. If  
tion is  
— 4

or a wh

2. If  
pieces l  
pie ? if  
what fra

It is i  
division,  
dividend  
fraction  
numera  
tient ar  
that is,  
same as

Henc  
sign of

1. expe

3. If  
part of

A six  
is 4 and

4. If  
of an  
apples,



It will be recollected, (§ 14, ex. 11,) that when a thing or unit is divided into 3 parts, the parts or fractions are called *thirds*; when into four parts, *fourths*; when into six parts, *sixths*; that is, the fraction takes its name or *denomination* from the number of parts, into which the unit is divided. Thus, if the unit be divided into 16 parts, the parts are called *sixteenths*, and 5 of these parts would be 5 *sixteenths*, expressed thus,  $\frac{5}{16}$ . The number below the short line, (16,) as before taught, (§ 17,) is called the *denominator*, because it gives the name or *denomination* to the parts; the number above the line is called the *numerator*, because it *numbers* the parts.

The *denominator* shows how many parts it takes to make a unit or whole thing; the *numerator* shows how many of these parts are expressed by the *fraction*.

1. If an orange be cut into 5 equal parts, by what fraction is 1 part expressed? — 2 parts? — 3 parts? — 4 parts? — 5 parts? how many parts make unity or a whole orange?

2. If a pie be cut into 8 equal pieces, and 2 of these pieces be given to Harry, what will be his fraction of the pie? if 5 pieces be given to John, what will be his fraction? what fraction or part of the pie will be left?

It is important to bear in mind, that fractions arise from *division*, (§ 17,) and that the *numerator* may be considered a *dividend*, and the *denominator* a *divisor*, and the *value* of the fraction is the *quotient*; thus,  $\frac{1}{2}$  is the quotient of 1 (the numerator) divided by 2, (the denominator;)  $\frac{1}{4}$  is the quotient arising from 1 divided by 4, and  $\frac{3}{4}$  is 3 times as much, that is, 3 divided by 4; thus, one fourth part of 3 is the same as 3 fourths of 1.

Hence, in all cases, a fraction is always expressed by the *sign of division*.

$\frac{3}{4}$  expresses the quotient, of which  $\left\{ \begin{array}{l} 3 \text{ is the dividend, or numerator.} \\ 4 \text{ is the divisor, or denominator.} \end{array} \right.$

3. If 4 oranges be equally divided among 6 boys, what part of an orange is each boy's share?

A sixth part of 1 orange is  $\frac{1}{6}$ , and a sixth part of 4 oranges is 4 such pieces, =  $\frac{4}{6}$ . *Ans.*  $\frac{2}{3}$  of an orange.

4. If 3 apples be equally divided among 5 boys, what part of an apple is each boy's share? if 4 apples, what? if 2 apples, what? if 5 apples, what?



5. What is the quotient of 1 divided by 31 — of 2 by 3? — of 1 by 4? — of 2 by 4? — of 3 by 4? — of 5 by 7? — of 6 by 8? — of 4 by 5? — of 2 by 14?

6. What part of an orange is a third part of 2 oranges? — one fourth of 2 oranges? —  $\frac{1}{4}$  of 3 oranges? —  $\frac{1}{2}$  of three oranges? —  $\frac{1}{2}$  of 4? —  $\frac{1}{2}$  of 2? —  $\frac{1}{2}$  of 5? —  $\frac{1}{2}$  of 3? —  $\frac{1}{2}$  of 2?

*A proper fraction.* Since the denominator shows the number of parts necessary to make a whole thing, or 1, it is plain, that, when the numerator is less than the denominator, the fraction is less than a unit, or whole thing; it is then called a proper fraction. Thus,  $\frac{1}{2}$ ,  $\frac{3}{4}$ , &c. are proper fractions.

*An improper fraction.* When the numerator equals or exceeds the denominator, the fraction equals or exceeds unity, or 1, and is then called an improper fraction. Thus,  $\frac{3}{2}$ ,  $\frac{5}{4}$ , &c. are improper fractions.

*A mixed number*, as already shown, is one composed of a whole number and a fraction. Thus,  $14\frac{1}{2}$ ,  $13\frac{3}{4}$ , &c. are mixed numbers.

7. A father bought 4 oranges, and cut each orange into 6 equal parts; he gave to Samuel 3 pieces, to James 5 pieces, to Mary 7 pieces, and to Nancy 9 pieces; what was each one's fraction?

Was James' fraction proper, or improper? Why?

Was Nancy's fraction proper, or improper? Why?

*To change an improper fraction to a whole or mixed number.*

¶ 41. It is evident, that every improper fraction must contain one or more whole ones, or integers.

How many whole apples are there in 4 halves ( $\frac{1}{2}$ ) of an apple? — in  $\frac{3}{2}$ ? — in  $\frac{5}{2}$ ? — in  $\frac{7}{2}$ ? — in  $\frac{9}{2}$ ? — in  $\frac{11}{2}$ ? — in  $\frac{13}{2}$ ? — in  $\frac{15}{2}$ ?

2. How many yards in  $\frac{3}{2}$  of a yard? — in  $\frac{5}{2}$  of a yard? — in  $\frac{7}{2}$ ? — in  $\frac{9}{2}$ ? — in  $\frac{11}{2}$ ? — in  $\frac{13}{2}$ ? — in  $\frac{15}{2}$ ? — in  $\frac{17}{2}$ ? — in  $\frac{19}{2}$ ?

3. How many bushels in 8 pecks? that is, in  $\frac{8}{4}$  of a bushel? — in  $\frac{10}{4}$ ? — in  $\frac{12}{4}$ ? — in  $\frac{14}{4}$ ? — in  $\frac{16}{4}$ ? — in  $\frac{18}{4}$ ? — in  $\frac{20}{4}$ ?

This finding how many integers, or whole things, are contained in any improper fraction, is called *reducing an improper fraction to a whole or mixed number*.

of 2 by 3?  
1 — of 5  
2 by 14?  
2 oranges 1  
ages 1 —  
—  $\frac{1}{2}$  of 51

ws the num-  
, it is plain,  
inator, the  
then called  
fractions.

quals or ex-  
ceeds unity,  
Thus,  $\frac{3}{2}$ ,  $\frac{5}{3}$ .

imposed of a  
 $\frac{1}{3}$ , &c. are

orange into  
to James 5  
s; what was

Why 1  
Why 1

mixed number.

fraction must

halves ( $\frac{1}{2}$ ) of  
— in  $\frac{3}{2}$ !

n  $\frac{3}{4}$  of a yard!  
— in  $\frac{1}{2}$ !

n  $\frac{1}{2}$  of a bush-  
— in  $\frac{3}{4}$ !

le things, are  
d reducing as

4. If I give 27 children  $\frac{1}{4}$  of an orange each, how many oranges will it take? It will take  $\frac{27}{4}$ ; and it is evident, OPERATION. that dividing the numerator, 27, (= the

4)27

Ans.  $6\frac{3}{4}$  oranges.

Hence, To reduce an improper fraction to a whole or mixed number,—RULE: Divide the numerator by the denominator; the quotient will be the whole or mixed number.

### EXAMPLES FOR PRACTICE.

5. A man, spending  $\frac{1}{8}$  of a pound a day, in 83 days would spend  $\frac{83}{8}$  of a pound; how many pounds would that be?

Ans.  $10\frac{3}{8}$ l.

6. In  $\frac{1417}{60}$  of an hour, how many whole hours?

The 60th part of an hour is a minute: therefore the question is evidently the same as if it had been, in 1417 minutes, how many hours?

Ans.  $23\frac{17}{60}$  hours.

7. In  $\frac{730\frac{1}{2}}{8}$  of a shilling, how many units or shillings?

Ans.  $730\frac{1}{2}$  shillings.

8. Reduce  $\frac{1417}{60}$  to a whole or mixed number.

9. Reduce  $\frac{38}{100}$ ,  $\frac{706}{100}$ ,  $\frac{275}{100}$ ,  $\frac{1788}{1000}$ ,  $\frac{3456}{1000}$ , to whole or mixed numbers.

To reduce a whole or mixed number to an improper fraction.

47 42. We have seen, that an improper fraction may be changed to a whole or mixed number; and it is evident, that, by reversing the operation, a whole or mixed number may be changed to the form of an improper fraction.

1. In 2 whole apples, how many halves of an apple? Ans. 4 halves; that is,  $\frac{4}{1}$ . In 3 apples, how many halves? in 4 apples? in 6 apples? in 10 apples? in 24? in 60? in 170? in 492?

2. Reduce 2 yards to thirds. Ans.  $\frac{2}{3}$ . Reduce 23 yards to thirds. Ans.  $\frac{23}{3}$ . Reduce 3 yards to thirds —  $3\frac{1}{3}$  yards. —  $3\frac{2}{3}$  yards. — 5 yards. —  $5\frac{1}{3}$  yards. —  $6\frac{2}{3}$  yards.

3. Reduce 2 bushels to fourths. —  $2\frac{1}{2}$ bu. — 6 bushels. —  $6\frac{1}{4}$  bushels. —  $7\frac{3}{4}$  bushels. —  $25\frac{3}{4}$  bushels.

4. In  $16\frac{1}{2}$  pounds, how many  $\frac{1}{12}$  of a pound?  $\frac{1}{12}$  make 1 pound: if, therefore, we multiply 16 by 12, that is, multiply the whole number by the denominator, the

product will be the number of 12ths in 16£ :  $16 \times 12 = 192$ , and this, increased by the numerator of the fraction, (5,) evidently gives the whole number of 12ths ; that is,  $\frac{197}{12}$  of a pound, *Ans.*

OPERATION.

$16\frac{5}{12}$  pounds.

12

$192 = 12$ ths in 16 pounds, or the whole number.

5 = 12ths contained in the fraction.

$197 = \frac{197}{12}$ , the answer.

Hence, *To reduce a mixed number to an improper fraction,—*

**RULE :** Multiply the whole number by the denominator of the fraction, to the product add the numerator, and write the result over the denominator.

### EXAMPLES FOR PRACTICE.

5. What is the improper fraction equivalent to  $23\frac{3}{4}$  hours ? *Ans.*  $\frac{187}{4}$  of an hour.

6. Reduce  $730\frac{3}{4}$  shillings to 12ths.

As  $\frac{1}{4}$  of a shilling is equal to 1 penny, the question is evidently the same as, in 730s. 3d., how many pence ?

*Ans.*  $\frac{8763}{4}$  of a shilling ; that is 8763 pence.

7. Reduce  $1\frac{1}{8}$ ,  $17\frac{2}{8}$ ,  $8\frac{7}{8}$ ,  $4\frac{1}{8}$ , and  $7\frac{1}{8}$  to improper fractions.

8. In  $156\frac{1}{4}$  days, how many 24ths of a day ?

*Ans.*  $\frac{3761}{4} = 3761$  hours.

9. In  $342\frac{3}{4}$  gallons, how many 4ths of a gallon ?

*Ans.*  $\frac{1371}{4}$  of a gallon = 1371 quarts.

*To reduce a fraction to its lowest or most simple terms.*

T 43. The numerator and the denominator, taken together, are called the *terms of the fraction*.

If  $\frac{1}{2}$  of an apple be divided into 2 equal parts, it becomes  $\frac{1}{4}$ . The effect on the fraction is evidently the same as if we had multiplied both of its terms by 2. In either case, the parts are made 2 times as MANY as they were before ; but they are only HALF AS LARGE ; for it will take 2 times as many fourths to make a whole one as it will take halves ; and hence it is that  $\frac{1}{4}$  is the same in value or quantity as  $\frac{1}{2}$ .

$\frac{1}{2}$  is 2 parts ; and if each of these parts be again divided into 2 equal parts, that is, if both terms of the fraction be

multi

rever

It

of the

out a

The

terms

ing a

of the

which

The

ducing

both to

both w

the sam

er than

A

out a

ure of

this is

1.

One

acre.

find, b

and, d

we fin

viding

in its

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

measu

12=192,  
tion, (5.)  
is,  $\frac{1}{12}$  of

ber.

fraction,—  
ominator of  
and write

at to 23 $\frac{1}{2}$   
of an hour.

question is  
pence 1  
763 pence.  
to inprop-

761 hours.  
n ?  
371 quarts.

e terms.  
taken to-

becomes  $\frac{1}{2}$ .  
if we had  
e, the parts  
ut they are  
as many  
ves; and  
y as  $\frac{1}{2}$ .  
in divided  
fraction be

multiplied by 2, it becomes  $\frac{1}{2}$ . Hence,  $\frac{1}{2} = \frac{2}{4} = \frac{1}{2}$ , and the reverse of this is evidently true, that  $\frac{1}{2} = \frac{2}{4} = \frac{1}{2}$ .

It follows therefore, by multiplying or dividing both terms of the fraction by the same number, we change its terms without altering its value.

Thus, if we reverse the above operation, and divide both terms of the fraction  $\frac{2}{4}$  by 2, we obtain its equal,  $\frac{1}{2}$ ; dividing again by 2, we obtain  $\frac{1}{4}$ , which is the most simple form of the fraction, because the terms are the least possible by which the fraction can be expressed.

The process of changing  $\frac{2}{4}$  into its equal  $\frac{1}{2}$ , is called *reducing the fraction to its lowest terms*. It consists in dividing both terms of the fraction by any number which will divide them both without a remainder, and the quotient thence arising in the same manner, and so on, till it appears that no number greater than 1 will again divide them.

A number, which will divide two or more numbers without a remainder, is called a *common divisor*, or *common measure* of those numbers. The greatest number that will do this is called the *greatest common divisor*.

1. What part of an acre are 128 rods ?

One rod is  $\frac{1}{160}$  of an acre, and 128 rods are  $\frac{128}{160}$  of an acre. Let us reduce this fraction to its lowest terms. We find, by trial, that 4 will exactly measure both 128 and 160, and, dividing, we change the fraction to its equal  $\frac{32}{40}$ . Again, we find that 8 is a divisor common to both terms, and, dividing, we reduce the fraction to its equal  $\frac{4}{5}$ , which is now in its lowest terms, for no greater number than 1 will again measure them. The operation may be presented thus :

$$\begin{array}{r} 8 \phantom{00} \\ 4 \overline{) \frac{128}{160}} = \frac{32}{40} = \frac{4}{5} \end{array} \text{ of an acre, answer.}$$

2. Reduce  $\frac{128}{160}$ ,  $\frac{128}{256}$ ,  $\frac{128}{400}$ , and  $\frac{128}{640}$  to their lowest terms.

Ans.  $\frac{4}{5}$ ,  $\frac{1}{2}$ ,  $\frac{2}{5}$ , and  $\frac{1}{5}$ .

*Note.* If any number ends with a cipher, it is evidently divisible by 10. If the two right hand figures are divisible by 4, the whole number is also. If it ends with an even number, it is divisible by 2; if with a 5 or 0, it is divisible by 5.

3. Reduce  $\frac{128}{160}$ ,  $\frac{128}{256}$ ,  $\frac{128}{400}$ , and  $\frac{128}{640}$  to their lowest terms.

**T 44.** Any fraction may evidently be reduced to its lowest terms by a single division, if we use the *greatest* common divisor of the two terms. The greatest common measure of any two numbers may be found by a sort of trial easily made. Let the numbers be the two terms of the fraction  $\frac{128}{160}$ . The common divisor cannot exceed the *less* number, for it must measure it. We will try, therefore, if the *less* number, 128, which measures itself, will also divide or measure 160.

$$128)160(1$$

$$128$$

$$32)128(4$$

$$128$$

128 in 160 goes 1 time, and 32 remain; 128, therefore, is not a divisor of 160. We will now try whether this remainder be not the divisor sought; for if 32 be a divisor of 128, the former divisor, it must also be a divisor of 160, which consists of 128 + 32. 32 in 128 goes 4 times, without any remainder. Consequently, 32 is a divisor of 128 and 160. And it is evidently the *greatest* common divisor of these numbers; for it must be contained at least *once more* in 160 than in 128, and no number greater than their difference, that is, greater than 32, can do it.

Hence, the rule for finding the greatest common divisor of two numbers:—Divide the greater number by the less, and that divisor by the remainder, and so on, always dividing the last divisor by the last remainder, till nothing remain. The last divisor will be the greatest common divisor required.

*Note.* It is evident, that, when we would find the greatest common divisor of more than two numbers, we may first find the greatest common divisor of two numbers, and then of that common divisor and one of the other numbers, and so on to the last number. Then will the greatest common divisor last found be the answer.

4. Find the greatest common divisor of the terms of the fraction  $\frac{31}{14}$ , and, by it, reduce the fraction to its *lowest* terms.

OPERATION.

$$21)35(1$$

$$21$$

$$14)21(1$$

$$14$$

Greatest divis. 7)14(2.

$$14$$

$$\text{Then, } 7) \frac{31}{35} = \frac{3}{5} \text{ Ans.}$$

to its low-  
est common  
measure of  
easily made.  
128. The  
for it must  
number, 128,  
be 160.  
and 32 re-  
a divisor of  
ther this re-  
ught; for if  
former divi-  
sists of 128  
remainder.  
And it is  
numbers;  
160 than in  
ce, that is,  
on divisor of  
he less, and  
ys dividing  
ing remain.  
or required.  
d the great-  
we may first  
s, and then  
umbers, and  
test common

terms of the  
lowest terms.

$$= \frac{3}{5} \text{ Ans.}$$

5. Reduce  $\frac{25}{111}$  to its lowest terms.

Ans.  $\frac{1}{11}$ .

*Note.* Let these examples be wrought by both methods; by several divisors, and also by finding the greatest common divisor.

6. Reduce  $\frac{111}{111}$  to its lowest terms.

Ans.  $\frac{1}{1}$ .

7. Reduce  $\frac{111}{111}$  to its lowest terms.

Ans.  $\frac{1}{1}$ .

8. Reduce  $\frac{111}{111}$  to its lowest terms.

Ans.  $\frac{1}{111}$ .

9. Reduce  $\frac{111}{111}$  to its lowest terms.

Ans.  $\frac{1}{1}$ .

*To divide a fraction by a whole number.*

¶ 45. 1. If 2 yards of cloth cost  $\frac{3}{4}$  of a pound, what does 1 yard cost? how much is  $\frac{3}{4}$  divided by 2?

2. If a cow consume  $\frac{1}{3}$  of a bushel of meal in 3 days, how much is that per day?  $\frac{1}{3} \div 3 =$  how much?

3. If a boy divide  $\frac{1}{2}$  of an orange among 2 boys, how much will he give each one?  $\frac{1}{2} \div 2 =$  how much?

4. A boy bought 5 cakes for  $\frac{1}{2}$  of a shilling; what did 1 cake cost?  $\frac{1}{2} \div 5 =$  how much?

5. If 2 bushels of apples cost  $\frac{1}{2}$  of a pound, what is that per bushel?

1 bushel is the half of 2 bushels; the half of  $\frac{1}{2}$  is  $\frac{1}{4}$ .  
Ans.  $\frac{1}{4}$  pound.

6. If 3 horses consume  $\frac{1}{2}$  of a ton of hay in a month, what will 1 horse consume in the same time?

$\frac{1}{2}$  are 12 parts; if 3 horses consume 12 such parts in a month, as many times as 3 are contained in 12, so many parts 1 horse will consume.  
Ans.  $\frac{1}{3}$  of a ton.

7. If  $\frac{3}{4}$  of a barrel of flour be divided equally among 5 families, how much will each family receive?

$\frac{3}{4}$  is 25 parts; 5 into 25 goes 5 times. Ans.  $\frac{3}{20}$  of a barrel.

The process in the foregoing examples is evidently dividing a fraction by a whole number; and consists, as may be seen, in dividing the *numerator*, (when it can be done without a remainder,) and under the quotient writing the *denominator*. But it not unfrequently happens, that the numerator will not contain the whole number without a remainder.

8. A man divided  $\frac{1}{2}$  of a pound equally among 2 persons; what part of a pound did he give to each?

$\frac{1}{2}$  of a pound divided into 2 equal parts will be 4ths.

Ans. He gave  $\frac{1}{4}$  of a pound to each.

9. A mother divided  $\frac{1}{4}$  a pie among 4 children ; what part of the pie did she give to each ?  $\frac{1}{4} \div 4 =$  how much ?

10. A boy divided  $\frac{1}{3}$  of an orange equally among 3 of his companions ; what was each one's share ?  $\frac{1}{3} \div 3 =$  how much ?

11. A man divided  $\frac{1}{2}$  of an apple equally between 2 children ; what part did he give to each ?  $\frac{1}{2}$  divided by 2 = what part of a whole one ?

$\frac{1}{2}$  is 3 parts : if each of these parts be divided into 2 equal parts, they will make 6 parts. He may now give 3 parts to one, and 3 to the other : but 4ths divided into 2 equal parts, become 8ths. The parts are now *twice so many*, but they are only *half so large* ; consequently,  $\frac{1}{4}$  is only half so much as  $\frac{1}{2}$ . *Ans.*  $\frac{1}{4}$  of an apple.

In these last examples, the fraction has been divided by *multiplying the denominator*, without changing the numerator. The reason is obvious ; for, by multiplying the denominator by any number, the parts are made so many times *smaller*, since it will take so many more of them to make a whole one ; and if no more of these *smaller* parts be taken than were before taken of the *larger*, that is, if the numerator be not changed, the value of the fraction is evidently made so many times less.

T 46. Hence, we have two ways to divide a fraction by a whole number :

I. Divide the numerator by the whole number, (if it will contain it without a remainder,) and under the quotient write the denominator. Otherwise,

II. Multiply the denominator by the whole number, and over the product write the numerator.

#### EXAMPLES FOR PRACTICE.

1. If 7 pounds of tobacco cost  $\frac{3}{4}$  of a pound, what is that per pound ?  $\frac{3}{4} \div 7 =$  how much ? *Ans.*  $\frac{3}{28}$  of a pound.

2. If  $\frac{1}{8}$  of an acre produce 24 bushels, what part of an acre will produce 1 bushel ?  $\frac{1}{8} \div 24 =$  how much ?

3. If 12 yards of silk cost  $\frac{1}{2}$  of a pound, what is that a yard ;  $\frac{1}{2} \div 12 =$  how much ?

4. Divide  $\frac{1}{2}$  by 16.

*Note.* When the divisor is a composite number, the intelligent pupil will perceive, that he can first divide by one component part, and the quotient thence arising by the oth-

er ; the last ex

5. D

6. D

*Note*  
an imp

*Ans.*

lowest

7. D

8. D

9. D

10. D

*Note.*

conveni

duce th

viding,

the who

by 7, a

$= \frac{1}{2}$  rem

$= 26 \frac{1}{2}$

11. D

12. D

13. D

14. D

15. D

16. D

17. D

18. D

19. D

20. D

21. D

22. D

23. D

24. D

25. D

26. D

27. D

28. D

29. D

30. D

31. D

32. D

33. D

34. D

35. D

36. D



what part  
much 1

g 3 of his  
3 = how

etween 2  
l by 2 =

to 2 equal  
e 3 parts  
o 2 equal  
many, but  
ly half so  
apple.

divided by  
e numera-  
g the de-  
so many  
f them to  
er parts be  
is, if the  
ion is evi-

fraction by

(if it will  
tient write

mber, and

, what is  
f a pound.  
part of an  
t is that a

er, the in-  
ide by one  
by the oth-

er; thus he may frequently shorten the operation. In the last example,  $16 \div 8 \times 2$ , and  $\frac{8}{3} \div 8 = \frac{1}{3}$ , and  $\frac{1}{3} \div 2 = \frac{1}{6}$ .

*Ans.*  $\frac{1}{6}$ .

5. Divide  $\frac{1}{10}$  by 12. Divide  $\frac{1}{10}$  by 21. Divide  $\frac{1}{10}$  by 24.

6. If 6 bushels of wheat cost £1  $\frac{1}{2}$  what is it per bushel?

*Note.* The mixed number may evidently be reduced to an improper fraction, and divided as before.

*Ans.*  $\frac{1}{4} = \frac{1}{4}$  of a pound, expressing the fraction in its lowest terms. (¶ 43.)

7. Divide £4  $\frac{1}{2}$  by 9.

*Quot.*  $\frac{1}{18}$  of a pound.

8. Divide 12  $\frac{1}{2}$  by 5.

*Quot.*  $\frac{1}{5} = 2\frac{1}{5}$ .

9. Divide 14  $\frac{1}{2}$  by 8.

*Quot.*  $1\frac{3}{8}$ .

10. Divide 184  $\frac{1}{2}$  by 7.

*Ans.* 26  $\frac{1}{4}$ .

*Note.* When the mixed number is large, it will be most convenient, first, to divide the whole number, and then reduce the remainder to an improper fraction; and, after dividing, annex the quotient of the fraction to the quotient of the whole number; thus, in the last example, dividing 184  $\frac{1}{2}$  by 7, as in whole numbers, we obtain 26 integers, with 2  $\frac{1}{2}$  =  $\frac{5}{2}$  remainder, which, divided by 7, gives  $\frac{5}{14}$  and 26  $\frac{5}{14}$  = 26  $\frac{1}{4}$ , *Ans.*

11. Divide 2786  $\frac{1}{2}$  by 6.

*Ans.* 464  $\frac{1}{3}$ .

12. How many times is 24 contained in 7646  $\frac{1}{2}$ ?

*Ans.* 318  $\frac{1}{4}$ .

13. How many times is 3 contained in 462  $\frac{1}{2}$ ?

*Ans.* 154  $\frac{1}{2}$ ?

To multiply a fraction by a whole number.

¶ 47. 1. If 1 yard of cloth cost  $\frac{1}{3}$  of a pound, what will 2 yards cost?  $\frac{1}{3} \times 2 =$  how much?

2. If a cow consume  $\frac{1}{3}$  of a bushel of meal in 1 day, how much will she consume in 3 days?  $\frac{1}{3} \times 3 =$  how much?

3. A boy bought 5 cakes, at  $\frac{1}{2}$  of a shilling each; what did he give for the whole?  $\frac{1}{2} \times 5 =$  how much?

4. How much is 2 times  $\frac{1}{3}$ ? — 3 times  $\frac{1}{3}$ ? — 2 times  $\frac{2}{3}$ ?

5. Multiply  $\frac{2}{3}$  by 3. —  $\frac{2}{3}$  by 2. —  $\frac{2}{3}$  by 7.

6. If a man spend  $\frac{1}{3}$  of a shilling per day, how much will he spend in 7 days?

$\frac{1}{3}$  is 3 parts. If he spend 3 such parts in 1 day, he will evidently spend 7 times 3, that is, 7 = 21 in 7 days.

Hence, we perceive, a fraction is multiplied by multiplying the numerator, without changing the denominator.

But it has been made evident, (¶ 46,) that multiplying the denominator produces the same effect on the value of the fraction, as dividing the numerator: hence, also, dividing the denominator will produce the same effect on the value of the fraction, as multiplying the numerator. In all cases, therefore, where one of the terms of the fraction is to be multiplied the same result will be effected by dividing the other; and where one term is to be divided, the same result may be effected by multiplying the other.

This principle, borne distinctly in mind, will frequently enable the pupil to shorten the operations of fractions. Thus, in the following example:

At  $\frac{2}{3}$  of a pound, for 1 pound of sugar, what will 11 pounds cost?

Multiplying the numerator by 11, we obtain for the product  $\frac{22}{3} = 7\frac{1}{3}$  of a pound for the answer.

¶ 48. But by applying the above principle, and dividing the denominator, instead of multiplying the numerator we at once come to an answer,  $\frac{2}{3}$ , in much lower terms. Hence, there are two ways to multiply a fraction by a whole number:

I. Divide the denominator by the whole number, (when it can be done without a remainder,) and over the quotient write the numerator.—Otherwise,

II. Multiply the numerator by the whole number, and under the product write the denominator. If then it be an improper fraction, it may be reduced to a whole or mixed number.

#### EXAMPLES FOR PRACTICE.

1. If one man consume  $\frac{1}{3}$  of a barrel of flour in a month, how much will 18 men consume in the same time? — 6 men? — 9 men? *Ans. to the last.  $1\frac{1}{2}$  barrels.*

2. What is the product of  $\frac{1}{12}$  multiplied by 40?  $\frac{1}{12} \times 40 =$  how much? *Ans.  $23\frac{1}{3}$ .*

3. Multiply  $\frac{1}{12}$  by 10. — by 18. — by 21. — by 36. — by 48. — by 60.

*Note.* When the multiplier is a composite number, the pupil will recollect (¶ 11.) that he may multiply first by one component part, and that product by the other. Thus,

ultiplying the

ultiplying the

of the frac-

value of the

cases, there-

be multiplied

e other; and

may be effect-

ill frequently

ctions. Thus,

what will 11

n for the pro-

le, and divid-

numerator we

terms. Hence,

whole number:

number, (when it

the quotient

number, and un-

then it be an

whole or mixed

or in a month,

time? — 6

t. 1½ barrels.

y 40?  $\frac{1}{2} \times$

Ans. 23½.

21. — by

number, the

ultiply first by

ther. Thus,

in the last example, the multiplier 60 is equal to  $12 \times 5$ ; therefore,  $\frac{1}{12} \times 12 = 1$ , and  $\frac{1}{5} \times 5 = 1$ , Ans.

4. Multiply  $5\frac{1}{2}$  by 7. Ans. 40½.

*Note.* It is evident, that the mixed number may be reduced to an improper fraction, and multiplied, as in the preceding examples; but the operation will usually be shorter, to multiply the fraction and whole number separately, and add the results together. Thus, in the last example, 7 times 5 are 35; and 7 times  $\frac{1}{2}$  are  $3\frac{1}{2} = 5\frac{1}{2}$ , which, added to 35, make 40½, Ans.

Or, we may multiply the fraction first, and, writing down the fraction, reserve the integers, to be carried to the product of the whole number.

5. What will  $9\frac{1}{2}$  tons of hay come to at 3£ per ton? Ans. 28£. 19s.

6. If a man travel  $2\frac{5}{10}$  miles in 1 hour, how far will he travel in 5 hours? — in 8 hours? — in 12 hours? — in 3 days, supposing he travel 12 hours each day? Ans. to the last, 77½ miles.

*Note.* The fraction is here reduced to its lowest terms; the same will be done in all the following examples.

*To multiply a whole number by a fraction.*

† 49. 1. If 36 pounds be paid for a piece of cloth, what costs  $\frac{3}{4}$  of it?  $36 \times \frac{3}{4}$  — how much?

$\frac{3}{4}$  of the quantity will cost  $\frac{3}{4}$  of the price;  $\frac{3}{4}$  a time 36 pounds, that is  $\frac{3}{4}$  of 36 pounds, implies that 36 be first divided into 4 equal parts, and then that 1 of these parts be taken 3 times; 4 into 36 goes 9 times, and 3 times 9 is 27. Ans. 27 pounds.

From the above example it plainly appears that the object in multiplying by a fraction, whatever may be the multiplicand, is to take of the multiplicand a part, denoted by the multiplying fraction; and that this operation is composed of two others, viz. a division by the denominator of the multiplying fraction, and a multiplication of the quotient by the numerator. It is a matter of indifference, as it respects the result, which of these operations precedes the other, for  $36 \times \frac{3}{4} = 27$ , the same as  $36 \div 4 \times 3 = 27$ .

Hence, — *To multiply by a fraction, whether the multiplicand be a whole number or a fraction, — RULE.*

Divide the multiplicand by the denominator of the multi-

plying fraction, and multiply the quotient by the numerator; or, (which will often be found more convenient in practice,) first multiply by the numerator, and divide the product by the denominator.

Multiplication, therefore, when applied to fractions, does not always imply augmentation or increase, as in whole numbers; for, when the multiplier is less than unity, it will always require the product to be less than the multiplicand, to which it would be only equal if the multiplier were 1.

We have seen, (¶ 10,) that, when two numbers are multiplied together, either of them may be made the multiplier, without effecting the result. In the last example, therefore, instead of multiplying 16 by  $\frac{2}{3}$ , we may multiply  $\frac{2}{3}$  by 16, (¶ 47,) and the result will be the same.

#### EXAMPLES FOR PRACTICE.

2. What will 40 barrels of meal come to at  $\frac{2}{3}$  of a pound per barrel?  $40 \times \frac{2}{3}$  = how much?

3. What will 24 yards of cloth cost at  $\frac{2}{3}$  of a pound per yard?  $24 \times \frac{2}{3}$  = how much?

4. How much is  $\frac{1}{2}$  of 90? —  $\frac{2}{3}$  of 360? —  $\frac{1}{10}$  of 45?

5. Multiply 45 by  $\frac{1}{10}$ . Multiply 20 by  $\frac{1}{4}$ .

*To multiply one fraction by another.*

¶ 50. 1. A man owning  $\frac{2}{3}$  of a farm, sold  $\frac{1}{3}$  of his share; what part of the whole farm did he sell?  $\frac{1}{3}$  of  $\frac{2}{3}$  is how much?

We have just seen, (¶ 49,) that to multiply by a fraction, is to divide the multiplicand by the denominator, and to multiply the quotient by the numerator.  $\frac{2}{3}$  divided by 3, the denominator of the multiplying fraction, (¶ 46,) is  $\frac{2}{9}$ , which, multiplied by 2, the numerator, (¶ 48,) is  $\frac{4}{9}$ . *Ans.*

The process, if carefully considered, will be found to consist in multiplying together the two numerators for a new numerator, and the two denominators for a new denominator.

#### EXAMPLES FOR PRACTICE.

2. A man, having  $\frac{2}{3}$  of a pound, gave  $\frac{1}{10}$  of it for a dinner what did the dinner cost him? *Ans.*  $\frac{2}{30}$  pound.

3. Multiply  $\frac{2}{3}$  by  $\frac{1}{4}$ . Multiply  $\frac{3}{10}$  by  $\frac{2}{5}$ . *Product,*  $\frac{2}{15}$ .

4. How much is  $\frac{1}{2}$  of  $\frac{2}{3}$  of  $\frac{3}{4}$  of  $\frac{1}{2}$ ?

*Note.* Fractions like the above, connected by the word

the numeration in practice the product

actions, does in whole unity, it will multiplicand, were 1. rs are multiplier, ple, there-multiply  $\frac{1}{2}$

of a pound pound per

$\frac{7}{10}$  of 45?

and  $\frac{3}{4}$  of his  $\frac{3}{4}$  of  $\frac{1}{2}$  is

a fraction, and to mul-3, the de- $\frac{1}{5}$ , which,

and to con-a new numerator.

for a din- $\frac{1}{10}$  pound. product,  $\frac{1}{10}$

the word

of, are sometimes called compound fractions. The word *or* implies their continual multiplication into each other.

Ans.  $\frac{1}{2} \times \frac{1}{3} = \frac{1}{6}$ .

When there are several fractions to be multiplied continually together, as the *several numerators* are factors of the new numerator, and the *several denominators* are factors of the new denominator, the operation may be shortened by dropping those factors which are the same in both terms, on the principle explained in ¶ 43. Thus, in the last example,  $\frac{1}{2}$ ,  $\frac{3}{4}$ ,  $\frac{1}{3}$ , we find a 4 and a 3 both among the numerators and among the denominators; therefore we drop them multiplying together only the remaining numerators,  $2 \times 7 = 14$ , for a new numerator, and the remaining denominators,  $5 \times 8 = 40$ , for a new denominator, making  $\frac{14}{40} = \frac{7}{20}$ , Ans. as before.

5.  $\frac{2}{3}$  of  $\frac{1}{2}$  of  $\frac{3}{4}$  of  $\frac{1}{5}$  of  $\frac{7}{8}$  of  $\frac{1}{10}$  of  $\frac{1}{12}$  of  $\frac{1}{15}$  = how much? Ans.  $\frac{1}{180}$ .

6. What is the continual product of 7,  $\frac{1}{2}$ ,  $\frac{1}{3}$  of  $\frac{1}{4}$  and  $3\frac{1}{2}$ ?

Note. The integer 7 may be reduced to the form of an improper fraction, by writing a unit under it for a denominator, thus,  $\frac{7}{1}$ .

Ans.  $2\frac{1}{2}$ .

7. At  $\frac{1}{2}$  of a pound a yard, what will  $\frac{1}{2}$  of a yard of cloth cost?

8. At  $1\frac{1}{2}$  pounds per barrel for flour, what will  $\frac{1}{10}$  of a barrel cost?

$1\frac{1}{2} = \frac{3}{2}$  then  $\frac{3}{2} \times \frac{1}{10} = \frac{3}{20}$  £. Ans.

9. At  $\frac{1}{2}$  of a pound, per yard, what cost  $7\frac{1}{2}$  yards?

Ans.  $6\frac{1}{4}$  £.

10. At  $\$2\frac{1}{2}$  per yard, what cost  $6\frac{1}{2}$  yards? Ans.  $\$14\frac{3}{4}$ .

11. What is the continued product of 3,  $\frac{2}{3}$ ,  $\frac{1}{2}$  of  $\frac{1}{3}$ ,  $2\frac{1}{2}$ , and  $\frac{1}{2}$  of  $\frac{1}{2}$  of  $\frac{1}{2}$ ?

Ans.  $\frac{3}{2}$ .

¶ 51. The RULE for the multiplication of fractions may now be presented at one view:

I. To multiply a fraction by a whole number, or a whole number by a fraction, —Divide the denominator by the whole number, when it can be done without a remainder; otherwise, multiply the numerator by it, and under the product write the denominator, which may then be reduced to a whole or mixed number.

II. To multiply a mixed number by a whole number, —Multiply the fraction and integers, separately, and add their products together.

III. To multiply one fraction by another, —Multiply together

er the numerators for a new numerator, and the denominators for a new denominator.

*Note.* If either or both are mixed numbers, they may first be reduced to improper fractions.

### EXAMPLES FOR PRACTICE.

1. At  $\frac{1}{2}$ £ per yard, what cost 4 yards of cloth? — 5 yds. — 6 yds. — 8 yds. — 20 yds.

*Ans.* to the last, 15£.

2. Multiply 148 by  $\frac{1}{2}$ . — by  $\frac{1}{4}$ . — by  $\frac{3}{8}$ . — by  $\frac{5}{16}$ .

*Last product, 44 $\frac{1}{16}$ .*

3. If 2  $\frac{1}{10}$  tons of hay keep 1 horse through the winter, how much will it take to keep 3 horses the same time? — 7 horses? — 13 horses? *Ans.* to the last, 37 $\frac{7}{10}$  tons.

4. What will 8 $\frac{1}{12}$  barrels of cider come to, at 7 shillings per barrel?

5. At 14 $\frac{1}{2}$ £ per cwt. what will be the cost of 147 cwt.?

6. A owned  $\frac{2}{5}$  of a note; B owned  $\frac{3}{10}$  of the same; the note amounted to 1000£; what was each one's share of the money?

7. Multiply  $\frac{1}{2}$  of  $\frac{3}{4}$  by  $\frac{2}{3}$  of  $\frac{1}{2}$ .

*Product,  $\frac{1}{4}$ .*

8. Multiply 7 $\frac{1}{2}$  by 2 $\frac{1}{15}$ .

*Product, 15 $\frac{1}{2}$ .*

9. Multiply  $\frac{3}{4}$  by 2 $\frac{3}{4}$ .

*Product, 2 $\frac{1}{4}$ .*

10. Multiply  $\frac{1}{2}$  of 6 by  $\frac{3}{4}$ .

*Product, 1.*

11. Multiply  $\frac{1}{2}$  of 2 by  $\frac{1}{3}$  of 4.

*Product, 3.*

12. Multiply continually together  $\frac{1}{2}$  of 8,  $\frac{2}{3}$  of 7,  $\frac{3}{4}$  of 9, and  $\frac{1}{5}$  of 10.

*Product, 20.*

13. Multiply 1000000 by  $\frac{1}{2}$ .

*Product, 555555 $\frac{1}{2}$ .*

*To divide a whole number by a fraction.*

¶ 52. We have already shown, (¶ 46,) how to divide a fraction by a whole number; we now proceed to show how to divide a whole number by a fraction.

1. A man divided 9£. among some poor people, giving them  $\frac{1}{3}$  of a pound each; how many were the persons who received the money?  $9 \div \frac{1}{3} =$  how many?

1 pound is  $\frac{1}{3}$ , and 9 pounds is 9 times as many, that is, 9 $\times 3$ ; then  $\frac{1}{3}$  is contained in 9 $\times 3$  as many times as 3 is contained in 36.

*Ans.* 12 persons,

That is, — *Multiply the dividend by the denominator of the dividing fraction, (thereby reducing the dividend to parts of the same magnitude as the divisor,) and divide the product by the numerator.*



the denomina-  
they may first

cloth 1 — 5  
the last, 15£.  
by 10  
product, 44  
the winter  
ne time 1—  
st, 37  $\frac{7}{10}$  tons.  
at 7 shillings

of 147 cwt. 1  
he same; the  
share of the

Product. 1.  
Product. 15  $\frac{1}{2}$ .  
Product. 2  $\frac{1}{2}$ .  
Product, 1.  
Product, 3.  
of 7,  $\frac{1}{2}$  of 9,  
Product, 20.  
ct, 555555  $\frac{1}{2}$ .

w to divide a  
to show how  
ple, giving  
persons who  
any, that is,  
as 3 is con-  
12 persons,  
nator of the  
end to parts  
side the pro-

2. How many times is  $\frac{1}{2}$  contained in  $818 \div \frac{1}{2}$  = how many ?

OPERATION.

8 Dividend.

5 Denominator,

Numerator, 3)40

Quotient, 13  $\frac{1}{2}$  times, the answer.

To multiply by a fraction, we have seen, (¶ 49,) implies two operations—a *division* and a *multiplication*; so, also, to divide by a fraction implies two operations—a *multiplication* and a *division*.

¶ 53. Division is the reverse of multiplication.

<p>To multiply by a fraction, whether the multiplicand be a whole number or a fraction, as has already been shown, (¶ 49,) we divide by the denominator of the multiplying fraction, and multiply the quotient by the numerator.</p>	<p>To divide by a fraction, whether the dividend be a whole number or a fraction, we multiply by the denominator of the dividing fraction, and divide the product by the numerator.</p>
--	---

*Note.* In either case, it is matter of indifference, as it respects the result, which of these operations precedes the other; but in *practice* it will frequently be more convenient, that the multiplication precede the division.

12 multiplied by $\frac{1}{2}$ , the product is 9.	12 divided by $\frac{1}{2}$ , the quotient is 16.
--	---

<p>In multiplication, the multiplier being <i>less</i> than unity, or 1, will require the product to be <i>less</i> than the multiplier; consequently will require the quotient to be <i>greater</i> than the dividend, to which it will be equal when the multiplier is 1, and greater when the multiplier is <i>more</i> than 1.</p>	<p>In division, the divisor being <i>less</i> than unity, or 1, will be contained a <i>greater</i> number of times; consequently will require the quotient to be <i>greater</i> than the dividend, to which it will be equal when the divisor is 1, and <i>less</i> when the divisor is <i>more</i> than 1.</p>
--	---

EXAMPLES FOR PRACTICE.

1. How many times is  $\frac{1}{2}$  contained in 7?  $7 \div \frac{1}{2}$  = how many ?



2. How many times can I draw  $\frac{1}{2}$  of a gallon of wine out of a cask containing 26 gallons ?

3. Divide 3 by  $\frac{1}{2}$ . — 6 by  $\frac{1}{3}$ . — 10 by  $\frac{1}{4}$ .

4. If a man drink  $\frac{1}{6}$  of a quart of rum a day, how long will 3 gallons last him ?

5. If  $2\frac{1}{2}$  bushels of oats sow an acre, how many acres will 22 bushels sow ?  $22 \div 2\frac{1}{2} =$  how many times ?

*Note.* Reduce the mixed number to an improper fraction,  $2\frac{1}{2} = \frac{5}{2}$ . *Ans.* 8 acres.

6. At  $1\frac{1}{2}$  £ a yard, how many yards of cloth may be bought for 37 £ ? *Ans.* 26  $\frac{2}{3}$  yards.

7. How many times  $\frac{8}{103}$  contained in 84 ?

*Ans.* 90  $\frac{1}{3}$  times.

8. How many times is  $\frac{3}{5}$  contained in 6 ?

*Ans.*  $\frac{10}{3}$  of 1 time.

9. How many times is  $8\frac{1}{2}$  contained in 53 ?

*Ans.* 6  $\frac{1}{2}$  times.

10. At  $\frac{2}{3}$  of a pound for building 1 rod of stone wall, how many rods may be built for 87 £ ?  $87 \div \frac{2}{3} =$  how many times ?

*To divide one fraction by another.*

¶ 54. 1. At  $\frac{2}{3}$  of a pound per barrel, how much rye may be bought for  $\frac{1}{3}$  of a pound ?  $\frac{1}{3}$  is contained in  $\frac{2}{3}$  how many times ?

Had the rye been 2 whole pounds per barrel, instead of  $\frac{2}{3}$  of a pound, it is evident, that  $\frac{2}{3}$  of a pound must have been divided by 2, and the quotient would have been  $\frac{1}{3}$ ; but the divisor is 3ds, and 3ds will be contained 3 times where a like number of whole ones are contained 1 time; consequently the quotient  $\frac{1}{3}$  is 3 times too small, and must therefore, in order to give the true answer, be multiplied by 3, that is, by the denominator of the divisor; 3 times  $\frac{1}{3} = \frac{1}{1}$  barrel, answer.

The process is that already described, ¶ 52 and ¶ 53. If carefully considered, it will be perceived, that the *numerator* of the divisor is multiplied into the denominator of the dividend, and the *denominator* of the divisor into the *numerator* of the dividend; wherefore, in practice, it will be more convenient to *invert the divisor*; thus,  $\frac{2}{3}$  inverted becomes  $\frac{3}{2}$ ; then multiply together the two upper terms for a numerator, and the two lower terms for a denominator, as in the multi-

on of wine

2.  
how long

many acres  
mes 1

proper frac-

ns. 8 acres.

th may be

26½ yards.

90½ times.

½ of 1 time.

. 6½ times.

stone wall,

÷ ½ = how

uch rye may

h ½ how ma-

instead of ¾

st have been

een 1½; but

times where

time; conse-

, and must

be multiplied

or; 3 times

nd ¶ 53. If

the numerator

r of the divi-

the numerator

be more con-

becomes ¾;

the numerator.

n the multi-

plication of one fraction by another. Thus, in the above example,  $3 \times 3 = 9$ , as before.

$$\frac{2}{2} \times \frac{5}{5} = \frac{10}{10}$$

### EXAMPLES FOR PRACTICE.

2. At  $\frac{1}{4}$  of a pound per bushel for wheat, how many bushels may be bought for  $\frac{1}{2}$  of a pound? How many times is  $\frac{1}{4}$  contained in  $\frac{1}{2}$ ? *Ans.*  $3\frac{1}{2}$  bushels.

3. If  $\frac{1}{4}$  of a yard of cloth cost  $\frac{1}{2}$  of a pound, what is that per yard? It will be recollected, (¶ 24,) that when the cost of any quantity is given to find the price of a unit, we divide the cost by the quantity. Thus,  $\frac{1}{2}$  (the cost) divided by  $\frac{1}{4}$  (the quantity) will give the price of 1 yard.

*Ans.*  $3\frac{1}{2}$  of a pound per yard.

**Proof.** If the work be right, (¶ 16, "Proof,") the product of the quotient into the divisor will be equal to the dividend; thus,  $3\frac{1}{2} \times \frac{1}{4} = \frac{1}{2}$ . This, it will be perceived, is multiplying the price of one yard ( $3\frac{1}{2}$ ) by the quantity ( $\frac{1}{4}$ ) to find the cost ( $\frac{1}{2}$ ); and is, in fact, reversing the question, thus, if the price of 1 yard be  $3\frac{1}{2}$  of a pound, what will  $\frac{1}{4}$  of a yard cost? *Ans.*  $\frac{1}{2}$  of a pound.

**Note.** Let the pupil be required to reverse and prove the succeeding examples in the same manner.

4. How many bushels of wheat, at  $\frac{1}{16}$  of a pound per bushel, may be bought for  $\frac{1}{2}$  of a pound? *Ans.*  $4\frac{1}{2}$  bushels.

5. If  $4\frac{1}{2}$  pounds of butter serve a family 1 week, how many weeks will  $36\frac{1}{2}$  pounds serve them?

The mixed numbers, it will be recollected, may be reduced to improper fractions.

*Ans.*  $8\frac{1}{4}$  weeks.

6. Divide  $\frac{1}{2}$  by  $\frac{1}{4}$ . Quot. 1

Divide  $\frac{1}{2}$  by  $\frac{1}{4}$ . Quot. 2.

7. Divide  $\frac{1}{2}$  by  $\frac{1}{4}$ . Quot. 3

Divide  $\frac{1}{2}$  by  $\frac{1}{4}$ . Quot.  $3\frac{1}{2}$ .

8. Divide  $2\frac{1}{4}$  by  $1\frac{1}{4}$ .

Divide  $10\frac{1}{4}$  by  $2\frac{1}{4}$ .

Quot.  $1\frac{1}{2}$ .

Quot.  $4\frac{1}{4}$ .

9. How many times  $\frac{1}{16}$  contained in  $\frac{1}{2}$ ? *Ans.* 4 times.

10. How many times is  $\frac{1}{4}$  contained in  $4\frac{1}{4}$ ? *Ans.*  $11\frac{1}{2}$  times.

11. Divide  $\frac{1}{2}$  of  $\frac{1}{4}$  by  $\frac{1}{4}$  of  $\frac{1}{4}$ . Quot. 4.

¶ 55. The RULE for division of fractions may now be presented at one view:—

1. To divide a fraction by a whole number,—Divide the

numerator by the whole number, when it can be done without a remainder, and under the quotient write the denominator; otherwise, multiply the denominator by it, and over the product write the numerator.

II. To divide a whole number by a fraction.—Multiply the dividend by the denominator of the fraction, and divide the product by the numerator.

III. To divide one fraction by another.—Invert the divisor, and multiply together the two upper terms for a numerator, and the two lower terms for a denominator.

Note. If either or both are mixed numbers, they may be reduced to improper fractions.

### EXAMPLES FOR PRACTICE.

1. If 7 lb. of tobacco cost  $\frac{83}{100}$  of a pound, what is it per pound?  $\frac{83}{100} \div 7 =$  how much?  $\frac{1}{7}$  of  $\frac{83}{100}$  is how much?

2. At  $\frac{1}{4}\text{£}$ . for  $\frac{2}{3}$  of a barrel of cider, what is that per barrel?

3. If 4 pounds of sugar cost  $\frac{3}{17}$  of a pound, what does 1 pound cost?

4. If  $\frac{7}{8}$  of a yard cost 13s. what is the price per yard?

5. If 14 $\frac{3}{4}$  yards cost 43 $\text{£}$ . what is the price per yard?

Ans. 2 $\frac{11}{14}$ .

6. At 4 $\frac{1}{2}$  pounds for 10 $\frac{1}{2}$  barrels of cider, what is that per barrel?

Ans. 3 $\text{£}$ .

7. How many times is  $\frac{3}{8}$  contained in 746? Ans. 1969 $\frac{1}{8}$ .

8. Divide  $\frac{1}{2}$  of  $\frac{2}{3}$  by  $\frac{1}{4}$ .

Divide  $\frac{2}{3}$  by  $\frac{1}{4}$  of  $\frac{2}{3}$

Quot.  $\frac{2}{3}$ .

Quot. 3 $\frac{2}{3}$ .

9. Divide  $\frac{1}{2}$  of  $\frac{2}{3}$  by  $\frac{2}{3}$  of  $\frac{2}{3}$ .

Quot.  $\frac{1}{2}$ .

10. Divide  $\frac{1}{2}$  of 4 by  $\frac{1}{15}$ .

Quot. 3.

11. Divide 4 $\frac{3}{4}$  by  $\frac{1}{2}$  of 4.

Quot. 2 $\frac{1}{2}$ .

12. Divide  $\frac{3}{4}$  of 4 by 4 $\frac{3}{4}$ .

Quot.  $\frac{2}{11}$ .

### ADDITION AND SUBTRACTION OF FRACTIONS.

T 56. 1. A boy gave to one of his companions  $\frac{1}{4}$  of an orange, to another  $\frac{1}{8}$ , to another  $\frac{1}{8}$ ; what part of an orange did he give to all?  $\frac{1}{4} + \frac{1}{8} + \frac{1}{8} =$  how much? Ans.  $\frac{1}{2}$ .

2. A cow consumes, in 1 month,  $\frac{1}{5}$  of a ton of hay; a horse, in the same time, consumes  $\frac{1}{10}$  of a ton; and a pair of oxen  $\frac{1}{20}$ ; how much do they all consume? how much more does the horse consume than the cow? — the oxen

than the horse?  $\frac{1}{2} + \frac{1}{3} + \frac{1}{6} =$  how much?  $\frac{1}{2} - \frac{1}{3} =$  how much?  $\frac{1}{2} - \frac{1}{6} =$  how much?

3.  $\frac{1}{2} + \frac{1}{3} + \frac{1}{6} =$  how much?  $\frac{1}{2} - \frac{1}{3} =$  how much?

4.  $\frac{1}{20} + \frac{1}{20} + \frac{1}{20} + \frac{1}{20} + \frac{1}{20} =$  how much?  $\frac{1}{20} - \frac{1}{20} =$  how much?

5. A boy, having  $\frac{3}{4}$  of an apple, gave  $\frac{1}{4}$  of it to his sister; what part of the apple had he left?  $\frac{3}{4} - \frac{1}{4} =$  how much?

When the denominators of two or more fractions are alike (as in the foregoing examples,) they are said to have a common denominator. The parts are then in the same denomination, and, consequently, of the same magnitude or value. It is evident, therefore, that they may be added or subtracted, by adding or subtracting their numerators, that is, the number of their parts, care being taken to write under the result their proper denominator. Thus,  $\frac{1}{7} + \frac{1}{7} = \frac{2}{7}$ ;  $\frac{3}{7} - \frac{1}{7} = \frac{2}{7}$ .

6. A boy, having an orange, gave  $\frac{2}{3}$  of it to his sister, and  $\frac{1}{3}$  to his brother; what part of the orange did he give away?

4ths and 8ths, being parts of different magnitudes, or value, cannot be added together. We must therefore first reduce them to parts of the same magnitude, that is, to a common denominator.  $\frac{2}{3}$  are 3 parts. If each of these parts be divided into 2 equal parts, that is, if we multiply both terms of the fraction  $\frac{2}{3}$  by 2, (§ 43,) it will be changed to  $\frac{4}{6}$ ; then  $\frac{4}{6}$  and  $\frac{1}{6}$  are  $\frac{5}{6}$ . *Ans.*  $\frac{5}{6}$  of an orange.

7. A man had  $\frac{2}{3}$  of a hogshead of molasses in one cask, and  $\frac{1}{3}$  of a hogshead in another; how much more in one cask than in the other?

Here, 3ds cannot be so divided as to become 5ths, nor can 5ths be so divided as to become 3ds; but if the 3ds be each divided into 5 equal parts, and the 5ths each into 3 equal parts, they will all become 15ths. The  $\frac{2}{3}$  will become  $\frac{10}{15}$ , and the  $\frac{1}{3}$  will become  $\frac{5}{15}$ ; then,  $\frac{10}{15}$  taken from  $\frac{10}{15}$  leaves  $\frac{5}{15}$ . *Ans.*

¶ 57. From the very process of dividing each of the parts, that is, of increasing the denominators by multiplying them, it follows, that each denominator must be a factor of the common denominator; now, multiplying all the denominators together will evidently produce such a number.

Hence,—To reduce fractions of different denominators to equivalent fractions, having a common denominator,—**RULE:** Multiply together all the denominators for a common denominator; and, as by this process each denominator is multiplied by all the others, so, to retain the value of each fraction, multiply each numerator by all the denominators, except its own, for a new numerator, and under it write the common denominator.

### EXAMPLES FOR PRACTICE.

1. Reduce  $\frac{3}{4}$ ,  $\frac{2}{5}$  and  $\frac{1}{6}$  to fractions of equal value, having a common denominator.

$3 \times 4 \times 5 = 60$ , the common denominator.

$2 \times 4 \times 5 = 40$ , the new numerator for the first fraction.

$3 \times 3 \times 5 = 45$ , the new numerator for the second fraction.

$3 \times 4 \times 4 = 48$ , the new numerator for the third fraction.

The new fractions, therefore, are  $\frac{40}{60}$ ,  $\frac{45}{60}$ , and  $\frac{48}{60}$ . By an inspection of the operation, the pupil will perceive, that the numerator and denominator of each fraction have been multiplied by the same numbers; consequently, (§ 43,) that their value has not been altered.

3. Reduce to equivalent fractions of a common denominator, and add together,  $\frac{1}{2}$ ,  $\frac{2}{3}$ , and  $\frac{1}{6}$ .

*Ans.*  $\frac{30}{60} + \frac{40}{60} + \frac{10}{60} = \frac{80}{60} = 1\frac{2}{3}$ , amount.

4. Add together  $\frac{1}{2}$  and  $\frac{2}{3}$ .

*Amount,*  $1\frac{2}{3}$ .

5. What is the amount of  $\frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5}$ ?

*Ans.*  $\frac{34}{60} = 1\frac{17}{30}$ .

6. What are the fractions of a common denominator equivalent to  $\frac{1}{2}$  and  $\frac{2}{3}$ ? *Ans.*  $\frac{12}{24}$  and  $\frac{16}{24}$ , or  $\frac{8}{12}$  and  $\frac{16}{12}$ .

We have already seen, (§ 56, ex. 7,) that the common denominator may be any number, of which each given denominator is a factor, that is, any number which may be divided by each of them without a remainder. Such a number is called a common multiple of all its common divisors, and the least number that will do this is called their least common multiple; therefore, the least common denominator of any fractions is the least common multiple of all their denominators. Though the rule already given will always find a common multiple of the given denominators, yet it will not always find their least common multiple. In the last example, 24 is evidently a common multiple of 4 and 6, for it will exactly measure both of them; but 12 will do the same, and as 12 is

the *least* number that will do this, it is the *least* common multiple of 4 and 6. It will therefore be convenient to have a *rule* for finding this least common multiple. Let the numbers be 4 and 6.

It is evident, that one number is a multiple of another, when the former contains all the factors of the latter. The factors of 4 are 2 and 2, ( $2 \times 2 = 4$ .) The factors of 6 are 2 and 3, ( $2 \times 3 = 6$ .) Consequently,  $2 \times 2 \times 3 = 12$  contains the factors of 4 that is,  $2 \times 2$ ; and also contains the factors of 6, that is,  $2 \times 3$ . 12, then, is a common multiple of 4 and 6 and it is the *least* common multiple, because it does not contain *any* factor, except those which make up the numbers 4 and 6; nor either of those repeated more than is necessary to produce 4 and 6. Hence it follows, that when any two numbers have a factor common to both, it may be once omitted; thus, 2 is a factor common both to 4 and 6, and is consequently once omitted.

7 58. On this principle is founded the *RULE* for finding the *least common multiple of two or more numbers*. Write down the numbers in a line, and divide them by any number that will measure two or more of them; and write the quotients and undivided numbers in a line beneath. Divide this line as before, and so on, until there are no two numbers that can be measured by the same divisor; then the continual product of all the divisors and numbers in the last line will be the least common multiple required.

Let us apply the rule to find the least common multiple of 4 and 6.

	4 and 6 may both be measured by 2; the
2) 4 . 6	quotients are 2 and 3. There is no number greater
—	than 1, which will measure 2 and 3. There-
2 . 3	fore, $2 \times 2 \times 3 = 12$ is the least common mul-
	tiply of 4 and 6.

If the pupil examine the process, he will see that the divisor 2 is a factor common to 4 and 6, and that dividing 4 by this factor gives for a quotient its other factor, 2. In the same manner, dividing 6 gives its other factor, 3. Therefore the divisor and quotients make up all the factors of the two numbers, which, multiplied together, must give the common multiple.

7. Reduce  $\frac{2}{3}$ ,  $\frac{1}{2}$ ,  $\frac{3}{4}$  and  $\frac{1}{6}$  to equivalent fractions of the least common denominator.

$$\begin{array}{r} \text{OPERATION} \\ 2) 4 . 2 . 3 . 6 \\ 3) 2 . 1 . 3 . 3 \\ \hline 2 . 1 . 1 . 1 \end{array}$$

Then,  $2 \times 3 \times 2 = 12$ , least common denominator. It is evident we need not multiply by the 1s, as this would not alter the number.

To find the new numerators that is, how many 12ths each fraction is, we may take  $\frac{2}{3}$ ,  $\frac{1}{2}$ ,  $\frac{3}{4}$ , and  $\frac{1}{6}$  of 12, Thus:

$$\left. \begin{array}{l} \frac{2}{3} \text{ of } 12 = 8 \\ \frac{1}{2} \text{ of } 12 = 6 \\ \frac{3}{4} \text{ of } 12 = 9 \\ \frac{1}{6} \text{ of } 12 = 2 \end{array} \right\} \begin{array}{l} \text{New numerators, which,} \\ \text{written over the com-} \\ \text{mon denominators, give} \end{array} \left\{ \begin{array}{l} \frac{8}{12} = \frac{2}{3} \\ \frac{6}{12} = \frac{1}{2} \\ \frac{9}{12} = \frac{3}{4} \\ \frac{2}{12} = \frac{1}{6} \end{array} \right.$$

*Ans.*  $\frac{8}{12}$ ,  $\frac{6}{12}$ ,  $\frac{9}{12}$ , and  $\frac{2}{12}$ .

8. Reduce  $\frac{1}{2}$ ,  $\frac{3}{4}$ , and  $\frac{1}{6}$  to fractions having the least common denominator, and add them together.

*Ans.*  $\frac{1}{2} + \frac{3}{4} + \frac{1}{6} = \frac{3}{4} + \frac{1}{2} = \frac{5}{4} = 1\frac{1}{4}$ , amount.

9. Reduce  $\frac{1}{2}$  and  $\frac{1}{3}$  to fractions of the least common denominator, and subtract one from the other.

*Ans.*  $\frac{1}{2} - \frac{1}{3} = \frac{1}{6}$ , difference.

10. What is the least number that 3, 5, 8 and 10 will measure?

*Ans.* 120.

11. There are 3 pieces of cloth, one containing  $7\frac{3}{4}$  yards, another  $13\frac{3}{4}$  yards, and the other  $15\frac{3}{4}$  yards; how many yards in the 3 pieces.

Before adding, reduce the fractional parts to their least common denominator; this being done, we shall have,

$$\left. \begin{array}{l} 7\frac{3}{4} = 7\frac{6}{8} \\ 13\frac{3}{4} = 13\frac{6}{8} \\ 15\frac{3}{4} = 15\frac{6}{8} \end{array} \right\} \begin{array}{l} \text{Adding together all the 24ths, viz. } 18 + 20 \\ + 21, \text{ we obtain } 59, \text{ that is, } \frac{59}{24} = 2\frac{11}{24}. \\ \text{We write down the fraction } \frac{11}{24} \text{ under the} \\ \text{other fractions, and reserve the 2 integers} \\ \text{to be carried to the amount of the other in-} \\ \text{tegers, making in the whole } 37\frac{11}{24} \end{array}$$

*Ans.*  $37\frac{11}{24}$

12. There was a piece of cloth containing  $34\frac{3}{4}$  yards, from which were taken  $12\frac{3}{4}$  yards; how much was there left?

$$\begin{array}{r} 34\frac{3}{4} = 34\frac{6}{8} \\ 12\frac{3}{4} = 12\frac{6}{8} \\ \hline \end{array}$$

*Ans.*  $21\frac{1}{2}$  yds.

We cannot take 16 twenty-fourths ( $\frac{16}{24}$ ) from 9 twenty-fourths ( $\frac{9}{24}$ ); we must, therefore, borrow 1 integer, = 24 twenty-fourths, ( $\frac{24}{24}$ ) which, with  $\frac{9}{24}$ , makes  $\frac{33}{24}$ ; we can now take  $\frac{16}{24}$  from  $\frac{33}{24}$ , and there will remain  $\frac{17}{24}$ ; but, as we borrowed, so also



we must carry 1 to the 12, which makes it 13, and 13 from 34 leaves 21. *Ans.*  $21\frac{1}{4}$ .

13. What is the amount of  $\frac{1}{2}$  of  $\frac{2}{3}$  of a yard,  $\frac{2}{3}$  of a yard, and  $\frac{1}{2}$  of 2 yards?

*Note.* The compound fraction may be reduced to a simple fraction; thus,  $\frac{1}{2}$  of  $\frac{2}{3} = \frac{1}{3}$ ; and  $\frac{1}{2}$  of 2 = 1; then,  $\frac{1}{3} + 1 + \frac{2}{3} = 1\frac{1}{3} = 1\frac{2}{3}$  yds., *answer.*

† 59. From the foregoing examples we derive the following RULE:—To add or subtract fractions, add or subtract their numerators, when they have a common denominator; otherwise, they must first be reduced to a common denominator.

*Note.* Compound fractions must be reduced to simple fractions before adding or subtracting.

# EXAMPLES FOR PRACTICE.

1. What is the amount of  $\frac{1}{2}$ ,  $4\frac{1}{2}$  and 12? *Ans.*  $17\frac{1}{2}$ .
2. A man bought a farm, and sold  $\frac{1}{2}$  of  $\frac{1}{2}$  of it; what part of the farm had he left? *Ans.*  $\frac{3}{4}$ .
3. Add together  $\frac{1}{2}$ ,  $\frac{1}{3}$ ,  $\frac{1}{4}$ ,  $\frac{1}{5}$ ,  $\frac{1}{6}$  and  $\frac{1}{7}$ ? *Amount.*  $2\frac{1}{42}$ .
4. What is the difference between  $14\frac{1}{2}$  and  $16\frac{1}{3}$ ? *Ans.*  $1\frac{1}{6}$ .  
*Remainder,*  $\frac{1}{2}$ .
5. From  $1\frac{1}{2}$  take  $\frac{1}{4}$ . *Remainder,*  $1\frac{1}{4}$ .
6. From 3 take  $\frac{1}{2}$ . *Remainder,*  $2\frac{1}{2}$ .
7. From  $147\frac{1}{2}$  take  $48\frac{1}{2}$ . *Rem.*  $99$ .
8. From  $\frac{1}{2}$  of  $\frac{1}{3}$  take  $\frac{1}{4}$  of  $\frac{1}{5}$ . *Rem.*  $\frac{1}{30}$ .
9. Add together  $112\frac{1}{2}$ ,  $311\frac{1}{2}$ , and  $1000\frac{1}{2}$ .
10. Add together 14, 11,  $4\frac{1}{2}$ ,  $\frac{1}{3}$  and  $\frac{1}{4}$ .
11. From  $\frac{2}{3}$  take  $\frac{1}{4}$ . From  $\frac{2}{3}$  take  $\frac{1}{4}$ .
12. What is the difference between  $\frac{1}{2}$  and  $\frac{1}{3}$ ?  $\frac{1}{3}$  and  $\frac{1}{4}$ ?  $\frac{1}{4}$  and  $\frac{1}{5}$ ?  $\frac{1}{5}$  and  $\frac{1}{6}$ ?  $\frac{1}{6}$  and  $\frac{1}{7}$ ?
13. How much is  $1 - \frac{1}{2}$ ?  $1 - \frac{1}{3}$ ?  $1 - \frac{1}{4}$ ?  $1 - \frac{1}{5}$ ?  $2 - \frac{1}{2}$ ?  $2 - \frac{1}{3}$ ?  $3 - \frac{1}{3}$ ?  $1000 - \frac{1}{10}$ ?

# REDUCTION OF FRACTIONS.

† 60. We have seen († 27,) that integers of one denomination may be reduced to integers of another denomination. It is evident, that fractions of one denomination, after the same manner, and by the same rules, may be reduced to fractions of another denomination; that is, fractions, like integers, may be brought into lower denomina-

tions by multiplication, and into higher denominations by division.

To reduce higher into LOWER denominations.

(RULE. See † 28.)

1. Reduce  $\frac{1}{280}$  of a pound to pence, or the fraction of a penny.

*Note.* Let it be recollected that a fraction is multiplied either by dividing its denominator, or by multiplying its numerator.

$$\frac{1}{280} \text{ £.} \times 20 = \frac{1}{14} \text{ s.} \times 12 = \frac{3}{7} \text{ d. Ans.}$$

Or thus:  $\frac{1}{280}$  of 20 of 12 =  $\frac{3}{7}$  of a penny, *Ans.*

2. Reduce  $\frac{1}{1280}$  of a pound to the fraction of a farthing.

$$\frac{1}{1280} \text{ £.} \times 20 = \frac{1}{64} \text{ s.} \times 12 = \frac{3}{16} \text{ d.} \times 4 = \frac{3}{4} \text{ q.}$$

Or thus:

*Num.* 1

20 s. in 1 £.

20

12 d. in 1 s.

240

4 q. in 1 d.

960

Then,  $\frac{1}{1280} = \frac{3}{4} \text{ q. Ans.}$

5. Reduce  $\frac{1}{2688}$  of a guinea to the fraction of a penny.

7. Reduce  $\frac{1}{4}$  of a guinea to the fraction of a pound.

Consult † 28, ex. 12.

9. Reduce  $\frac{1}{4}$  of a moidore, at 1 £. 10s. to the fraction of a guinea.

11. Reduce  $\frac{1}{27}$  of a pound, Troy, to the fraction of an ounce.

To reduce lower into HIGHER denominations.

(RULE. See † 28.)

2. Reduce  $\frac{1}{4}$  of a penny to the fraction of a pound.

*Note.* Division is performed either by dividing the numerator, or by multiplying the denominator.

$$\frac{1}{4} \text{ d.} \div 12 = \frac{1}{48} \text{ s.} \div 20 = \frac{1}{960} \text{ £. Ans.}$$

Or thus:  $\frac{1}{4}$  of  $\frac{1}{12}$  of  $\frac{1}{20}$  =  $\frac{1}{960} \text{ £. Ans.}$

4. Reduce  $\frac{1}{4}$  of a farthing to the fraction of a pound.

$$\frac{1}{4} \text{ q.} \div 4 = \frac{1}{16} \text{ d.} \div 12 = \frac{1}{192} \text{ s.} \div 20 = \frac{1}{3840} \text{ £.}$$

Or thus:

*Denom.* 4

4 q. in 1 d.

16

12 d. in 1 s.

192

20 s. in 1 £.

3840

Then  $\frac{1}{3840} = \frac{1}{1280} \text{ £. Ans.}$

6. Reduce  $\frac{1}{4}$  of a penny to the fraction of a guinea.

8. Reduce  $\frac{1}{4}$  of a pound to the fraction of a guinea.

10. Reduce  $\frac{3}{4}$  of a guinea to the fraction of a moidore.

12. Reduce  $\frac{1}{4}$  of an ounce Troy, to the fraction of a pound

Troy.

nations by

HIGHER

28.)

penny to

und.

s perform-

g the nu-

ultiplying the

. ÷ 20 =

r of 20 =

a farthing

pound.

12 = 132s.

£.

Ans.

penny to

inea.

a pound to

inea.

f a guinea

moidore.

an ounce

a pound

13. Reduce  $\frac{1}{2}$  of a pound, avoirdupois, to the fraction of a pound an ounce.

15. A man has  $\frac{1}{2}$  of a hogshead of wine; what part is that of a pint?

17. A cucumber grew to the length of  $\frac{1}{3500}$  of a mile; what part is that of a foot?

19. Reduce  $\frac{3}{4}$  of  $\frac{1}{2}$  of a pound to the fraction of a shilling.

21. Reduce  $\frac{1}{2}$  of  $\frac{1}{2}$  of 3 pounds to the fraction of a penny.

14. Reduce  $\frac{1}{2}$  of an ounce to the fraction of a pound avoirdupois.

16. A man has  $\frac{1}{2}$  of a pint of wine; what part is that of a hogshead?

18. A cucumber grew to the length of 1 foot 4 inches =  $\frac{13}{4}$  of a foot; what part is that of a mile?

20.  $\frac{3}{4}$  of a shilling is  $\frac{3}{4}$  of what fraction of a pound?

22.  $\frac{1}{100}$  of a penny is  $\frac{1}{100}$  of what fraction of 3 pounds?

$\frac{1}{100}$  of a penny, is  $\frac{1}{100}$  of what part of 3 pounds?  $\frac{1}{100}$  of a penny is  $\frac{1}{100}$  of how many pounds?

¶ 61. It will frequently be required to find the value of a fraction, that is to reduce a fraction of a greater denomination to integers of less denomination.

1. What is the value of  $\frac{3}{4}$  of a pound? In other words, Reduce  $\frac{3}{4}$  of a pound to shillings and pence.

$\frac{3}{4}$  of a £ is  $4^s = 12^d$  shillings; it is evident from  $\frac{1}{4}$  of a shilling may be obtained some pence;  $\frac{1}{4}$  of a shilling is  $3^d = 4^d$ . That is,—Multiply the numerator by that number which will reduce it to the next less denomination, and divide the product by the denominator; if there be a remainder, multiply and divide as before, and so on; the several quotients, placed one after another, in their order, will be the answer.

It will frequently be required to reduce integers to the fraction of a greater denomination.

2. Reduce 13s. 4d. to the fraction of a pound.

13s. 4d. is 160 pence; there are 240 pence in a pound?

therefore, 13s. 4d. is  $\frac{160}{240} = \frac{4}{6} = \frac{2}{3}$  of a pound. That is,—Reduce the given sum or quantity to the least denomination mentioned in it, for a numerator; then reduce an integer to that greater denomination (to a fraction of which it is required to reduce the given sum or quantity) to the same denomination, for a denominator, and they will form the fraction required.

## EXAMPLES FOR PRACTICE.

3. What is the value of  $\frac{3}{4}$  of a shilling?

OPERATION.

$$\begin{array}{r} \text{Numer. } 3 \\ 12 \\ \hline \text{Denom. } 8 \end{array} \quad \begin{array}{r} 36 \\ 32 \\ 4 \\ 4 \end{array} \quad \begin{array}{l} (4d. \text{ } 2q. \text{ } Ans. \\ 16(2q. \\ 16 \end{array}$$

4. Reduce 4d. 2q. to the fraction of a shilling.

OPERATION.

$$\begin{array}{r} 4d. \text{ } 2q. \text{ } 1s. \\ 4 \\ \hline 18 \end{array} \quad \begin{array}{r} \text{Numer. } 12 \\ 4 \\ \hline 48 \end{array} \quad \begin{array}{l} \text{Denom.} \end{array}$$

$$\frac{1}{4} = \frac{3}{12} \text{ } Ans.$$

5. What is the value of  $\frac{3}{4}$  of a pound Troy?

7. What is the value of  $\frac{3}{4}$  of a pound avoirdupois?

6. Reduce 7 oz. 4 pwt. to the fraction of a pound Troy.

8. Reduce 8 oz. 14 $\frac{3}{4}$  dr. to the fraction of a pound avoirdupois.

*Note.* Both the numerator and the denominator must be reduced to 9ths of a dr.

9.  $\frac{3}{4}$  of a month is how many days, hours, and minutes?

11. Reduce  $\frac{3}{4}$  of a mile to its proper quantity.

13. Reduce  $\frac{3}{4}$  of an acre to its proper quantity.

15. What is the value of  $\frac{3}{4}$  of a dollar in shillings, pence, &c.?

17. What is the value of  $\frac{3}{4}$  of a yard?

19. What is the value of  $\frac{3}{4}$  of a ton.

10. 3 weeks, 1d. 9h. 36m. is what fraction of a month?

12. Reduce 4 fur. 125 yds. 2ft. 1 in. 2 $\frac{1}{2}$  bar. to the fraction of a mile.

14. Reduce 1 rood 30 poles to the fraction of an acre.

16. Reduce 4s. 8 $\frac{1}{2}$ d. to the fraction of a dollar.

18. Reduce 2 ft. 8 in. 1 $\frac{1}{2}$ b. to the fraction of a yard.

20. Reduce 4 cwt. 2qr. 12 lb. 14 oz. 12 $\frac{3}{4}$  dr. to the fraction of a ton.

*Note.* Let the pupil be required to reverse and prove the following examples:

21. What is the value of  $\frac{3}{4}$  of a guinea?

22. Reduce 3 roods 17 $\frac{1}{2}$  poles to the fraction of an acre.

23. A man bought 27 gal. 3 qts. 1 pt. of molasses ; what part is that of a hogshead ?

24. A man purchased  $\frac{1}{3}$  of 7 cwt. of sugar ; how much sugar did he purchase ?

25. 13h. 42m. 51s. is what part or fraction of a day ?

# SUPPLEMENT TO FRACTIONS.

1. What are *fractions* ? 2. Whence is it that the parts into which any thing or any number may be divided, take their name ? 3. How are fractions *represented* by figures ? 4. What is the number *above* the line called ? — Why is it so called ? 5. What is the number *below* the line called ? — Why is it so called ? — What does it show ? 6. What is it which determines the *magnitude* of the parts ? — Why ? 7. What is a *simple* or *proper* fraction ? — an *improper* fraction — a *mixed* number ? 8. How is an improper fraction reduced to a whole or mixed number ? 9. How is a mixed number reduced to an improper fraction ? — a whole number ? 10. What is understood by the *terms* of the fraction ? 11. How is a fraction reduced to its most *simple* or *lowest* terms ? 12. What is understood by a common divisor ? — by the greatest common divisor ? 13. How is it found ? 14. How many ways are there to multiply a fraction by a whole number ? 15. How does it appear, that *dividing the denominator multiplies the fraction* ? 16. How is a *mixed* number multiplied ? 17. What is implied in multiplying by a fraction ? 18. Of how many operations does it consist ? — What are they ? 19. When the multiplier is *less* than a unit, what is the product compared with the multiplicand ? 20. How do you multiply a whole number by a fraction ? 21. How do you multiply one fraction by another ? 22. How do you multiply a mixed number by a mixed number ? 23. How does it appear, that in multiplying both terms of the fraction by the same number the value of the fraction is not altered ? 24. How many ways are there to divide a fraction by a whole number ? — What are they ? 25. How does it appear that a *fraction is divided by multiplying its denominator* ? 26. How does *dividing* by a fraction differ from *multiplying* by a fraction ? 27. When the *divisor* is *less* than a unit, what is the quotient compared with the dividend ? 28. What is understood by a *common denominator* ? — the *least common denominator* ? 29. How does it appear, that each *given denominator* must be a factor of the *common denominator* ? 30. How is the common denominator to two or more fractions found ? 31. What is understood by a *multiple* ? — by a *common multiple* ? — by the *least common multiple* ? — What is the process of finding it ? 32. How are fractions added and subtracted ? 33. How is a fraction of a greater denomination reduced to one of a less ? — of a less to a greater ? 34. How are fractions of a greater denomination reduced to integers of a less ? — integers of a less denomination to the fraction of a greater ?

## EXERCISES.

1. What is the amount of  $\frac{1}{2}$  and  $\frac{3}{4}$ ? — of  $\frac{1}{2}$  and  $\frac{3}{4}$ ? — of  $12\frac{1}{2}$ ,  $3\frac{3}{4}$  and  $4\frac{1}{2}$ ? *Ans. to the last,  $20\frac{1}{2}$ .*

2. To  $\frac{1}{2}$  of a pound add  $\frac{3}{4}$  of a shilling. *Amount,  $18\frac{1}{4}$ s.*

*Note.* First reduce both to the same denomination.

3.  $\frac{1}{2}$  of a day added to  $\frac{1}{4}$  of an hour, make how many hours? — what part of a day? *Ans. to the last,  $8\frac{1}{2}$  d.*

4. Add  $\frac{1}{2}$  lb. Troy to  $\frac{1}{12}$  of an oz.

*Amount, 6 oz. 11 pwt. 16 gr.*

5. How much is  $\frac{1}{2}$  less  $\frac{1}{3}$ ?  $\frac{1}{12}$  —  $\frac{1}{3}$  of  $\frac{2}{3}$  of  $\frac{1}{2}$ ?

*Ans. to the last,  $\frac{1}{12}$ .*

6. From  $\frac{1}{2}$  shilling take  $\frac{1}{4}$  of a penny. *Rem.  $5\frac{1}{4}$ d.*

7. From  $\frac{1}{2}$  of an ounce take  $\frac{1}{8}$  of a pwt.

*Rem. 11 pwt. 3 grs.*

8. From 4 days  $7\frac{1}{2}$  hours, take 1 d.  $9\frac{1}{4}$  h.

*Rem. 2 d. 22 h. 20 m.*

9. At  $\frac{1}{2}$ £. per yard, what costs  $\frac{1}{2}$  of a yard of cloth?

**T 62.** The price of unity, or 1, being given, to find the cost of any quantity, either less or more than unity, *multiply the price by the quantity.* On the other hand, the cost of any quantity, either less or more than unity, being given, to find the price of unity, or 1, *divide the cost by the quantity.*

*Ans.  $\frac{1}{2}$ £.*

1. If  $\frac{1}{12}$  lb. of sugar cost  $\frac{1}{15}$  of a shilling, what will  $\frac{2}{3}$  of a pound cost?

This example will require two operations: first, as above, to find the price of 1 lb.; secondly, having found the price of 1 lb. to find the cost of  $\frac{2}{3}$  of a pound.  $\frac{1}{15}$ s.  $\div$   $\frac{1}{12}$  ( $\frac{1}{15}$  of  $\frac{1}{15}$ s.  $\pi$  54) =  $\frac{2}{15}$ s. the price of 1 lb. Then,  $\frac{2}{15}$ s.  $\times$   $\frac{2}{3}$  ( $\frac{2}{3}$  of  $\frac{2}{15}$ s.  $\pi$  50) =  $\frac{4}{15}$ s. = 4d.  $3\frac{4}{15}$  q. the answer.

Or we may reason thus: first to find the price of 1 lb:  $\frac{1}{12}$  lb. costs  $\frac{1}{15}$  s. If we knew what  $\frac{1}{12}$  lb. would cost, we might repeat this 12 times, and the result would be the price of 1 lb.  $\frac{1}{12}$  is 12 parts. If  $\frac{1}{12}$  lb. costs  $\frac{1}{15}$  s. it is evident  $\frac{1}{12}$  lb. will cost  $\frac{1}{12}$  of  $\frac{1}{15}$  =  $\frac{1}{180}$  s. and  $\frac{1}{12}$  lb. will cost 12 times as much, that is,  $\frac{12}{180}$ s. = the price of 1 lb. Then,  $\frac{2}{3}$  of  $\frac{1}{180}$  s. =  $\frac{2}{270}$  s. the cost of  $\frac{2}{3}$  of a pound.  $\frac{2}{270}$ s. = 4d.  $3\frac{4}{15}$  q. as before. This process is called *solving the question by analysis.*

After the same manner let the pupil solve the following questions:

and  $\frac{3}{4}$ ?—

the last,  $20\frac{1}{2}$ .

mount,  $18\frac{1}{2}$ s.

ination.

how many

the last,  $8\frac{3}{4}$  d.

pwt. 16 gr.

$\frac{1}{4}$  lb.

the last,  $\frac{1}{16}$  lb.

Rem.  $5\frac{1}{4}$  d.

pwt. 3 grs.

22 h. 20 m.

of cloth?

n, to find the

unity, multi-

d, the cost of

being given,

the quantity.

Ans.  $\frac{1}{16}$  £,

what will  $\frac{1}{4}$  £

st, as above,

and the price

is  $\frac{1}{16}$  £ ( $\frac{1}{16}$  £

$\times \frac{1}{16}$  £.

the answer.

price of 1 lb :

ould cost, we

ould be the

is. it is ev-

lb. will cost

lb. Then,

nd.  $28\frac{1}{2}$  s.

solving the

e following

2. If 7 lb. of tobacco cost  $\frac{1}{4}$  of a pound, what is that a pound?  $\frac{1}{4}$  of  $\frac{1}{4}$  = how much? What is it for 4 lb.?  $\frac{1}{4}$  of  $\frac{1}{4}$  = how much? What for 12 lb.?  $\frac{1}{4}$  of  $\frac{1}{4}$  = how much?

Ans. to the last,  $1\frac{1}{2}$  £.

3. If  $6\frac{1}{2}$  yds of cloth cost 3 £. what cost  $9\frac{1}{2}$  yards?

Ans. 4 £. 5s.  $4\frac{1}{2}$  d.

4. If 2 oz. of silver cost 11s. 3d. what costs  $\frac{1}{4}$  oz.?

Ans. 4s. 2d.  $2\frac{1}{2}$  q.

5. If  $\frac{1}{4}$  oz. costs 4s. 1d. what costs 1 oz. Ans. 5s. 8d.

6. If  $\frac{1}{4}$  lb. less by  $\frac{1}{8}$  costs  $13\frac{1}{2}$  d. what costs 14 lb. less by  $\frac{1}{4}$  of 2 lb.?

Ans. 4 £. 9s.  $9\frac{3}{4}$  d.

7. If  $\frac{1}{4}$  yd cost  $\frac{1}{4}$  £., what will  $40\frac{1}{2}$  yds. cost.

Ans. 59 £. 1s.  $2\frac{1}{2}$  d.

8. If  $\frac{1}{16}$  of a ship costs 251 £. what is  $\frac{1}{32}$  of her worth?

Ans. 53 £. 15s.  $8\frac{1}{2}$  d.

9. At  $3\frac{1}{2}$  £. per cwt. what will  $9\frac{3}{4}$  lb. cost?

10. A merchant, owning  $\frac{1}{4}$  of a vessel, sold  $\frac{1}{8}$  of his share for 39 £. 5s. what was the vessel worth?

Ans. 448 £. 11s.  $10\frac{1}{2}$  d.

11. If  $\frac{1}{8}$  yds. cost  $\frac{1}{4}$  £. what will  $\frac{1}{16}$  of an ell Eng. cost.

Ans. 17s. 1d.  $2\frac{1}{2}$  q.

12. A merchant bought a number of bales of cloth, each containing  $129\frac{1}{4}$  yards, at the rate of 7 £ for 5 yards, and sold them out at the rate of 11 £ for 7 yards, and gained 200 £ by the bargain; how many bales were there?

First find for what he sold 5 yards; then what he gained on 5 yards—what he gained on 1 yard. Then, as many times as the sum gained on 1 yd. is contained in 200 £, so many yards there must have been. Having found the number of yards reduce them to bales.

Ans. 9 bales.

13. If a staff,  $5\frac{1}{2}$  ft. in length, cast a shadow of 6 ft. how high is that steeple whose shadow measures 153 feet?

Ans.  $144\frac{1}{2}$  feet.

14. If 16 men finish a piece of work in  $28\frac{1}{2}$  days, how long will it take 12 men to do the same work?

First find how long it would take 1 man to do it; then 12 men will do it in  $\frac{1}{12}$  of that time.

Ans.  $37\frac{1}{2}$  days.

15. How many pieces of merchandise, at  $20\frac{1}{2}$ s. apiece, must be given for 240 pieces, at  $12\frac{1}{2}$ s. apiece? Ans.  $149\frac{1}{16}$  T.

16. How many yards of backing that is  $1\frac{1}{2}$  yd. wide will be sufficient to line 20 yds. of camlet that is  $\frac{1}{4}$  of a yard wide?



First find the contents of the camlet in square measure ; then it will be easy to find how many yards in length of bocking that is  $1\frac{1}{2}$  yd. wide it will take to make the same quantity.

Ans. 12 yards of camlet.

17. If  $1\frac{1}{2}$  yd. in breadth require  $20\frac{1}{2}$  yds. in length to make a cloak, what in length that is  $\frac{3}{4}$  yd. wide will be required to make the same ?

Ans.  $34\frac{1}{2}$  yds.

18. If 7 horses consume  $2\frac{1}{2}$  tons of hay in 6 weeks, how many tons will 12 horses consume in 8 weeks ?

If we knew how much 1 horse consumed in 1 week, it would be easy to find how much 12 horses would consume in 8 weeks.

$2\frac{1}{2} = \frac{5}{2}$  tons. If 7 horses consume  $\frac{5}{2}$  tons in 6 weeks, 1 horse will consume  $\frac{1}{7}$  of  $\frac{5}{2} = \frac{5}{14}$  of a ton in 6 weeks ; and if a horse consume  $\frac{5}{14}$  of a ton in 6 weeks, he will consume  $\frac{1}{6}$  of  $\frac{5}{14} = \frac{5}{84}$  of a ton in 1 week. 12 horses will consume 12 times  $\frac{5}{84} = \frac{5}{7}$  in 1 week, and in 8 weeks they will consume 8 times  $\frac{5}{7} = \frac{40}{7} = 5\frac{5}{7}$  tons, answer.

19. A man with his family, which in all were 5 persons, did usually drink  $7\frac{1}{2}$  gallons of cider in 1 week ; how much will they drink in  $22\frac{1}{2}$  weeks when 3 persons more are added to the family ?

Ans.  $280\frac{1}{2}$  gallons.

20. If 9 students spend  $10\frac{1}{2}\text{£}$  in 18 days, how much will 20 students spend in 30 days ?

Ans.  $39\text{£}$  18s.  $4\frac{3}{4}\text{d}$ .

## DECIMAL FRACTIONS.

† 63. We have seen, that an individual thing or number may be divided into any number of equal parts, and that these parts will be called halves, thirds, fourths, fifths, sixths, &c., according to the number of parts into which the thing or number may be divided ; and that each of these parts may be again divided into any other number of equal parts, and so on. Such are called *common* or *vulgar fractions*. Their denominators are not uniform, but vary with every varying division of a unit. It is this circumstance which occasions the chief difficulty in the operations to be performed on them ; for when numbers are divided into different kinds or parts, they cannot be so easily compared. This difficulty led to the invention of *decimal fractions*, in which an individual thing, or number, is supposed to be divided first into *ten* equal parts,

re measure ;  
in length of  
ke the same  
ds of camlet.  
n length to  
e will be re-  
ns. 34½ yds.  
weeks, how  
1 week, it  
ould consume  
in 6 weeks,  
weeks ; and  
will consume  
will consume  
eks they will  
r.  
re 5 persons,  
; how much  
ns more are  
280½ gallons.  
ow much will  
£ 18s. 4d.

PS.

ning or num-  
arts, and that  
fifths, sixths,  
ich the thing  
se parts may  
parts, and so  
their denom-  
ying division  
ons the chief  
n them ; for  
r parts, they  
ed to the in-  
ual thing, or  
equal parts,

which will be *tenths* ; and each of these parts to be again di-  
vided into ten *other* equal parts, which will be *hundredths* ;  
and each of these parts to be still further divided into ten  
other equal parts, which will be *thousandths* ; and so on.  
Such are called *decimal fractions*, (from the Latin word *decem*,  
which signifies *ten*,) because they increase and decrease, in a  
*tenfold* proportion, in the same manner as whole numbers.

7 64. In this way of dividing a unit, it is evident, that  
the denominator to a decimal fraction will always be 10,  
100, 1000, or 1 with a number of ciphers annexed ; conse-  
quently, the denominator to a decimal fraction need not be  
expressed, for the numerator only, written with a point  
before it (·) called the *separatrix*, is sufficient of itself to  
express the true value. Thus,

$\frac{6}{10}$  are written '6.  
 $\frac{27}{100}$  .... '27.  
 $\frac{685}{1000}$  .... '685.

The denominator to a decimal fraction, although not ex-  
pressed, is always understood, and is 1 with as many ci-  
phers annexed as there are places in the numerator. Thus,  
'3765 is a decimal consisting of four places ; consequently,  
1 with four ciphers annexed (10000) is its proper denomina-  
tor. Any decimal may be expressed in the form of a com-  
mon fraction by writing under it its proper denominator.  
Thus, '3765 expressed in the form of a common fraction,  
is  $\frac{3765}{10000}$ .

When whole numbers and decimals are expressed to-  
gether, in the same number, it is called a *mixed number*.  
Thus, 25·63 is a mixed number, 25, or all the figures on the  
left hand of the decimal point, being whole numbers, and  
'63, or all the figures on the right hand of the decimal point,  
being decimals.

The names of the places to ten-millionths, and, generally,  
how to read or write decimal fractions, may be seen from  
the following



hand, and decimals continually *decreasing*, in the same proportion, towards the right hand. But as decimals decrease towards the right hand, it follows of course, that they increase towards the left hand, in the same manner as whole numbers.

¶ 65. The value of every figure is determined by its place from *units*. Consequently, ciphers placed at the *right hand* of decimals do *not* alter their value, since every significant figure continues to possess the same place from unity. Thus, '5, '50, '500 are all of the same value, each being equal to  $\frac{5}{10}$ , or  $\frac{1}{2}$ .

But every cipher, placed at the *left hand* of decimal fractions, *diminishes* them tenfold, by removing the significant figures further from unity, and consequently making each part ten times as small. Thus, '5, '05, '005, are of different value, '5 being equal to  $\frac{5}{10}$ , or  $\frac{1}{2}$ ; '05 being equal to  $\frac{5}{100}$ , or  $\frac{1}{20}$ ; and '005 being equal to  $\frac{5}{1000}$ , or  $\frac{1}{200}$ .

Decimal fractions, having *different denominators*, are readily reduced to a *common denominator*, by annexing ciphers until they are equal in number of places. Thus, '5, '06, '234 may be reduced to '500, '060, '234, each of which has 1000 for common denominator.

¶ 66. Decimals are read in the same manner as whole numbers, giving the name of the lowest denomination, or right hand figure, to the whole. Thus, '6853 (the lowest denomination, or right hand figure, being ten-thousandths) is read, 6853 ten-thousandths.

Any whole number may evidently be reduced to decimal parts, that is, to tenths, hundredths, thousandths, &c. by annexing ciphers. Thus, 25 is 250 tenths, 2500 hundredths, 25000 thousandths, &c. Consequently, any mixed number may be read together giving it the name of the lowest denomination or right hand figure. Thus, 25'63 may be read 2563 hundredths, and the whole may be expressed in the form of a common fraction, thus,  $\frac{2563}{100}$ .

The denominations in *FEDERAL MONEY* are made to correspond to the decimal divisions of a unit now described, *dollars being units, or whole numbers, dimes tenths, cents hundredths, and mills thousandths of a dollar*; consequently, the

*expression of any sum in dollars, cents, and mills, is simply the expression of a mixed number in decimal fractions.*

Forty-six and seven tenths  $= 46\frac{7}{10} = 46.7$ .

Write the following numbers in the same manner :

Eighteen and thirty-four hundredths.

Fifty-two and six hundredths.

Nineteen and four hundred eighty-seven thousandths.

Twenty and forty-two thousandths.

One and five thousandths.

135 and 3784 ten-thousandths.

9000 and 342 ten-thousandths.

10000 and 15 ten-thousandths.

974 and 102 millionths.

320 and 3 tenths, 4 hundredths and 2 thousandths.

500 and 5 hundred-thousandths.

47 millionths.

Four hundred and twenty-three thousandths.

### ADDITION AND SUBTRACTION OF DECIMAL FRACTIONS.

T 67. As the value of the parts in decimal fractions increases in the same proportion as units, tens, hundreds, &c. and may be read *together*, in the same manner as whole numbers, so, it is evident that *all the operations on decimal fractions may be performed in the same manner as on whole numbers*. The only difficulty, if any, that can arise, must be in finding *where to place the decimal point* in the result. This, in addition and subtraction, is determined by the same rule; consequently, they may be exhibited together.

1. A man bought a barrel of flour for \$8, a firkin of butter for \$3.50, 7 pounds of sugar for 83½ cents, an ounce of pepper for 6 cents; what did he give for the whole?

*Note.* See the table of Federal Money, T 27. Let the pupil go back now and read carefully all that is said respecting federal money, in Reduction. From what is there stated it is plain, that we may readily reduce any sums in federal money to the same denominations, as to cents or mills,

lls, is simply  
tions.

anner :

ousandths.

andths.

s.

## DECIMAL

l fractions in-  
hundreds, &c.  
er as whole  
ns on decimal  
r as on whole  
n arise, must  
in the result.  
d by the same  
gether.

a firkin of but-  
an ounce of  
whole?

27. Let the  
said respect-  
is there stated  
ms in federal  
ts or mills,

and add or subtract them as simple numbers. Or, what is the same thing, we may set down the sums, taking care to write dollars under dollars, cents under cents, and mills under mills, in such order, that the separating points of the several numbers shall fall directly under each other, and add them as simple numbers, placing the separatrix in the amount directly under the other points.

### OPERATION.

$$\begin{array}{rcl} \$8 & = & 8000 \text{ mills, or } 1000\text{ths of a dollar} \\ 3\cdot50 & = & 3500 \text{ mills, or } 1000\text{ths.} \\ \cdot835 & = & 835 \text{ mills, or } 1000\text{ths.} \\ \cdot06 & = & 60 \text{ mills, or } 1000\text{ths.} \end{array}$$

$$\text{Ans. } \$12\cdot395 = 12395 \text{ mills, or } 1000\text{ths.}$$

As the denominations of federal money correspond with the parts of decimal fractions, so the rules for adding and subtracting decimals are exactly the same as for the same operations in federal money.

2. A man owing \$375, paid \$175\cdot75; how much did he then owe?

### OPERATION.

$$\begin{array}{rcl} \$375 & = & 37500 \text{ cents, or } 100\text{ths of a dollar.} \\ 175\cdot75 & = & 17575 \text{ cents, or } 100\text{ths of a dollar.} \\ \$199\cdot25 & = & 19925 \text{ cents, or } 100\text{ths.} \end{array}$$

Wherefore,—In the addition and subtraction of decimal fractions,—**RULE**: Write the numbers under each other, tenths under tenths, hundredths under hundredths, according to the value of their places, and point off in the results as many places for decimals as are equal to the greatest number of decimal places in any of the given numbers.

### EXAMPLES FOR PRACTICE.

3. Bought 1 barrel of flour for 6 dollars 75 cents, 10 lb. of coffee for 2 dollars 30 cents, 7 lb. of sugar for 92 cents, 1 lb. of raisins for 12½ cents, and 2 oranges, for 6 cents; what was the whole amount? *Ans. \$10\cdot155*

4. A man is indebted to A, \$237\cdot62; to B, \$350; to C, \$86\cdot12½ cents; to D, \$9\cdot62½, and to E, \$0\cdot834; what is the amount of his debts? *Ans. \$684\cdot204*

5. A man has three notes specifying the following sums, viz. three hundred dollars, fifty dollars sixty cents, and

nine dollars eight cents ; what is the amount of the three notes ?

*Ans.* 8359·68.

6. A man gave 4 dollars 75 cents for a pair of boots, and 2 dollars 12½ cents for a pair of shoes ; how much did the boots cost more than the shoes ?

OPERATION.

4750 mills.

or,

2125 mills.

OPERATION.

84·75

82·125

2625 mills = 82·625 *Ans.* 82·625 *Ans.*

7. A man bought a cow for eighteen dollars, and sold her again for twenty-one dollars thirty-seven and a half cents ; how much did he gain ?

*Ans.* 83·375.

8. A man bought a horse for 82 dollars, and sold him again for seventy-nine dollars seventy-five cents ; did he gain or lose ? and how much ?

9. A man sold wheat at several times as follows, viz. 13·25 bushels ; 8·4 bushels ; 23,051 bushels ; 6 bushels, and 75 of a bushel ; how much did he sell in the whole ?

*Ans.* 51·451 bushels.

10. What is the amount of 429, 21<sup>37</sup>/<sub>100</sub>, 355<sup>3</sup>/<sub>1000</sub>, 1<sup>7</sup>/<sub>100</sub> and 1<sup>7</sup>/<sub>10</sub> ?

*Ans.* 808<sup>143</sup>/<sub>1000</sub>, or 808·143.

11. What is the amount of 2 tenths, 80 hundredths, 89 thousandths, 6 thousandths, 9 tenths, and 5 thousandths ?

*Ans.* 2.

12. What is the amount of three hundred twenty-nine, and seven tenths ; thirty-seven and one hundred sixty-two thousandths, and sixteen hundredths ?

13. A man, owing \$4316, paid \$376·865 ; how much did he then owe ?

*Ans.* \$3939·135.

14. From thirty-five thousand take thirty-five thousandths.

*Ans.* 34999·965.

15. From 5·83 take 4·2793.

*Ans.* 1·5507.

16. From 480 take 245·0075.

*Ans.* 234·9925.

17. What is the difference between 1793·13 and 817·05693 ?

*Ans.* 976·07307.

18. From 4<sup>8</sup>/<sub>100</sub> take 2<sup>1</sup>/<sub>10</sub>-

*Remainder,* 1<sup>88</sup>/<sub>100</sub>, or 1·98.

19. What is the amount of 29<sup>3</sup>/<sub>10</sub>, 374<sup>8</sup>/<sub>1000000</sub>, 97<sup>353</sup>/<sub>1000</sub>, 315<sup>4</sup>/<sub>1000</sub>, 27, and 100<sup>4</sup>/<sub>10</sub> ?

*Ans.* 942·957009.



f the three  
8359·68.  
boots, and  
uch did the

ON.

5 Ans.  
and sold her  
half cents ;  
ns. 83·375.  
and sold him  
ts ; did he

ollows, viz.  
; 6 bushels,  
the whole ?  
451 bushels.

1000, 1000  
or 808·143.  
ndredths, 89  
ousandths ?

Ans. 2.  
twenty-nine,  
ed sixty-two

; how much  
\$3939·135.  
y-five thou-  
\$34999·965.

Ans. 1·5507.  
s. 234·9925.  
13 and 817  
e 976,07307.  
00, or 1·98.

00, 97 333  
942·957009.

# MULTIPLICATION OF DECIMAL FRACTIONS.

¶ 68. 1. How much hay is 7 loads, each containing 23·571 cwt?

OPERATION.

$$\begin{array}{r} 23\cdot571 \text{ cwt.} = 23571 \text{ 1000ths of a cwt.} \\ 7 \qquad \qquad \qquad 7 \end{array}$$

$$\text{Ans. } 164,997 \text{ cwt.} = 164997 \text{ 1000ths of a cwt.}$$

We may here (¶ 66) consider the multiplicand so many *thousandths* of a cwt., and then the product will evidently be *thousandths*, and will be reduced to a mixed or whole number by pointing off 3 figures, that is, the same number as are in the multiplicand ; and as either factor may be made the multiplier, so, if the decimals had been in the *multiplier*, the same number of places must have been pointed off for decimals. Hence it follows, *we must always point off in the product as many places for decimals as there are decimal places in both factors.*

2. Multiply '75 by '25.

OPERATION.

$$\begin{array}{r} \cdot75 \\ \cdot25 \\ \hline 375 \\ 150 \\ \hline \end{array}$$

1875 *Product.* If we consider the two factors as common or vulgar fractions. Thus, '75 is  $\frac{75}{100}$ , and '25 is  $\frac{25}{100}$  : now,  $\frac{75}{100} \times \frac{25}{100} = \frac{1875}{10000} = 1875$ , *Ans.* same as before.

3. Multiply '125 by '03.

OPERATION.

$$\begin{array}{r} \cdot125 \\ \cdot03 \\ \hline \end{array}$$

00375 *Product.*

Here, as the number of significant figures in the product is not equal to the number of decimals in both factors, the deficiency must be supplied by prefixing ciphers, that is, placing them at the left hand. The correctness of the rule may appear from the following process : '125 is  $\frac{125}{1000}$ , and '03 is  $\frac{3}{100}$  : now,  $\frac{125}{1000} \times \frac{3}{100} = \frac{375}{100000} = 00375$ , the same as before.

These examples will be sufficient to establish the following

## RULE.

In the multiplication of decimal fractions, multiply as in whole numbers, and from the product point off so many figures for decimals as there are decimal places in the multiplicand and multiplier counted together, and, if there are not so many figures in the product, supply the deficiency by prefixing ciphers.

As the denominations of federal money correspond with the parts of decimal fractions; the rules for the multiplication and division of both are the same.

## EXAMPLES FOR PRACTICE.

4. At \$5'47 per yard, what cost 8'3 yards of cloth?  
*Ans.* \$45'401.
5. At \$'07 per pound, what cost 26'5 pounds of rice?  
*Ans.* \$1'855.
6. If a barrel contain 1'75 cwt. of flour, what will be the weight of '63 of a barrel?  
*Ans.* 1'1025 cwt.
7. If a melon be worth \$'00, what is '7 of a melon worth?  
*Ans.* 6 $\frac{3}{10}$  cents.
8. Multiply five hundredths by seven thousandths.  
*Product,* '00035.
9. What is '3 of 116?  
*Ans.* 34'8.
10. What is '85 of 3872?  
*Ans.* 3121'2.
11. What is '37 of '0563?  
*Ans.* '020831.
12. Multiply 572 by '58.  
*Product,* 331'76.
13. Multiply eighty-six by four hundredths.  
*Product,* 3'44.
14. Multiply '2062 by '0008.
15. Multiply forty-seven tenths by one thousand eighty-six hundredths.
16. Multiply two hundredths by eleven thousandths.
17. What will be the cost of thirteen hundredths of a ton of hay, at \$11 a ton?
18. What will be the cost of three hundred seventy-five thousandths of a cord of wood at \$2 a cord?
19. If a man's wages be seventy-five hundredths of a dollar a day, how much will he earn in 4 weeks, Sundays excepted?
20. What will 250 bushels of rye come to, at \$0'88 $\frac{1}{2}$  per bushel?  
*Ans.* \$221'25.
21. What is the value of 87 barrels of flour, at \$6'37 $\frac{1}{2}$  a barrel?

multiply as in  
so many fig-  
the multipli-  
there are not  
efficiency by

spond with  
multiplica-

f cloth ?  
s. \$45.401.  
s of rice ?  
Ans. \$1.855.  
t will be the  
1.1025 cwt.  
melon worth?  
s. 6 <sup>10</sup>/<sub>100</sub> cents.  
ndths.

uct, .00035.  
Ans. 34.8.  
Ans. 3121.2.  
ns. .020831.  
uct, 331.76.

product, 3.44.

and eighty-

sandths.  
redths of a

seventy-five

redths of a  
ks, Sundays

\$0.88 <sup>1</sup>/<sub>2</sub> per  
s. \$221.25.  
t \$6.37 <sup>1</sup>/<sub>2</sub> a

22. What will be the cost of a hogshead of molasses, containing 63 gallons, at 28 <sup>1</sup>/<sub>2</sub> cents a gallon? Ans. \$17.955.

23. If a man spend 12 <sup>1</sup>/<sub>2</sub> cents a day, what will that amount to in a year of 365 days? what will it amount to in 5 years? Ans. \$228.12 <sup>1</sup>/<sub>2</sub> in 5 years.

## DIVISION OF DECIMAL FRACTIONS.

169. Multiplication is proved by division. We have seen, in multiplication, that the decimal places in the product must always be equal to the number of decimal places in the multiplicand and multiplier counted together. The multiplicand and multiplier, in proving multiplication, become the divisor and quotient in division. It follows of course, in division, that the number of decimal places in the divisor and quotient, counted together, must always be equal to the number of decimal places in the dividend. This will still further appear from the examples and illustrations which follow :

1. If 6 barrels of flour cost \$44.718, what is that a barrel?

By taking away the decimal point,  $44.718 = 44718$  mills, or 1000ths, which, divided by 6, the quotient is 7453 mills, = \$7.453, the answer.

Or, retaining the decimal point, divide as in whole numbers?

OPERATION.

6)44.718

Ans. 7.453

As the decimal places in the divisor and quotient, counted together, must be equal to the number of decimal places in the dividend, there being no decimals in the divisor,—therefore point off three figures for decimals in the quotient, equal to the number of decimals in the dividend, which brings us to the same result as before.

2. At \$4.75 a barrel for cider, how many barrels may be bought for \$31?

In this example, there are decimals in the divisor, and none in the dividend.  $4.75 = 475$  cents, and \$31, by annexing two ciphers, = 3100 cents; that is, reduce the dividend to parts of the same denomination as the divisor.

Then, it is plain, as many times as 475 cents are contained in 3100 cents, so many barrels may be bought.

475)3100(6 $\frac{4}{5}$  barrels the answer; that is, 6 barrels and  $\frac{4}{5}$  of another barrel.

2850

250

But the remainder, 250, instead of being expressed in the form of a common fraction, may be reduced to 10ths by annexing a cipher, which, in effect, is multiplying it by 10, and the division continued, placing the decimal point after the 6, or whole ones already obtained, to distinguish it from the decimals which are to follow. The points may be withdrawn or not from the divisor and dividend.

#### OPERATION.

475)31'00(6'526+ barrels, the answer, that is, 6 barrels and 526 thousandths of another barrel.

2850

2500

2375

1250

950

3000

2850

150

By annexing a cipher to the first remainder, thereby reducing it to 10ths, and continuing the division, we obtain from it '5, and a still further remainder of 125, which, by annexing another cipher, is reduced to 100ths, and so on.

The last remainder, 150, is  $\frac{1}{10}$  of a thousandth part of a barrel, which is of so trifling a value, as not to merit notice.

If now we count the decimals of the dividend, (for every cipher annexed to the remainder is evidently to be counted a decimal of the dividend,) we shall find them to be five, which corresponds with the number of decimal places in the divisor and quotient counted together.

3. Under T 68, ex. 3, it was required to multiply '125 by '03; the product was '00375. Taking this product for a dividend, let it be required to divide '00375 by '125. One operation will prove the other. Knowing that the number of decimal places in the quotient and divisor, counted together, will be equal to the decimal places in the dividend, we may divide as in whole numbers, being careful to retain the decimal points in their proper places. Thus,

OPERATION.

‘125)‘00375(‘03

375

—

000

The divisor, 125, in 375 goes 3 times and no remainder. We have only to place the decimal point in the quotient, and the work is done.

There are five decimal places in the dividend; consequently there must be five in the divisor and quotient counted together; and, as there are *three* in the divisor, there must be two in the quotient; and since we have but one figure in the quotient, the *deficiency* must be supplied by prefixing a cypher.

The operation by vulgar fractions will bring us to the same result. Thus, ‘125 is  $\frac{125}{1000}$ , and ‘00375 is  $\frac{375}{100000}$ ; now,  $\frac{375}{100000} \div \frac{125}{1000} = \frac{375000}{12500000} = \frac{3}{100} = .03$  the same as before.

† 70. The foregoing examples and remarks are sufficient to establish the following

RULE.

In the division of decimal fractions, divide as in whole numbers, and from the right hand of the quotient point off as many figures for decimals, as the decimal figures in the dividend exceed those in the divisor, and if there are not so many figures in the quotient, supply the deficiency by prefixing ciphers.

If at any time there is a remainder, or if the decimal figures in the divisor exceed those in the dividend, ciphers may be annexed to the dividend or the remainder, and the quotient carried to any necessary degree of exactness; but the ciphers annexed must be counted so many decimals of the dividend.

EXAMPLES FOR PRACTICE.

4. If \$472.875 be divided equally between 13 men, how much will each one receive? *Ans.* \$36.375.

5. At \$.75 per bushel, how many bushels of rye can be bought for \$141? *Ans.* 188 bushels.

6. At 6¢ cents apiece, how many oranges may be bought for \$8? *Ans.* 133 oranges.

7. If ‘6 of a barrel of flour cost \$5, what is that per barrel? *Ans.* \$8.333+

8. Divide 2 by 53.1. *Quot.* ‘037+

9. Divide '012 by '005.

Quot. 2'4.

11. Divide three thousandths by four hundredths.

Quot. '075.

11. How many times is '17 contained in 8?

12. If I pay \$468'75 for 750 pounds of wool, what is the value of 1 pound? Ans. \$0'625; or thus \$0'62½.

13. If a piece of cloth, measuring 125 yards, cost \$181'25 what is that a yard? Ans. \$1'45.

14. If 536 quintals of fish cost \$1913'52, how much is that a quintal? Ans. \$3'57.

15. Bought a farm, containing 84 acres, for \$3213; what did it cost me per acre? Ans. \$38'25.

16. At \$954 for 3816 yards of flannel, what is that per yard? Ans. 80'25.

## REDUCTION OF COMMON OR VULGAR FRACTIONS TO DECIMALS.

¶ 71. 1. A man has  $\frac{1}{5}$  of a barrel of flour; what is that expressed in decimal parts?

As many times as the denominator of a fraction is contained in the numerator, so many whole ones are contained in the fraction. We can obtain no whole ones in  $\frac{1}{5}$ , because the denominator is not contained in the numerator. We may, however, reduce the numerator to *tenths*, (¶ 69, ex. 2.) by annexing a cipher to it, (which, in effect, is multiplying it by 10,) making 40 tenths, or 4'0. Then, as many times as the denominator, 5, is contained in 40, so many *tenths* are contained in the fraction. 5 into 40 goes 8 times, and no remainder. Ans. '8 of a bushel.

2. Express  $\frac{3}{4}$  of a dollar in decimal parts.

The numerator, 3, reduced to tenths, is 30, 3'0, which, divided by the denominator, 4, the quotient is 7 tenths, and a remainder of 2. This remainder must now be reduced to *hundredths* by annexing another cipher, making 20 hundredths. Then, as many times as the denominator 4, is contained in 20, so many *hundredths* also may be obtained. 4 into 20 goes 5 times, and no remainder.  $\frac{3}{4}$  of a dollar, therefore, reduced to decimals, is 7 tenths and 5 hundredths, that is, '75 of a dollar.

Quot. 2'4.  
 edths.  
 Quot. '075.

ool, what is  
 thus 80'62½.  
 cost 8181'25  
 Ans. 81'45.  
 ow much is  
 Ans. 83'57.  
 3213; what  
 Ans. 838'25.  
 at is that per  
 Ans. 80'25.

# AR FRAC-

what is that

n is contain-  
 contained in  
 n ½, because  
 erator. We  
 169, ex. 2.)  
 multiplying  
 many times  
 many tenths  
 8 times, and  
 of a bushel.

3'0, which,  
 tenths, and  
 e reduced to  
 ing 20 hun-  
 or 4, is con-  
 obtained. 4  
 of a dollar,  
 hundredths,

The operation may be presented in form as follows :—

$$\begin{array}{r} \text{Num.} \\ \text{Denom. } 4)3'0('75 \text{ of a dollar, the answer.} \\ \underline{28} \end{array}$$

$$\begin{array}{r} 20 \\ \underline{20} \end{array}$$

3. Reduce  $\frac{4}{88}$  to a decimal fraction.

The numerator must be reduced to *hundredths* by annex-  
 ing two ciphers, before the division can begin.

66) 4'00( '0606+, the answer.

$$\begin{array}{r} 396 \\ \underline{\phantom{000}} \end{array}$$

$$\begin{array}{r} 400 \\ \underline{396} \end{array}$$

$$\begin{array}{r} 4 \end{array}$$

As there can be no *tenths*, a cipher must  
 be placed in the quotient, in tenth's place.

*Note.*  $\frac{4}{88}$  cannot be reduced *exactly*; for, however long  
 the division be continued, there will still be a remainder.\*  
 It is sufficiently exact for most purposes, if the decimal be  
 extended to three or four places.

\*Decimal figures which *continually repeat*, like '06, in this exam-  
 ple, are called *Repetends* or *Circulating Decimals*. If only *one figure*  
 repeats, as '3333 or '7777, &c. it is called a *single repetend*. If *two or*  
*more figures* circulate alternately, as '060606, '234234234, &c. it is  
 called a *compound repetend*. If other figures arise *before* those which  
 circulate, as '743333, '143010101, &c. the decimal is called a *mixed*  
*repetend*.

A *single repetend* is denoted by writing only the *circulating figure*  
 with a point over it: thus, '3, signifies that the 3 is to be continually  
 repeated, forming an *infinite* or *never ending series* of 3's.

A *compound repetend* is denoted by a point over the *first* and *last re-*  
*peating figure*: thus, '234 signifies that 234 is to be continually re-  
 peated.

It may not be amiss, here to show how the *value* of any *repetend*  
 may be found, or in other words, how it may be *reduced to its equiva-*  
*lent vulgar fraction*.

If we attempt to reduce  $\frac{1}{3}$  to a *decimal*, we obtain a con-  
 tinual repetition of the figure 1: thus, '11111, that is, the  
*repetend* '1. The value of the repetend '1, then is  $\frac{1}{3}$ ; the value  
 of '222, &c. the repetend '2 will be *twice* as much; that is,



From the foregoing examples we may deduce the following general RULE:—*To reduce a common to a decimal fraction:—Annex one or more ciphers, as may be necessary, to the numerator, and divide it by the denominator. If then there be a remainder, annex another cipher, and divide as before, and so continue to do so long as there shall continue*

2. In the same manner,  $'3 = \frac{3}{10}$ , and  $'4 = \frac{4}{10}$ , and  $'5 = \frac{5}{10}$ , and so on to 9, which  $= \frac{9}{10}$ .

1. What is the value of  $'8$ ?      *Ans.  $\frac{8}{10}$ .*

2. What is the value of  $'6$ ?      *Ans.  $\frac{6}{10}$ .* What is the value of  $'3$ ? — of  $'7$ ? — of  $'4$ ? — of  $'5$ ? — of  $'9$ ? — of  $'1$ ?

If  $\frac{1}{99}$  be reduced to a decimal, it produces  $'010101$ , or the repetend  $'01$ . The repetend  $'02$ , being 2 times as much, must be  $\frac{2}{99}$  and  $'03 = \frac{3}{99}$ , and  $'48$ , being 48 times as much, must be  $\frac{48}{99}$ , and  $'74 = \frac{74}{99}$ , &c.

If  $\frac{1}{999}$  be reduced to a decimal, it produces  $'001$ ; consequently,  $'002 = \frac{2}{999}$ , and  $'037 = \frac{37}{999}$ , and  $425 = \frac{425}{999}$ , &c. As this principle will apply to any number of places, we have this general RULE for reducing a circulating decimal to a vulgar fraction.—*Make the given repetend the numerator, and the denominator will be as many 9s as there are repeating figures.*

3. What is the vulgar fraction equivalent to  $'704$ ?

*Ans.  $\frac{704}{999}$ .*

4. What is the value of  $'003$ ? —  $'014$  —  $'324$  —  $'01021$ ? —  $'2463$ ? —  $'002103$ ?      *Ans. to the last,  $\frac{701}{999}$ .*

5. What is the value of  $'43$ ?

In this fraction, the repetend begins in the second place, or place of hundredths. The first figure, 4, is  $\frac{4}{10}$ , and the repetend, 3, is  $\frac{3}{100}$ , that is,  $\frac{3}{10}$ ; these two parts must be added together.  $\frac{4}{10} + \frac{3}{100} = \frac{43}{100} = \frac{43}{100}$ , *ans.* Hence, to find the value of a mixed repetend,—Find the value of the two parts separately, and add them together.

6. What is the value of  $'153$ ?  $\frac{15}{100} + \frac{3}{1000} = \frac{153}{1000} = \frac{153}{1000}$ , *ans.*

7. What is the value of  $'138$ ? —  $'16$ ? —  $'4123$ ?

It is plain, that circulating may be added, subtracted, multiplied, and divided, by first reducing them to their equivalent vulgar fractions.

duce the follow-  
a decimal frac-  
be necessary, to  
inator. If then  
r, and divide as  
e shall continue

, and  $5 = \frac{5}{1}$ , and

Ans.  $\frac{1}{10}$ .

What is the  
— '91 — '1?  
s '010101, or the  
ing 2 times as  
eing 48 times as

ces '001; conse-  
425 =  $\frac{425}{1000}$ , &c.  
r of places, we  
culating decimal  
extend the numer-  
9s as there are

t to '704?

Ans.  $\frac{704}{1000}$ .

— '324? —

the last,  $\frac{701}{1000}$ .

the second place,  
is  $\frac{4}{10}$ , and the  
two parts must be  
Hence, to find  
value of the two

=  $\frac{138}{1000} = \frac{23}{100}$ , ans.

6? — '4123?

subtracted, mul-  
to their equiva-

to be a remainder, or until the fraction shall be reduced to any necessary degree of exactness. The quotient will be the decimal required, which must consist of as many decimal places as there are ciphers annexed to the numerator; and if there are not so many figures in the quotient, the deficiency must be supplied by prefixing ciphers.

### EXAMPLES FOR PRACTICE.

4. Reduce  $\frac{1}{2}$ ,  $\frac{1}{4}$ ,  $\frac{1}{16}$ , and  $\frac{9}{1125}$  to decimals.  
Ans. '5; '25; '025; '00797+
5. Reduce  $\frac{37}{1000}$ ,  $\frac{1000}{1785}$ , and  $\frac{5000}{5000}$  to decimals.  
Ans. '002+; '003; '0028+; '000183+
6. Reduce  $\frac{475}{1000}$ ,  $\frac{10}{100}$ ,  $\frac{10}{1000}$  to decimals.
7. Reduce  $\frac{1}{10}$ ,  $\frac{5}{100}$ ,  $\frac{5}{1000}$ ,  $\frac{1}{10}$ ,  $\frac{3}{10}$ ,  $\frac{1}{10}$ ,  $\frac{1}{10}$ ,  $\frac{5}{100}$  to decimals.
8. Reduce  $\frac{1}{10}$ ,  $\frac{3}{10}$ ,  $\frac{5}{10}$ ,  $\frac{1}{10}$ ,  $\frac{2}{10}$ ,  $\frac{3}{10}$ ,  $\frac{4}{10}$ ,  $\frac{1}{20}$ ,  $\frac{1}{20}$ ,  $\frac{7}{10}$  to decimals.

### REDUCTION OF DECIMAL FRACTIONS.

¶ 72. Fractions, we have seen (¶ 60,) like *integers*, are reduced from low to higher denominations by *division*, and from high to lower denominations by *multiplication*.

To reduce a compound num-  
ber to a decimal of the highest  
denomination.

1. Reduce 7s. 6d. to the  
decimal of a pound.

6d. reduced to the decimal  
of a shilling, that is, divided  
by 12, is '5s., which annexed  
to the 7s. making 7'5s., and  
divided by 20, is '375£ the  
answer.

The process may be pre-  
sented in form of a *rule*, thus:—  
Divide the *lowest* denomina-  
tion given, annexing to it one  
or more ciphers, as may be  
necessary, by that number  
which it takes of the same to  
make one of the next higher  
denomination, and annex the

To reduce the decimal of a  
higher denomination to *integers*  
of lower denominations.

2. Reduce '375£ to inte-  
gers of lower denominations.

'375£ reduced to shillings,  
that is, multiplied by 20, is  
7'50s.; then the *fractional*  
part, '50s., reduced to pence,  
that is, multiplied by 12, is  
6d. Ans. 7s. 6d.

That is,—Multiply the given  
decimal by that number which  
it takes of the next *lower* de-  
nomination to make *one* of this  
higher, and from the right  
hand of the product point off  
as many figures for decimals  
as there are figures in the  
given decimal, and so contin-

quotient, as a decimal to that value to do through all the de- higher denomination; so con- nominations; the several num-  
 continue to do, until the whole bers at the left hand of the  
 shall be reduced to the deci- decimal points will be the  
 mal required. value of the fraction in the  
 proper denominations.

### EXAMPLES FOR PRACTICE

3. Reduce 1 oz. 10pwt. to the fraction of a pound.

#### OPERATION.

20)10'0 pwt.

12)1'5 oz.

125 lb. *Ans.*

5. Reduce 4cwt. 2 $\frac{3}{4}$ qrs to the decimal of a ton.

*Note.* 2 $\frac{3}{4}$  = 2'6.

7. Reduce 38gals. 3'52qts. of beer, to the decimal of a hhd.

9. Reduce 1qr. 2n. to the decimal of a yard.

11. Reduce 17h. 6m. 43s. to the decimal of a day.

13. Reduce 21s. 10 $\frac{1}{2}$ d. to the decimal of a guinea.

15. Reduce 3cwt. 0qr. 7lbs. 8oz. to the decimal of a ton.

4. Reduce 125lbs. Troy to integers of lower denominations.

#### OPERATION.

lb. 125

12

oz. 1'500

20

pwt 10'000. *Ans.* 1oz. 10pwt

6. What is the value of 2325 of ton?

8. What is the value of 72 of a hogshead of beer?

10. What is the value of 375 of a yard?

12. What is the value of 713 of a day?

14. What is the value of 78125 of a guinea?

16. What is the value of 15334821 of a ton?

Let the pupil be required to reverse and prove the following examples:

17. Reduce 4 rods to the decimal of an acre.

18. What is the value of 7 of a lb. of silver?

19. Reduce 18 hours, 15m. 50'4sec. to the decimal of a day.

20. What is the value of 67 of a league?

21. Reduce 10s. 9 $\frac{1}{2}$ d. to the fraction of a pound.

773. There is a method of reducing shillings, pence,

and farthings to the decimal of a pound, by *inspection*, more simple and concise than the foregoing. The reasoning in relation to it is as follows :

$\frac{1}{20}$  of 20s is 2s. ; therefore every 2s. is  $\frac{1}{10}$ , or '1£. Every shilling is  $\frac{1}{20} = \frac{1}{100}$ , or '05£. Pence are readily reduced to farthings. Every farthing is  $\frac{1}{400}$ £. Had it so happened, that 1000 farthings, instead of 960, had made a pound, then every farthing would have been  $\frac{1}{1000}$ , or '001£.  $960 + 40 = 1000$  ; that is,  $\frac{1}{4}$  of 960 added to 960 is 1000. Taking  $\frac{1}{4}$  of a number, and adding to that number, is the same as multiplying the number by unity and the fraction  $\frac{1}{4}$ ,  $1\frac{1}{4}$ . Suppose you have the fraction  $\frac{24}{960}$ . If you multiply both the numerator, and denominator by  $1\frac{1}{4}$ , you do not change the value of the fraction. Do this, and you obtain  $\frac{36}{1200}$ .  $\frac{36}{1200}$  then is equal to  $\frac{36}{1200}$ .  $\frac{36}{1200}$ £ is 24 farthings ; of course it follows that 24 farthings is equal to  $\frac{36}{1200}$ £. Wherefore, if the number of farthings, in the given pence and farthings, be more than 12,  $\frac{1}{4}$  part will be more than  $\frac{1}{4}$  ; therefore add 1 to them : if they be more than 36,  $\frac{1}{4}$  part will be more than  $1\frac{1}{4}$  ; therefore add 2 to them : then call them so many thousandths, and the result will be correct within less than  $\frac{1}{4}$  of  $\frac{1}{1000}$  of a pound. Thus, 17s. 5½d. is reduced to the decimal of a pound as follows : 16s. = '8£ and 1s. = '05£. Then, 5½d = 23 farthings, which, increased by 1, (the number being more than 12, but not exceeding 36,) is '024£, and the whole is '874£ the answer.

Wherefore, to reduce shillings, pence and farthings to the decimal of a pound by inspection,—Call every two shillings one tenth of a pound ; every odd shilling, five hundredths ; and the number of farthings, in the given pence and farthings, so many thousandths, adding one, if the number be more than twelve and not exceeding thirty-six, and two, if the number be more than thirty-six.

§ 74. Reasoning as above, the result, or the three first figures in any decimal of a pound, may readily be reduced back to shillings, pence and farthings, by *inspection*. Double the first figure, or tenths, for shillings, and, if the second figure, or hundredths, be five or more than five, reckon another shilling ; then, after the five is deducted, call the figures

in the second and third place so many farthings, abating one when they are above twelve, and two when above thirty-six, and the result will be the answer, sufficiently exact for all practical purposes. Thus, to find the value of '876 £ by inspection :—

'8 tenths of a pound - - - = 16 shillings.  
 '05 hundredths of a pound - - - = 1 shilling.  
 '026 thousandths, abating 1, = 25 farthings = 0 s. 6½ d.  
 '876 of a pound - - - = 17 s. 6½ d.  
*Ans.*

### EXAMPLES FOR PRACTICE.

1. Find, by inspection, the decimal expressions of 9s. 7d. and 12s. 0½d. *Ans.* '479 £, and '603 £.

2. Find, by inspection, the value of '523 £, and '694 £. *Ans.* 10s. 5½d., and 13s. 10½d.

3. Reduce to decimals, by inspection, the following sums, and find their amount, viz : 15s. 3d.; 8s. 11½d.; 10s. 6½d.; 1s. 8½d.; ½d. and 2½d. *Amount, £1'833.*

4. Find the value of '47 £.

*Notes.* When the decimal has but two figures, after taking out the shillings, the remainder, to be reduced to thousandths will require a cipher to be annexed to the right hand, or supposed to be so. *Ans.* 9s. 4½d.

5. Value the following decimals, by inspection, and find their amount, viz. '785 £.; '357 £.; '916 £.; '74 £.; '5 £.; '25 £.; '09 £.; and '008 £. *Ans.* 3 £. 12s. 11d.

## SUPPLEMENT TO DECIMAL FRACTIONS.

### QUESTIONS.

1. What are decimal fractions? 2. Whence is the term derived? 3. How do decimals differ from common fractions? 4. How are decimal fractions written? 5. How can the proper denominator to a decimal fraction be known, if it be not expressed? 6. How is the value of every figure determined? 7. What does the first figure on the right hand of the decimal point signify? — the second figure? — the third figure? — fourth figure? 8. How do ciphers, placed at the right hand of decimals affect their value? 9. Placed at the left hand how do they affect their value? 10. How are decimals read? 11. How are decimal fractions, having different denominators, reduced to a

ings, abating  
 en above thir-  
 iciently exact  
 value of '876 £

= 16 shillings.  
 = 1 shilling.  
 = 0 s.  $6\frac{1}{2}$  d.  
 = 17 s.  $6\frac{1}{2}$  d.  
*Ans.*

sions of 9s. 7d.  
 £, and '603 £.  
 £, and '694 £.  
 and 13s.  $10\frac{1}{2}$  d.  
 following sums,  
 $\frac{1}{2}$  d.; 10s.  $6\frac{1}{2}$  d.;  
 amount, £1'833.

es, after taking  
 d to thousandths  
 right hand, or  
*Ans.* 9s.  $4\frac{1}{2}$  d.  
 ection, and find  
 ; '74£.; '5£.;  
 11d.

## FRACTIONS.

the term derived?  
 4. How are deci-  
 ominator to a dec-  
 How is the value  
 figure on the right  
 figure? — the  
 ers, placed at the  
 ed at the left hand  
 is read? 11. How  
 ters, reduced to a

common denominator? 12. What is a mixed number? 13. How may any whole number be reduced to decimal parts? 14. How can any mixed number be read together, and the whole expressed in the form of a common fraction? 15. What is observed respecting the denominations in federal money? 16. What is the rule for addition and subtraction of decimals, particularly as respects placing the decimal point in the results? — multiplication? — division? 17. How is a common or vulgar fraction reduced to a decimal? 18. What is the rule for reducing a compound number to a decimal of the highest denomination contained in it? 19. What is the rule for finding the value of any given decimal of a higher denomination in terms of a lower? 20. What is the rule for reducing shillings, pence and farthings to the decimal of a pound, by inspection? 21. What is the reasoning in relation to this rule? 22. How may the three first figures of any decimal of a pound be reduced to shillings, pence and farthings, by inspection?

## EXERCISES.

1. A merchant had several remnants of cloth, measuring as follows :

7 $\frac{3}{4}$ yds. }	How many yards in the whole, and what
6 $\frac{3}{4}$ " }	would the whole come to at \$3'67 per yard?
1 $\frac{1}{2}$ " }	<i>Note.</i> Reduce the common fractions to deci-
9 $\frac{1}{2}$ " }	mals. Do the same wherever they occur in the
8 $\frac{1}{4}$ " }	examples which follow.
3 $\frac{1}{10}$ " }	<i>Ans.</i> 36'475 yards. \$133'863 X, cost.

2. From a piece of cloth, containing  $36\frac{3}{4}$  yds. a merchant sold, at one time,  $7\frac{3}{10}$  yds. and, at another time,  $12\frac{3}{4}$  yds.; how much of the cloth had he left? *Ans.* 16'7 yds.

3. A farmer bought 7 yards of broadcloth for  $8\frac{1}{2}$  £., a barrel of flour for  $2\frac{1}{5}$  £., a cask of lime for 1£., and 7lbs. of rice for  $\frac{1}{4}$  £.; he paid 1 ton of hay at  $3\frac{7}{10}$  £., 1 cow at 6£. and the balance in pork at  $\frac{1}{10}$  £. per lb.; how many were the pounds of pork?

*Note.* In reducing the common fractions in this example, it will be sufficiently exact if the decimal be extended to three places.

4. At  $12\frac{1}{2}$  cents per lb. what will  $37\frac{1}{2}$  lbs. of butter cost? *Ans.* 108¢ lb.

*Ans.* \$4'718¢.

5. At \$17'37 per ton for hay, what will  $11\frac{1}{2}$  tons cost? *Ans.* \$201'92¢.

6. The above example reversed. At \$201'92¢ for  $11\frac{1}{2}$  tons of hay, what is that per ton? *Ans.* \$17'37.

7. If '45 of a ton of hay cost \$9, what is that per ton? *Ans.* \$20.  
*Consult T 62.*



8. At  $\frac{1}{4}$  of a dollar a gallon, what will  $\frac{1}{25}$  of a gallon of molasses cost ? Ans.  $\$1$ .

9. At  $\$9$  per cwt., what will 7 cwt. 3 qrs. 16 lbs. of sugar cost ?

*Note.* Reduce the 3 qrs. 16 lbs. to the decimal of a cwt. extending the decimal in this, and the examples which follow to four places. Ans.  $71\cdot935+$ .

10. At  $\$69\cdot875$  for 5 cwt. 1qr. 14lbs. of raisins, what is that per cwt. Ans.  $\$13$ .

11. What will 2300lbs of hay come to at 7 mills per lb. Ans.  $\$16\cdot10$

12. What will 765 $\frac{1}{2}$ lbs. of coffee come to at 18 cents per lb. Ans.  $\$137\cdot79$ .

13. What will 12 gallons, 3 qts. 1 pt. of gin cost, at 28 cents a quart ?

*Note.* Reduce the whole quantity to quarts and the decimal of a quart. Ans.  $\$14\cdot42$ .

14. Bought 16 yds. 2qrs. 3na. of broadcloth for  $\$100\cdot125$ ; what was that per yard ? Ans.  $\$6$ .

15. At  $\$1\cdot92$  per bushel, how much wheat may be purchased for  $\$72$  ? Ans. 1 peck 4 qts.

16. At  $\$92\cdot72$  per ton, how much iron may be purchased for  $\$60\cdot268$  ? Ans. 13cwt.

17. Bought a load of hay for  $\$9\cdot17$ , paying at the rate of  $\$16$  per ton ; what was the weight of the hay ?

Ans. 11cwt. 1qr. 23lbs.

18. At  $\$302\cdot4$  per tun, what will 1 hhd. 15 gals. 3 qts. of wine cost ? Ans.  $\$94\cdot50$ .

19. *The above reversed.* At  $\$94\cdot50$  for 1 hhd. 15 gals. 3qts. of wine, what is that per tun ? Ans.  $\$302\cdot4$ .

*Note.* The following examples reciprocally prove each other, excepting when there are some fractional losses, as explained above, and even then the results will be sufficiently exact for all practical purposes. If, however greater exactness be required, the decimals must be extended to a greater number of places.

20. At  $\$1\cdot80$  for  $3\frac{1}{2}$  qts. of wine, what is that per gal. ? 21. At  $\$2\cdot215$  per gallon, what cost  $3\frac{1}{2}$  qts. ?

22. If  $\frac{1}{4}$  of a ton of potash cost  $\$60\cdot45$ , what is that pot-ashes, what will  $\frac{1}{4}$  of a ton per ton ? 23. At  $\$96\cdot72$  per ton for pot-ashes, what will  $\frac{1}{4}$  of a ton cost ?



# REDUCTION OF CURRENCIES.

In the United States, since the act of Congress in 1786, establishing Federal Money, calculations in money, have generally been made in dollars, cents and mills. In England the denominations, though the same in name as the currency of this Province, are different in value. In the United States, previously to the act of Congress, it was the custom to reckon in pounds, shillings, &c. and now, though all accounts are kept in federal money, small sums are mentioned frequently in these denominations. There are different currencies of the same name in different parts of the United States. It may be necessary often in commercial dealings, and in the course of ordinary business, to change values in foreign currencies into the currency of the Provinces.

Supposing there is a sum in federal money—\$24'604. We find by the table of coins, ¶ 27, that 1 dollar is equal to 5 shillings, and of course 4 dollars equal 1 pound, there being 4 times 5 shillings in 20 shillings. The value of pounds, then it is clear, is 4 times that of dollars, and of course *dollars are reduced to pounds by dividing the given sum by 4.*

4) 24'604. dollars.

6 pounds.

There remain, however, '604, 60 cents and 4 mills, to be changed to Halifax currency. By referring to Decimal Fractions, ¶ 66, we see that dollars are the units in federal money, and cents and mills decimal parts; cents hundredths, and mills thousandths. We have then simply to divide these decimals of a dollar by 4, and the quotient will be in decimal parts of a pound, thus—

4) '604 of a dollar.

'151 of a pound.

This can be reduced to shillings, pence and farthings by inspection, (See ¶ 73,) as follows: '151£ equal 3s. 0d. 1qr. We find that \$24'604 is equal to 6£ 3s. 0d. 1qr. *Ans.*

The following then, is the general rule, to reduce federal money to Halifax currency,—*Divide the given sum by 4, and the quotient will be in pounds and decimal parts of a pound, which can be reduced to shillings, pence and farthings by inspection.*

## EXAMPLES FOR PRACTICE.

Reduce \$500 to Halifax currency,

*Ans.* 125£.

" 27'304 "

6£ 11s. 6d. 1qr.

" 118'25 "

29£ 11s. 3d.

" 236'50 "

59£ 2s. 6d.

Reduce \$490 to Halifax currency. — \$56'03, — \$93'814, — \$85'63, — \$1977'642 respectively to Halifax currency.

To reduce Halifax currency to federal money, we must reverse the process in the above examples. The rule is as follows: Reduce the shillings, pence, &c. if any, to the decimal of a pound, by inspection,

and multiply the given sum by 4, the product will be in the denominations of federal money.

## EXAMPLES FOR PRACTICE.

Red. 125£ Halifax currency to federal money.  
 " 59£ 2s. 6d. "

Ans. \$500.  
 \$239.50.

In order to change English sterling money, and the currencies in some degree in use in the different parts of the United States, into Halifax currency; since the denominations are the same in name, it will be necessary to take some other currency, the denominations of which are different, as a *common object of comparison* for these currencies, and for Halifax currency. By this means we shall be able to ascertain the values of the former relatively to those of the latter. We will take federal money as this common object of comparison; and will compare with a unit of one of its denominations, *the dollar*, one or more units of a denomination of Halifax currency, and the before mentioned currencies, *the shilling*.

In Halifax currency, - - - - - 5s. = \$1.

In English, or sterling money, (4s. 6d.)  $4\frac{1}{2}$ s. = \$1.

In New-England currency, - - - - - 6s. = \$1.

In New-York currency, - - - - - 8s. = \$1.

In Pennsylvania currency (7s. 6d.)  $7\frac{1}{2}$ s. = \$1.

In Halifax currency 5s. are equal to \$1, and in English sterling money, 4 shillings and a half are equal to \$1. Sterling money, then, is to Halifax currency as 5 to  $4\frac{1}{2}$ , or to avoid the fraction, as 10 to 9, since  $2 \times 5 = 10$ , and  $2 \times 4\frac{1}{2} = 9$ . Therefore, to change sterling money into Halifax currency,—Multiply by  $\frac{9}{10}$ , or take once the given sum, and  $\frac{1}{10}$ , thus,—

9)48£ 12s. 9d. sterling money.

5 8 1

54 0 10 Halifax currency.

Reduce 56£ 17s. 6d. sterling to Halifax currency. Ans. 63£ 3s. 10d.

Reduce 92 4 6 " " "

In New-England currency 6s. are reckoned to the dollar. New-England currency, then, is to Halifax currency as 5 to 6. Therefore, to reduce New-England currency to Halifax currency,—

Take  $\frac{5}{6}$  of the given sum, thus,—

6) 14£ 5s. 4d. New-England currency.

2 7  $5\frac{2}{3}$   
 5

11 17  $4\frac{1}{3}$  Halifax currency.

Reduce 60£ 4s. 10d. N. E. currency to Halifax currency.  
 Ans. 50£ 4s. 0 $\frac{2}{3}$ d.

n the denomin-

Ans. \$500.  
\$236 50.

a currencies in  
States, into Hal-  
in same, it will  
ations of which  
ese currencies,  
able to ascer-  
atter. We will  
rison; and will  
a dollar, one or  
the before men-

. = \$1.  
s. = \$1.  
s. = \$1.  
s. = \$1.  
s. = \$1.

English sterling  
g money, then,  
the fraction,  
Therefore, to  
y,—Multiply  
s,—

s. 63£ 3s. 10d.

dollar. New  
G. Therefore,

fax currency.  
50£ 4s. 0½d.

To reduce Halifax currency to sterling money, or to re-  
duce Halifax to New-England, it is only necessary to re-  
verse the process in the foregoing operations.

To reduce sterling to Halifax we multiply by  $\frac{10}{9}$ , there-  
fore, to reduce Halifax currency to sterling money,—Divide  
the given sum by  $\frac{10}{9}$ , or, what is the same, multiply by  $\frac{9}{10}$ ,  
that is, take  $\frac{9}{10}$  of the given sum; e. g.

10) 54£ 0s. 10d. Halifax currency.

$$\begin{array}{r} 5 \quad 8 \quad 1 \\ \hline 48 \quad 12 \quad 9 \text{ sterling money.} \end{array}$$

In the same manner, to reduce Halifax currency to New-  
England,—Take  $\frac{2}{3}$  of the given sum, or, add  $\frac{1}{3}$  to the giv-  
en sum.

From the foregoing rules and illustrations, the pupil him-  
self will be able, by pursuing a similar course, to reduce,  
with facility, any currency, the denominations of which are  
pounds, shillings, &c. to any other, in which the denomina-  
tions are the same.

The following is the *general rule* for finding a *multiplier*  
to reduce any currency to the par of another: Make the  
number of shillings that are equal to a dollar in the curren-  
cy to be reduced, the denominator of a fraction; and over this,  
for a numerator, write the number of shillings that are e-  
qual to a dollar in the currency to which the given sum is to  
be reduced.

Let the pupil find multipliers to reduce New-York and  
Pennsylvania, currencies to Halifax, and then Halifax cur-  
rency to those.

## INTEREST.

¶ 75. Interest is an allowance made by a debtor to a  
creditor for the use of money. It is computed at a certain  
number of pounds for the use of each hundred pounds, or  
so many dollars for each hundred dollars, &c. one year, and  
in the same proportion for a greater or less sum, or for a  
longer or shorter time.

The number of pounds so paid for the use of a hundred

pounds, one year, is called the *rate per cent* or *per centum*; the words *per cent*, or *per centum*, signifying *by the hundred*.

The highest rate allowed by law in the Canadas is 6 per cent\* that is, 6 pounds for 100 pounds, 6 shillings for 100 shillings; in other words  $\frac{6}{100}$  of the sum lent or due is paid for the use of it one year. This is called *legal interest*, and will be understood when no other rate is mentioned.

Let us suppose the sum lent, or due, to be 1£. The 100th part of 1£ or  $\frac{1}{100}$  of a pound is, decimally expressed, thus, '01, and  $\frac{6}{100}$  of a pound, the legal interest, written as a decimal fraction, is - - - - - '06.

So of any rate per cent.

1 per cent, expressed as a common fraction, is

$\frac{1}{100}$ ; decimally, - - - - - '01.

$\frac{1}{2}$  per cent, is a half of 1 per cent that is, - - - '005.

$\frac{1}{4}$  per cent is a fourth of 1 per cent that is, - - - '0025.

$\frac{3}{4}$  per cent is 3 times  $\frac{1}{4}$  per cent, that is, - - - '0075.

*Note.* The rate per cent is a decimal carried to two places, that is, to *hundredths*; all decimal expressions lower than hundredths are parts of 1 per cent.  $\frac{3}{8}$  per cent for instance, is '625 of 1 per cent that is, '00625.

Write  $2\frac{1}{2}$  per cent as a decimal fraction.

2 per cent is '02, and  $\frac{1}{2}$  per cent is '005. Ans. '025.

Write 4 per cent as a decimal fraction. —  $4\frac{1}{2}$  per cent.

—  $4\frac{1}{4}$  per cent. — 5 per cent. —  $7\frac{1}{4}$  per cent. —

8 per cent. —  $8\frac{1}{4}$  per cent. — 9 per cent. —  $9\frac{1}{4}$  per

cent. 10 per cent. — (10 per cent is  $\frac{10}{100}$ ; decimally, '10,)

—  $10\frac{1}{2}$  per cent. — 11 per cent. —  $12\frac{1}{4}$  per cent —

15 per cent.

1. If the interest on 1£ for 1 year, be '06 of a pound, what will be the interest on 25£ for the same time?

It will be 25 times 6, or 6 times 25, which is the same thing:—

25£.

'06

1'50 answer; that is, 1£ and 5 tenths. The 5 tenths must be reduced to shillings, pence and farthings by the rule

\*In the New-England States the legal rate is the same as in the Canadas. In England it is 5 per cent.

or per centum;  
by the hundred.  
anadas is 6 per  
hillings for 100  
or due is paid  
al interest, and  
is mentioned.  
he 1£. The  
ally expressed,  
rest, written as  
- - - '06.

on, is  
- - - '01.  
- - - '005.  
is, - - '0025.  
- - - '0075.  
carried to two pla-  
expressions lower  
½ per cent for  
25.

Ans. '025.  
— 4½ per cent.  
per cent. —  
nt. — 9½ per  
decimally, '10.)  
2½ per cent —

'06 of a pound,  
ame time 1  
which is the same

p. The 5 tenths  
things by the rule

the same as in the

for the reduction of decimals ; or with sufficient exact-  
ness by inspection. See ¶ 73. '50, or '5 of a pound equal 10  
shillings. The interest of 25£. for a year is then 1£. 10s.

To find the interest on any sum for 1 year, it is evident  
we need only multiply it by the rate per cent written as a  
decimal fraction. The product, observing to place the point  
as directed in multiplication of decimal fractions, will be  
the interest required.

*Note.* PRINCIPAL is the money due, for which interest is  
paid. AMOUNT is the principal and interest added together.

2. What will be the interest of 32£. 3s. 1 year, at 4½  
per cent?

We are to multiply the principal by the rate per cent, 4½,  
expressed in the form of a decimal, '045 ; we must therefore  
reduce the 3s. in the principal, to decimals by inspection.

We find 3s. equal to '15. There being five decimal places  
in the multiplicand and mul-  
tiplic 5 figures must be point-  
ed out for decimals from the  
product, which gives the an-  
swer 1 pound and 44675 hun-  
dred thousandths. Any thing  
less than thousandths need not

32'15£. principal.  
'045 rate per cent.

16075  
12860

1'44675£.

be regarded; hence, 1'446£ is sufficiently exact for the an-  
swer. The '446 must be reduced to shillings, pence and  
farthings by inspection. Double the '4 for shillings, equals  
8s.; call the '046 so many farthings, deducting two because  
one 36 equals 44 farthings. In 44 qrs. there are 11d.  
'446£=8s. 11d. The interest, then, of 32£. 3s. for 1  
year, at 4½ per cent, is 1£. 8s. 11d. answer.

Always, then, if there are shillings, pence and farthings,  
or either denomination, in the given principal, reduce them  
to the decimal of a pound by inspection, before multiplying by  
the rule. After obtaining the answer in decimals, reduce the  
tenths, hundredths and thousandths, to shillings, pence and far-  
things, by inspection. The method of effecting each reduc-  
tion, is exhibited in ¶ 73, and 74, and must be made per-  
fectly familiar to the pupil's mind.

3. What will be the interest of 11£. 3s. 4d. for 1 year,  
at 3 per cent ? — at 5½ per cent ? — at 6 per cent ? —  
at 7½ per cent ? — at 8½ per cent ? — at 9½ per cent ?  
— at 10 per cent ? — at 10½ per cent ? — at 11 per

cent? — at  $11\frac{1}{2}$  per cent? — at 12 per cent? — at  $12\frac{1}{2}$  per cent?

4. A tax on a certain town is 406£. 15s. 10 $\frac{1}{2}$ d., on which the collector is to receive  $2\frac{1}{2}$  per cent for collecting; what will he receive for collecting the whole tax at that rate?

In this example, the shillings, &c. reduced to the decimal of a pound equal .795. Multiply therefore, 406.795£. by the rate  $2\frac{1}{2}$ , that is, .025. The answer, in decimals, is 10.169£.; the tenths, &c. reduced to shillings, &c. equal 3s. 4 $\frac{1}{2}$ d. The answer, then, is, 10£. 3s. 4d.

*Note.* In the same way are calculated commission, insurance, buying and selling stocks, loss and gain, or any thing else rated at so much per cent *without respect to time.*

5. What must a man, paying  $37\frac{1}{2}$  per cent on his debts, pay on a debt of 132£. 5s. ? *Ans.* 49£. 11s. 10 $\frac{1}{2}$ d.

6. A merchant, having purchased goods to the amount of 580£. sold them so as to gain  $12\frac{1}{2}$  per cent, and in the same proportion for a greater or less sum; what was his whole gain, and what was the whole amount for which he sold the goods?

*Ans.* His whole gain was 72£. 10s. whole amount, 652£. 10s.

7. A merchant bought a quantity of goods for 173£. 15s. how much must he sell them for to gain 15 per cent?

*Ans.* 199£. 16s. 3d.

176. **COMMISSION** is an allowance of so much per cent to a person called a *correspondent*, *factor*, or *broker*, for assisting merchants and others in purchasing and selling goods.

8. My correspondent sends me word that he has purchased goods to the amount of 1286£, on my account; what will his commission come to at  $2\frac{1}{2}$  per cent? *Ans.* 32£. 3s.

9. What must I allow my correspondent for selling goods to the amount of 2317£ 9s. 2 $\frac{1}{2}$ d. at a commission of  $3\frac{1}{2}$  per cent? *Ans.* 75£ 6s. 4d.

**INSURANCE** is an exemption from hazard, obtained by the payment of a certain sum, which is generally so much per cent. on the estimated value of the property insured.

**PREMIUM** is the sum paid by the insured for the insurance.



ent! — at

d., on which  
ecting; what  
that rate?  
to the decimal  
400.795£. by  
decimals, is  
gs, &c. equal

mission, in-  
gain, or any  
spect to time.  
on his debts,  
0£. 11s. 10½d.  
to the amount  
nt, and in the  
what was his  
for which he

amount, 652£.

for 173£. 15s.  
per cent?  
199£. 16s. 3d.

of so much per  
r, or broker, for  
ng and selling

at he has pur-  
account; what  
Ans. 32£. 3s.  
for selling goods  
ision of 3¼ per  
ts. 75£ 6s. 4d.

rd, obtained by  
herally so much  
erty insured.  
or the insur-

**POLICY** is the name given to the instrument or writing, by which the contract of indemnity is effected between the insurer and insured.

10. What will be the premium for insuring a ship from Montreal to Liverpool, valued at 9450£, at  $4\frac{1}{2}$  per cent?

Ans. 425£ 5s.

11. What will be the annual premium for insurance on a house against loss by fire, valued at 875£ at  $\frac{3}{4}$  per cent?

By removing the separatrix 2 figures towards the left, it is evident, the sum itself may be made to express the premium at 1 per cent, of which the given rate parts may be taken; thus, 1 per cent on 875£ is 8.75£, and  $\frac{3}{4}$  of 875£ is 6.562£.

Ans. 6£ 11s. 3d.

12. What will be the premium for insurance on a ship and cargo valued at 6310£, at  $\frac{1}{2}$  per cent? — at  $\frac{3}{4}$  per cent? — at  $\frac{1}{2}$  per cent? — at  $\frac{3}{4}$  per cent? — at  $\frac{1}{2}$  per cent? Ans. At  $\frac{1}{2}$  per cent the premium is 39£ 7s. 8½d.

**Stock** is a general name, for the capital of any trading company or corporation, or of a fund established by government.

The value of stock is variable. When 100 pounds of stock sells for 100 pounds in *money*, the stock is said to be *at par*, which is a Latin word signifying *equal*; when for *more*, it is said to be *above par*; when for *less*, it is said to be *below par*.

13. What is the value of 756£ of stock, at  $112\frac{1}{2}$  per cent? that is, when 1 pound of stock sells for 1 pound  $12\frac{1}{2}$  hundredths in *money*, which is  $12\frac{1}{2}$  per cent above par, or  $2\frac{1}{2}$  per cent *advance*, as it is sometimes called.

Ans. 850£ 11s.

14. What is the value of 3700£ of bank stock, at  $95\frac{1}{2}$  per cent, that is,  $4\frac{1}{2}$  per cent *below par*? Ans. 3533£ 10s.

15. What is the value of 120£ of stock, at  $92\frac{1}{2}$  per cent? — at  $86\frac{1}{2}$  per cent? — at  $67\frac{1}{2}$  per cent? — at  $104\frac{1}{2}$  per cent? — at  $108\frac{1}{2}$  per cent? — at 115 per cent? — at  $37\frac{1}{2}$  per cent *advance*?

**LOSS AND GAIN.** 16. Bought a hogshead of molasses for 15£; for how much must I sell to gain 20 per cent?

Ans. 18£.

17. Bought broadcloth at 12s. 6d. per yard; but, it be-



ing damaged, I am willing to sell it so as to lose 12 per cent. ; how much will it be per yard ? *Ans.* 11s.

18. Bought calico at 1s. per yard ; how must I sell it to gain 5 per cent. ? — 10 per cent. ? — 15 per cent. ? — to lose 20 per cent. ? *Ans. to the last, 9 $\frac{3}{4}$ d. per yard.*

177. We have seen how interest is cast on any sum of money when the time is *one year* ; but it is frequently necessary to cast interest for months and days.

Now, the interest on 1£ for 1 year, at 6 per cent, being '06, is

'01, one hundredth for 2 months,

'005 five thousandths (or  $\frac{1}{2}$  a hundredth) for 1 month of 30 days, (for so we reckon a month in casting interest,) and

'001 one thousandth for every 6 days ; 6 being contained 5 times in 30.

Hence, it is very easy to cast in the mind, the interest on 1£, at 6 per cent for *any given time*. The *hundredths*, it is evident, will be equal to *half* the greatest even number of months ; the *thousandths* will be 5 for the odd month, if there be one, and 1 for every time 6 is contained in the given number of the days.

Suppose the interest of 1£, at 6 per cent, be required for 9 months, and 18 days. The greatest even number of the months is 8, half of which will be the hundredths, '04 ; the thousandths, reckoning 5 for the odd month, and 3 for the 18 (3 times 6 = 18) days, will be '008, which, united with the hundredths ('048) give 4 hundredths and 8 thousandths, 4 hundredths, and 8 thousandths, or, '048£ reduced = 11d.

*Ans.* 11d

1. What will be the interest on 1£ for 5 months 6 days ? — 6 months 12 days ? — 7 months ? — 8 months 24 days ? — 9 months 12 days ? — 10 months ? — 11 months 6 days ? — 12 months 18 days ? — 13 months 6 days ? — 16 months ?

ODD DAYS. 2. What is the interest of 1£ for 13 months 16 days ?

The hundredths will be 6, and the thousandths 5, for the odd month, and 2 for 2 times 6 = 12 days, and there is remainder of 4 days, the interest for which will be suc

76, 77.  
lose 12 per  
Ans. 11s.  
must I sell it to  
15 per cent. 1  
9d. per yard.  
on any sum  
is frequently  
er cent, being  
1 month of 30  
casting inter-  
being contained  
nd, the interest  
The hundredths,  
atest even num-  
the odd month,  
contained in the  
be required for  
n number of the  
redths, '04; the  
h, and 3 for the  
ich, united with  
d 8 thousandths,  
reduced=11d.  
Ans. 11d.  
6 months 6 days  
— 8 months  
0 months 1  
days? — 15  
1£ for 13 month  
sandths 5, for th  
s, and there is  
ich will be suc

part of 1 thousandth as 4 days is part of 6 days, that is,  $\frac{4}{6}$  of a thousandth. Ans. '067 $\frac{1}{3}$ .

3. What will be the interest of 1£ for 1 month 8 days 1  
— 2 months 7 days? — 3 months 15 days? — 4  
months 22 days? — 5 months 11 days? — 6 months  
17 days? — 7 months 3 days? — 8 months 11 days?  
— 9 months 2 days? — 10 months 15 days? — 11  
months 4 days? — 12 months 3 days?

Note. If there is no odd month, and the number of days be less than 6, so that there are no thousandths, it is evident, a cipher must be put in the place of thousandths; thus, in the last example,—12 months 3 days,—the hundredths will be '00, the thousandths 0, the 3 days  $\frac{1}{2}$  a thousandth.

Ans. 1s. 2 $\frac{1}{2}$ d.

4. What will be the interest of 1£ for 2 months 1 day?  
— 4 months 2 days? — 6 months 3 days? — 8 months  
4 days? — 10 months 5 days? — for 3 days? —  
for 1 day? — for 2 days? — for 4 days? — for 5  
days?

5. What is the interest of 56£ 2s. 7 $\frac{1}{2}$ d. for 8 months 5  
days? The interest of 1£, for the given time, is '040 $\frac{1}{2}$ ;  
therefore,

$\frac{1}{2}$ ) and  $\frac{1}{2}$ ) 56.13£. principal.

'040 $\frac{1}{2}$  interest of 1£ for the given time.

224520 interest for 8 months.

2806 interest for 3 days.

1871 interest for 2 days.

2.29197£ = 2£ 5s. 9 $\frac{1}{2}$ d.

5 days = 3 days + 2 days. As the multiplicand is taken once for every 6 days, for 3 days take  $\frac{1}{2}$ , for 2 days take  $\frac{1}{3}$ , of the multiplicand.  $\frac{1}{2} + \frac{1}{3} = \frac{5}{6}$ . So also, if the odd days be 4 = 2 days + 2 days, take  $\frac{1}{3}$  of the multiplicand twice; for 1 day, take  $\frac{1}{6}$ .

From the illustrations now given, it is evident,—To find the interest of any sum in Halifax currency, or any other currency of which the denominations are pounds, shillings, &c. at 6 per cent, it is only necessary to multiply the given principal, after having reduced the shillings and pence in it to the decimal of a pound by inspection, by the interest of 1£ for the given time, found as above directed and written as a deci-

mal fraction ; after pointing off as many places for decimals in the product as there are decimal places in both the factors counted together, these can be reduced back again to shillings and pence by inspection.

### EXAMPLES FOR PRACTICE.

6. What is the interest of 87£ 3s. 9½d. for 1 year 3 months ? *Ans.* 6£ 10s. 9½d.

7. Interest of 116£ 1s. 7½d. for 11 mo. 19 days ? *Ans.* 6£ 15s. 0½d.

8. Interest of 200£ for 8 mo. 4 days ? *Ans.* 8£ 2s. 7½d.

9. " of 17s. for 19 mo. 1 " *Ans.* 1s. 7½d.

10. " of 8£ 10s. for 1 year 9 mo. 12 days ? *Ans.* 18s. 2½d.

11. " of 675£ for 1 mo. 21 days ? *Ans.* 5£ 14s. 8½d.

12. " of 8673£ for 10 days ? *Ans.* 14£ 9s. 1½d.

13. " of 14s. 7½d. for 10 mo. 1 " *Ans.* 8½d.

14. " of 96£ for 3 days ? *Note.* The

15. " of 73£ 10s. for 2 days ? interest of 1£

16. " of 180£ 15s. for 5 days ? for 6 days be-

17. " of 15000£ for 1 day ? ing 1 thou-

sandth, the pounds *themselves* express the interest in *thousandths* for six days, of which we may take parts.

Thus, 6)15000 thousandths,

2500, that is, 2£ 10s *Ans.* to the last.

When the interest is required for a large number of years, it will be more convenient to find the interest for one year, and multiply it by the number of years ; after which find the interest for the months and days, if any, as usual.

18. What is the interest of 1000£ for 120 years ? *Ans.* 7200£.

19. What is the interest of 520£ 0s. 9½d. for 30 years and 6 months ? *Ans.* 951£ 13s. 5½d.

20. What is the interest on 400£ for 10 years 3 months and 6 days ? *Ans.* 246£ 8s.

21. What is the interest of 220£ for 5 years ? — for 12 years ? — 50 years ? *Ans.* to the last 660£.

22. What is the amount of 86£, at interest 7 years ? *Ans.* 122£ 2s. 4½d.

23. What is the interest of 943.30 for 1 year ?

It must be clear to the pupil's mind, that to obtain the

for decimals  
both the fac-  
back again to

r 1 year 3  
5£ 10s. 9½d.  
ays?

3£ 15s. 0½d.  
8£ 2s. 7½d.  
1s. 7½d.

ays?  
18s. 2½d.  
5£ 14s. 8½d.  
14£ 9s. 1½d.  
8½d.

Note. The  
interest of 1£  
or 6 days be-  
ing 1 thou-  
erest in thou-  
arts.

s. to the last.

number of years,  
for one year,  
er which find  
s usual.

years 1

Ans. 7200£.

for 30 years

1£ 13s. 5½d.

ars 3 months

ns. 246£ 8s.

ars 1 — for

he last 600£.

7 years 1

22£ 2s. 4½d.

ar?

to obtain the

the interest upon any sum in federal money, for any time, we proceed just as we do in Halifax currency; only we are not compelled to reduce any part of the given sum to decimals, since all the denominations of federal money are in a decimal ratio. The answer to the last example is \$2890.

What is the interest of \$64 for 2 years? *Ans.* \$7.08.

What is the interest of \$98.50 for 7 years, 6 months and 10 days? *Ans.* \$44.489.

78. 1. What is the interest of 36 pounds for 8 months, at 4½ per cent?

Note. When the rate is any other than six per cent, first find the interest at 6 per cent, then divide the interest so found by such part as the interest, at the rate required, exceeds or falls short of the interest, at 6 per cent, and the quotient added to or subtracted from the interest at 6 per cent, as the case may be, will give the interest required.

36£

04

144

36

4½ per cent is ¾ of 6 per cent; therefore, from the interest at 6 per cent subtract ¼; the remainder will be the interest at 4½ per cent.

1.08£. 1£ 1s. 7½d. answer.

2. Interest of 54£ 16s. 2½d. for 18 months, at 5 per cent 1

*Ans.* 4£ 2s. 2½d.

3. Interest of 500£ for 9 months 9 days, at 8 per cent?

*Ans.* 31£.

4. Interest of 62£ 2s. 4½d. for 1 mo. 20 days, at 4 per cent?

*Ans.* 6s. 10½d.

5. Interest of 85£ for 10 months 15 days, at 12½ per cent?

*Ans.* 9£ 5s. 10½d.

6. What is the amount of 53£ at 10 per cent for 7 mo.?

*Ans.* 56£ 1s. 9½d.

The time, rate per cent. and amount given, to find the principal.

79. 1. What sum of money, put at interest at 6 per cent, will amount to 61£ 4½d. in 1 year 4 months?

The amount of 1£ at the given rate and time, is 1.08£; hence,  $61.02£ \div 1.08£ = 56.50$ , the principal required; that is,—Find the amount of 1£ at the given rate and time, by which divide the given amount; the quotient will be the principal required.

*Ans.* 56£ 10s.

2. What principal, at 8 per cent, in 1 year 6 months, will amount to 85£ 2s. 4 $\frac{1}{2}$ d. ? *Ans.* 76£.

3. What principal, at 6 per cent, in 11 months 9 days, will amount to 99£ 6s. 2 $\frac{1}{2}$ d. ? *Ans.* 94£.

4. A factor receives 988£ to lay out after deducting his commission of 4 per cent; how much will remain to be laid out ?

It is evident, he ought not to receive commission on his own money. This question, therefore, in principle, does not differ from the preceding.

*Note.* In questions like this, where no respect is had to time, add the rate to 1£. *Ans.* 950£.

5. A factor receives 1008£ to lay out after deducting his commission of 5 per cent, what does his commission amount to ? *Ans.* 48£.

**DISCOUNT.** 6. Suppose I owe a man 397£ 10s. to be paid in 1 year, without interest, and I wish to pay him now; how much ought I to pay him when the usual rate is 6 per cent. ?

I ought to pay him such a sum, as put at interest, would, in 1 year, amount to 397£. 10s. The question, therefore, does not differ from the preceding. *Ans.* 375£.

*Note.* An allowance made for the payment of any sum of money before it becomes due, as in the last example, is called *Discount*.

The sum which, put at interest, would, in the time and at the rate per cent, for which discount is to be made, amount to the given sum, or debt, is called the *present worth*.

7. What is the present worth of 834£, payable in 1 year 7 months and 6 days, discounting at the rate of 7 per cent ? *Ans.* 750£.

8. What is the discount on 321£ 12s. 7 $\frac{1}{2}$ d. due 4 years hence, discounting at the rate of 6 per cent ? *Ans.* 62£ 5s. 2 $\frac{1}{2}$ d.

9. How much ready money must be paid for a note of 18£, due 15 months hence, discounting at the rate of 6 per cent ? *Ans.* 16£ 14s. 10 $\frac{1}{2}$ d.

10. Sold goods for 650£, payable one half in 4 months, and the other half in 8 months; what must be discounted for present payment ? *Ans.* 18£.

7 79.

6 months,  
*Ans.* 76£.

hs 9 days,  
*Ans.* 94£.

ducting his  
main to be

sion on his  
le, does not

it is had to  
*Ans.* 950£.

educting his  
sion amount

*Ans.* 48£.

. to be paid  
him now ;

is 6 perct.†  
rest, would,

, therefore,  
*Ans.* 375£.

of any sum  
example, is

he time and  
ade, amount

orth.

le in 1 year  
7 per cent ?

*Ans.* 750£.  
due 4 years

£ 5s. 2½d.  
or a note of

ate of 6 per  
14s. 10½d.

a 4 months,  
discounted

*Ans.* 18£.

11. What is the present worth of 56£ 4s. payable in 1 year 8 months, discounting at 6 per cent ? — at 4½ per cent ? — at 5 per cent ? — at 7 per cent ? — at 7½ per cent ? — at 9 per cent ?

*Ans. to the last* 48£ 17s. 4½d.

*The time, rate per cent., and interest being given, to find the principal.*

7 80. 1. What sum of money, put at interest 16 months, will gain 10£ 10s. at 6 per cent ?

1£ at the given rate and time, will gain '08 ; hence,  $10\cdot50\text{£} \div '08\text{£} = 131\cdot25\text{£}$  the principal required ; that is, — *Find the interest of 1£, at the given rate and time, by which divide the given gain, or interest ; the quotient will be the principal required.*

*Ans.* 131£ 5s.

2. A man paid 4£ 10s. 4½d. interest at the rate of 6 per cent at the end of 1 year 4 months ; what was the principal ?

*Ans.* 56£ 10s.

3. A man received, for interest on a certain note, at the end of 1 year, 20£ ; what was the principal, allowing the rate to have been 6 per cent ?

*Ans.* 333£ 6s. 8d.

*The principal, interest and time being given, to find the rate per cent.*

7 81. 1. If I pay 3£ 15s. 7½d. interest, for the use of 90£ for 1 year and 6 months, what is that per cent ?

The interest on 36£ at 1 per cent, the given time, is '54£ ; hence,  $3\cdot78\text{£} \div '54\text{£} = '07$ , the rate required ; that is, *Find the interest on the given sum, at 1 per cent, for the given time, by which divide the given interest ; the quotient will be the rate at which interest was paid.*

*Ans.* 7 per cent.

2. At 2£ 6s. 9½d. for the use of 468£ 1 month, what is the rate per cent ?

*Ans.* 6 per cent.

3. At 46£ 16s. for the use of 520£, 2 years, what is that per cent ?

*Ans.* 4½ per cent.

*The prices at which goods are bought and sold, being given, to find the rate per cent. of GAIN or LOSS.*

7 82. 1. If I purchase cloth at 1£ 2s. a yard, and sell it at 1£ 7s. 6d. per yard ; what do I gain per cent ?

This question does not differ essentially from those in the foregoing paragraph. Subtracting the cost from the price

at sale, it is evident I gain '275£ on a yard, that is,  $\frac{275}{1000}$  of the first cost.  $\frac{275}{1000} = 25$  per cent the answer. That is, — Make a common fraction, writing the gain or loss for the numerator, and the price at which the article was bought for the denominator; then reduce it to a decimal.

2. A merchant purchases goods to the amount of 550£; what per cent profit must he make to gain 66£?

*Ans.* 12 per cent.

3. — What per cent profit must he make on the same purchase to gain 38£ 10s.? — to gain 24£ 15s.? — to gain 2£ 15s.?

*Note.* The last gain gives for a quotient '005, which is  $\frac{1}{200}$  per cent. The rate per cent, it will be recollected, (T 75, note,) is a decimal carried to two places, or hundredths; all decimal expressions lower than hundredths are parts of 1 per cent.

4. Bought a hogshead of liquor, containing 114 gallons, at '96£ per gallon, and sold it at 1£ 0s. 0d.  $3\frac{1}{4}$  qrs. per gallon; what was the whole gain, and what was the gain per cent? *Ans.* 4£ 18s. 5½d. whole gain.  $4\frac{1}{2}$  gain per cent.

5. A merchant bought a quantity of tea for 365£, which, proving to have been damaged, he sold for 332£ 3s. what did he lose per cent? *Ans.* 9 per cent.

6. If I buy cloth at 2£ per yard, and sell it for 2£ 10s. per yard, what should I gain in laying out 100£.

*Ans.* 25£

7. Bought indigo at 6s. per lb. and sold the same at 4s. 6d. per lb.; what was the loss per cent? *Ans.* 25 per cent.

8. Bought 30 hogsheads of liquors at 600£; paid in duties 20£ 13s. 2½d.; for freight, 40£ 15s. 7½d.; for portorage, 6£. 1s.; and for insurance, 30£ 16s. 9½d.; if I sell them at 26£ per hogshead, how much shall I gain per cent?

*Ans.* 11'695 per cent.

*The principal, rate per cent, and interest being given, to find the time.*

T 83. 1. The interest on a note of 36£, at 7 per cent, was 3£ 15s. 7½d., what was the time?

The interest on 36£ 1 year, at 7 per cent, is 2£ 10s. 4½d.  $3'78£ \div 2'52£ = 1'5$  years, the time required; that is, — Find the interest for 1 year on the principal given, at the given



rate, by which divide the given interest; the quotient will be the time required, in years and decimal parts of a year; the latter may then be reduced to months and days.

Ans. 1 year 6 months.

2. If 31£ 14s. 2½d. interest be paid on a note of 226£ 10s. what was the time, the rate being 6 per cent?

Ans. 2:334=2 years 4 months.

3. A note of 600£, paid interest 20£, at 8 per cent: what was the time?

Ans. 416÷=5 months so nearly as to be called 5, and would be exactly 5 but for the fraction lost.

4. The interest on a note of 217£ 5s. at 4 per cent, was 28£ 4s. 10d.; what was the time? Ans. 3 y. 3 mo.

*Note.* When the rate is 6 per cent, we may divide the interest by  $\frac{1}{2}$  the principal, removing the separatrix two places to the left, and the quotient will be the answer in months.

The method given above, of finding the interest upon any sum in Halifax currency, for any time, and at any rate, will be found sufficiently exact in practice, and as simple and concise, perhaps, as any that could be proposed.

The teacher will do well to see that the scholar understands perfectly, the process by which the reciprocal reductions are effected by inspection, and the reason of this process.

If greater exactness be required, the reductions can be effected by the ordinary rules for the reduction of decimal fractions.

The following is a method of casting interest by vulgar fractions.

To obtain the interest upon any sum, for any time, at any rate:—Multiply the lowest terms of a fraction, the numerator of which is the given rate, and the denominator 100, by the given number of years; multiply the lowest terms of a fraction, the numerator of which is the given rate and the denominator 1200, by the given number of months; multiply the lowest terms of a fraction the numerator of which is the given rate, and the denominator 3600, by the given number of days; then reduce these several fractions to one common denominator; add them together, and by the resulting fraction multiply the given principal.

Find the interest of 100£ for 2 years, 6 months and 10 days, at 6 per cent.

$$\frac{6}{100} \text{ (or, } \frac{1}{16\frac{2}{3}}) \times 2 \text{ years} = \frac{6}{80}$$

$$\frac{6}{200} \text{ (or, } \frac{1}{33\frac{1}{3}}) \times 6 \text{ months} = \frac{6}{200}$$

$$\frac{6}{6000} \text{ (or, } \frac{1}{1000}) \times 10 \text{ days} = \frac{6}{600} = \frac{1}{100}$$

$\frac{6}{80}$ ,  $\frac{6}{200}$ ,  $\frac{6}{600}$  are to be reduced to one common denominator. Neglect the ciphers in the denominators—

$$5 \times 2 \times 6 = 60; 1 + 2 + 2 = 5, \text{ the number of ciphers.}$$

The common denominator is then 60, and 5 ciphers.

$$6 \times 2 \times 6 = 72; \text{ this with 4 ciphers is 1st numerator.}$$

$$5 \times 6 \times 6 = 180; \text{ this with 3 ciphers is 2d numerator.}$$

$$5 \times 2 \times 1 = 10; \text{ this with 3 ciphers is 3d numerator.}$$

Each numerator has as many as 3 ciphers; cut off 3 from each and 3 from the common denominator;  $\frac{72}{6000} + \frac{180}{2000} + \frac{10}{600} = \frac{80}{600} = \frac{8}{60} = \frac{2}{15}$ . Then 100£, the given principal, multiplied by  $\frac{2}{15} = \frac{200}{15} = 13\frac{1}{3}$ £ = 15£ 3s. 4d.

The reasons of the different steps in the foregoing process will appear—When the rate, as in the above example, is 6 per cent it is obvious, that the interest of any given principal for one year is  $\frac{6}{100}$ , or  $\frac{3}{50}$  of that principal. For any number of years, the interest must be as many times  $\frac{3}{50}$  of the principal, as there are units in the given number of years. In the example 2 is the given number of years; multiply then  $\frac{3}{50}$  by 2; or multiply the lowest terms of a fraction the numerator of which is the given rate, and the denominator 100, by the given number of years.  $\frac{6}{1200}$  of the given principal then is the interest for 2 years.  $\frac{6}{1200}$  of the given principal is the interest for 1 month; for there are 12 months in a year, and  $\frac{1}{1200} < \frac{1}{12} = \frac{1}{1200}$ , or  $\frac{1}{200}$ .  $\frac{6}{2000}$  of the given principal is the interest for 1 day; for there are 30 days in 1 month, and  $\frac{1}{2000} \times \frac{1}{30} = \frac{1}{6000} = \frac{1}{6000}$ . We have then  $\frac{6}{600}$  of given principal, as the interest for 1 year;  $\frac{6}{200}$  of same, for 1 month, and  $\frac{6}{6000}$  for 1 day. For 2 years, we have  $\frac{6}{600} \times 2 = \frac{12}{600}$ ; for 6 months  $\frac{6}{200} \times 6 = \frac{36}{200}$ ; for 10 day.,  $\frac{6}{6000} \times 10 = \frac{60}{6000} = \frac{1}{100}$ .  $\frac{12}{600}$ ,  $\frac{36}{200}$  and  $\frac{1}{100}$  then of the given principal are the interest of 100£ for 2 years, 6 months and 10 days. It is clear now, why we reduce these several fractions to one common denominator, add them together, and by the resulting fraction multiply the given principal.

Find the interest upon 78£ 4s. for 3 years, 9 months and 6 days, by this method, at 6 per cent, and also at 5 per cent.

To find

any p  
notes  
metho  
try, a  
tent in  
Ru  
the no  
payme  
of each  
al amo  
mount  
strum

1.  
or ord  
100

On

I  
J  
S  
J  
A

Wi

Th

which  
15s.  
for w  
132£  
for w  
10£  
time  
which  
7 mo  
which  
3d.  
ment  
date  
teres

To find the interest due on notes, &c. when partial payments have been made.

¶ 84. There is no statute in this Province, prescribing any particular form or method of casting interest upon notes or other obligations. It is believed the following method is generally allowed before the courts of the country, and also is that which has obtained to the greatest extent in mercantile transactions.

RULE.—Compute the interest upon the value for which the note, or other instrument was given, to the time of payment, which add to the principal; find the amount also of each endorsement to the time of payment, which several amounts add together, and the sum subtract from the amount of the value upon the face of the note, or other instrument.

1. For value received, I promise to pay Louis Rousseau, or order, one hundred pounds fifteen shillings, with interest.  
100£ 15s. JOHN BURTON.

May 1, 1822.

On this note, were the following endorsements:

Dec. 25, 1822,	received	10£.
July 10, 1823,	"	1£ 4s.
Sept. 1, 1824,	"	3£ 6s.
June 14, 1825,	"	21£ 15s.
April 15, 1826,	"	54£ 9s.

What was due Aug. 3, 1827? Ans. 31£ 3s. 1d.

The whole time is, from May 1st, 1822, to Aug. 3, 1827, which is 5 years, 3 months, 2 days. The interest of 100£ 15s. for this time is 31£. 15s. 4½d. This added to the value for which the note was given is, 100£ 15s. + 31£ 15s. 4½d. = 132£ 10s. 4½d. which is equal to the amount of the value for which the note was given. The first endorsement is 10£; the date of this endorsement is Dec. 25, 1822; the time of payment is Aug. 3, 1827. The time, therefore, for which interest is to be cast upon this endorsement, is 4 yrs. 7 mo. 8 days. The interest for this time is 2£ 15s. 3d. which added to the endorsement makes its amount 12£ 15s. 3d. In the same way find the amount of each other endorsement, by casting the interest upon it from the day of its date to the day of the payment of the note, and add this interest to the principal, that is, the endorsement.

The 2d endorsement is	- - - - -	12 4s.
" 3d "	- - - - -	3 6s.
" 4th "	- - - - -	21 15s.
" 5th "	- - - - -	31 2s.

The time for which interest is to be cast upon the					
2nd endorsement is	- - - -	4 years 0 months 23 day			
3d "	- - - -	2 " 11 " 2 "			
4th "	- - - -	2 " 1 " 19 "			
5th "	- - - -	1 " 3 " 18 "			

The interest upon the 2nd endorsement is		0£ 5s. 10d.
" " 3d "		0£ 11s. 6½d.
" " 4th "		2£ 15s. 8½d.
" " 5th "		4£ 4s. 11½d.

The amount of the 2nd endorsement is		1£ 9s. 10d.
" " 3d "		3£ 17s. 6½d.
" " 4th "		24£ 10s. 8½d.
" " 5th "		53£ 13s. 11½d.

The amount of 1st endorsement we found to be 12£ 15s. 3d.

The sum of the amounts of all the endorsements, 101£ 7s. 3½d.

The value upon the face of the note is 100£ 15s.

The amount of this value is, 132£ 10s. 4½d.

Subtract the sum of the amounts of the endorsements, 101£ 7s. 3½d.

Balance due Aug. 3d, 1827, 31£ 3s. 1d.

2. For value received, I promise to pay Thomas Wilson, or order, two hundred thirty-eight pounds eighteen shillings, with interest.

238£ 18s.

CHARLES STUART.

January 6, 1820.

On this note were the following endorsements: viz.

April 16, 1823, received - - - - 23£ 10s.

April 16, 1825, " - - - - 19£ 4s.

Jan. 1, 1826, " - - - - 87£ 19s.

What was due July 11, 1827?

### COMPOUND INTEREST.

185. A promises to pay B 256£ in 3 years, with in-

terest  
conve  
on the  
pal.

Now  
princi  
for bo  
at the  
Com  
the pr  
amount  
I. V  
at 6 pe

2  
3  
2  
48

2. At  
and wha  
amount  
for 6 year

It is p  
will be 2  
of 3£ w  
Hence

terest annually; but at the end of 1 year, not finding it convenient to pay the interest, he consents to pay interest on the interest from that time, the same as on the principal.

*Note.* Simple interest is that which is allowed for the principal only; compound interest is that which is allowed for both principal and interest, when the latter is not paid at the time it becomes due.

Compound interest is calculated by adding the interest to the principal at the end of each year, and making the amount the principal for the next succeeding year.

1. What is the compound interest of 256£ for 3 years, at 6 per cent?

256£ given sum or first principal.

06

15.36 interest, }  
256.00 principal, } to be added together.

271.36 amount, or principal for 2d year.

06

16.2816 compound interest 2d year, } added to-  
271.36 principal, do. } gether.

287.6416 amount, or principal for 3d year.

06

17.258496 compo. interest 3d year, } added to-  
287.641 principal, do. } gether.

304.899 amount.

256 first principal subtracted.

48,899£ compound interest for 3 years.

*Ans.* 48£ 17s. 11½d.

2. At 6 per cent, what will be the compound interest, and what the amount, of 1£ for 2 years? — what the amount for 3 years? — for 4 years? — for 5 years? — for 6 years? — for 7 years? — for 8 years?

*Ans.* to the last, 1£ 11s. 10½d.

It is plain that the amount of 2£, for any given time, will be 2 times as much as the amount of 1£; the amount of 3£ will be 3 times as much, &c.

Hence, we may form the amounts of 1£, for several years,

into a table of *multipliers* for finding the amount of any sum for the same time.

## TABLE,

Showing the amount of 1£, or 1s, &c. for any number of years; not exceeding 24, at the rates of 5 and 6 per cent compound interest.

Years.	5 per cent.	6 per cent.	Yrs.	5 per cent.	6 per cent.
1	1.05	1.06	13	1.88564	2.13292
2	1.1025	1.1236	14	1.97993	2.26090
3	1.15762	1.19101	15	2.07892	2.39655
4	1.21550	1.26247	16	2.18287	2.54035
5	1.27628	1.33822	17	2.29201	2.69277
6	1.34009	1.41851	18	2.40661	2.85483
7	1.40710	1.50363	19	2.52695	3.02559
8	1.47745	1.59384	20	2.65329	3.20713
9	1.55132	1.68947	21	2.78596	3.39956
10	1.62889	1.79084	22	2.92526	3.60353
11	1.71033	1.89829	23	3.07152	3.81974
12	1.79585	2.01219	24	3.22509	4.04893

*Note 1.* Four decimals in the above numbers will be sufficiently accurate for most operations.

*Note 2.* When there are months and days, you may first find the amount for the *years*, and on that amount cast the interest for the months and days; this, added to the amount, will give the answer.

3. What is the amount of 600£ 10s. for 20 years, at 5 per cent, compound interest? — at 6 per cent?

1£ at 5 per cent, by the table is 2.65329£; therefore,  $2.65329 \times 600.50 = 1593.30£ +$  is 1593£ 6s. *Ans.* at 5 per cent; and  $3.20713 \times 600.50 = 1925.881£ +$  is 1925£ 17s. 7½d. *Ans.* at 6 per cent.

4. What is the amount of 40£ 4s. at 6 per cent, compound interest, for 4 years? — for 10 years? — for 15 years? — for 12 years? — for 3 years and 4 months? — for 24 years 6 months and 18 days?

*Ans.* to the last 168£ 2s. 8½d.

*Note.* Any sum at compound interest will double itself in 11 years 10 months and 22 days.

From what has now been advanced we deduce the following general

I.

find th  
to tim  
Multi  
the sh  
the ra  
ing off  
are de  
mals b  
est req

II.

to find  
Multi  
inspect  
by insp  
require  
and per

III.

given t  
greate  
pound,  
for the  
for ever

IV.

inations  
necessity  
ning an  
in Halic

V. I.

per cent  
given t  
vide the  
terest, a  
terest a  
or subtr  
require.

*Note.*

at 6 per  
right ha  
terest in

of any sum

RULE.

number of  
6 per cent

6 per cent.

2:13292

2:20090

2:30655

2:54035

2:09277

2:85483

3:02559

3:20713

3:39956

3:60353

3:81974

4:04893

rs will be suf

you may first  
amount cast the  
to the amount.

20 years, at 5  
cent ?

92; therefore,

Ans. at 5 per

is 1925£ 17s

per cent, com

rs ? — for 1

and 4 months

at 168£ 2s. 8d

will double itse

deduce the fo

I. To find the interest when the time is 1 year, or, to find the rate per cent on any sum of money, without respect to time, as the premium for insurance, commission, &c.,—Multiply the principal, or given sum, after having reduced the shillings, and pence in it to the decimal of a pound, by the rate per cent, written as a decimal fraction; after pointing off as many places for decimals in the product as there are decimals in both the factors, and reducing these decimals back to shillings and pence, we shall obtain the interest required.

II. When there are months and days in the given time, to find the interest on any sum of money at 6 per cent,—Multiply the principal, reducing the shillings and pence by inspection, by the interest on 1£ for the given time, found by inspection, and the product, as before, will be the interest required, taking care to reduce the decimal parts to shillings and pence by inspection.

III. To find the interest on 1£, at 6 per cent, for any given time by inspection,—It is only to consider, that half the greatest even number of months will denote *hundredths* of a pound, and that there will be five thousandths of a pound for the odd month, (if there be one,) and one thousandth for every 6 days.

IV. If the sum given be in federal money,—The denominations being in a decimal ratio, we are saved from the necessity of effecting the reciprocal reductions, at the beginning and end of the process, otherwise proceed precisely as in Halifax currency.

V. If the interest required be at any other rate than 6 per cent, (if there be months, or months and days, in the given time,)—First find the interest at 6 per cent; then divide the interest so found by such part or parts, as the interest, at the rate required, exceeds, or falls short of the interest at 6 per cent, and the quotient, or quotients, added to or subtracted from the interest at 6 per cent, as the case may require, will give the interest at the rate required.

Note. The interest on any number of pounds, for 6 days, at 6 per cent, is readily found by cutting off the unit or right hand figure; those at the left hand will show the interest in hundredths for 6 days.



## EXAMPLES FOR PRACTICE.

1. What is the interest of 1000£ for 1 year and 3 months?

*Ans.* 120£

2. What is the interest of 5£ 16s. for 1 year 11 months?

*Ans.* 13s. 4d.

3. What is the interest of 2£ 5s. 9 $\frac{1}{2}$ d. for 1 month 19 days, at 3 per cent?

*Ans.* 2 $\frac{1}{2}$ d.

4. What is the interest of 18£ for 2 years 14 days, at 7 per cent?

*Ans.* 2£ 11s. 4 $\frac{1}{2}$ d.

5. What is the interest of 17£ 13s. 7 $\frac{1}{2}$ d. for 11 months 28 days?

*Ans.* 1£ 1s. 1d.

6. What is the interest of 200£ for 1 day? — 2 days? — 3 days? — 4 days? — 5 days?

*Ans.* for 5 days, 3s. 3 $\frac{1}{2}$ d.

7. What is the interest of half '001£ for 567 years?

*Ans.* 4d.

8. What is the interest of 81£, for 2 years 14 days, at  $\frac{1}{2}$  per cent? —  $\frac{3}{4}$  per cent? —  $\frac{1}{2}$  per cent? — 2 per cent? — 3 per cent? — 4 $\frac{1}{2}$  per cent? — 5 per cent? — 6 per cent? — 7 per cent? — 7 $\frac{1}{2}$  per cent? — 8 per cent? — 9 per cent? — 10 per cent? — 12 per cent? — 12 $\frac{1}{2}$  per cent?

*Ans.* to last, 20£ 12s. 10 $\frac{1}{2}$ d.

9. What is the interest of '09£ for 45 years, 7 months, 11 days?

*Ans.* 4s. 10 $\frac{1}{2}$ d.

10. A's note of 175£ was given Dec. 6, 1798, on which was endorsed one year's interest; what was there due Jan. 1, 1803?

*Note.* Consult ex. 16, Supplement to Subtraction of Compound Numbers.

*Ans.* 207£ 4s. 4 $\frac{1}{2}$ d.

11. B's note of 56£ 15s. was given June 6, 1801, on interest after 90 days; what was there due Feb. 9, 1802?

*Ans.* 58£ 3s. 9 $\frac{1}{2}$ d.

12. C's note of 365£ was given Dec. 3, 1797; June 7, 1800, he paid 97£ 3s. 2 $\frac{1}{2}$ d.; what was there due Sept. 11, 1800?

*Ans.* 327£ 0s. 7 $\frac{1}{2}$ d.

13. Supposing a note of 422£, dated July 5, 1797, on which were endorsed the following payments, viz. Sept. 13, 1799, 208£ 4s.; March 10, 1800, 96£; what was there due Jan. 1, 1801.

stood  
G. —  
ance  
What  
14. —  
how  
hund  
missio  
out re  
how n  
1£, at  
is inte  
given  
inter  
ney, h  
cent.,  
terest  
when p  
be four  
is unde  
the pri  
20. Ho  
goods a  
found, t  
time fou  
How m  
soning  
compou

1/ V  
days,  
2. V  
days,  
3. I  
after 3  
there d  
4. F  
terest  
was th  
5. V  
a house  
6. V

## SUPPLEMENT TO INTEREST.

## QUESTIONS.

1. What is interest? 2. How is it computed? 3. What is understood by rate per cent? 4. — by principal? 5. — by amount? 6. — by legal interest? 7. — by commission? 8. — insurance? 9. — premium? 10. — policy? 11. — stock? 12. What is understood by stock being at *par*? 13. — above *par*? 14. — below *par*? 15. The rate per cent. is a decimal carried to how many places? 16. What are decimal expressions *lower* than hundredths? 17. How is interest, (when the time is 1 year,) commission, insurance, or any thing else rated at so much per cent. without respect to time, found? 18. When the rate is one per cent., or less, how may the operation be contracted? 19. How is the interest on 1£, at 6 per cent. for any given time, found by inspection? 20. How is interest cast, at 6 per cent., when there are months and days in the given time? 21. When the given time is *less* than 6 days, how is the interest most readily found? 22. If the sum given be in federal money, how is interest cast? 23. When the rate is any other than 6 per cent., if there be months and days in the given time, how is the interest found? 24. What is the rule for casting interest on notes, &c. when partial payments have been made? 25. How may the principal be found, the time, rate per cent. and amount being given? 26. What is understood by *discount*? 27. — by *present worth*? 28. How is the principal found, the time, rate per cent. and interest being given? 29. How is the rate per cent. of gain or loss found, the prices at which goods are bought and sold being given? 30. How is the rate per cent. found, the principal, interest, and time being given? 31. How is the time found, the principal, rate per cent. and interest being given? 32. How may interest be cast by *vulgar fractions*? 33. What is the reasoning in regard to this rule? 34. What is simple interest? 35. — compound interest? 36. How is compound interest computed?

## EXERCISES

1. What is the interest of 273£ 10s. 2½d. for 1 year 10 months, at 7 per cent? *Ans.* 19£ 13s. 6¾d.  
 2. What is the interest of 486£ for 1 year, 3 months, 19 days, at 8 per cent? *Ans.* 50£ 13s. 4¾d.  
 3. D's note of 203£ was given Oct. 5, 1808, on interest after 3 months; January 5, 1809, he paid 50£; what was there due May 2, 1811? *Ans.* 175£ 7s. 2d.  
 4. E's note of 870£, was given Nov. 17, 1800, on interest after 90 days; Feb. 11, 1805, he paid 186£; what was there due Dec. 23, 1807? *Ans.* 1009£ 11s. 6¾d.  
 5. What will be the annual insurance, at ½ per cent. on a house valued at 1600£? *Ans.* 10.0£  
 6. What will be the insurance of a ship and cargo, valued

at 5643£ at  $1\frac{1}{2}$  per cent? — at  $\frac{1}{2}$  per cent? — at  $\frac{1}{4}$  per cent? — at  $\frac{1}{8}$  per cent? — at  $\frac{1}{16}$  per cent? — at  $\frac{1}{32}$  per cent?

*Note.* Consult ¶ 76, ex. 11.

*Ans.* at  $\frac{1}{2}$  per cent. 42£ 6s. 5½d.

7. A man having compromised with his creditors at 62½ per cent, what must he pay on a debt of 137£ 9s. 2½d.

*Ans.* 85£ 18s. 3d.

8. What is the value of 800£ Montreal Bank stock, at 112½ per cent?

*Ans.* 900£.

9. What is the value of 560£ 15s. of stock, at 93 per cent?

*Ans.* 521£ 9s. 11½d.

10. What principal at 7 per cent, will, in 9 months 18 days amount to 422£ 8s.

*Ans.* 400£.

11. What is the present worth of 426£, payable in 4 years and 12 days, discounting at the rate of 5 per cent?

In large sums, to bring out the hundredths and thousandths correctly, it will sometimes be necessary to extend the decimal in the divisor to five places.

*Ans.* 354£ 10s. 1½d.

12. A merchant purchased goods for 250£, ready money, and sold them again for 300£, payable in 9 months; what did he gain, discounting at 6 per cent?

*Ans.* 37£ 1s. 7½d.

13. Sold goods for 3120£, to be paid, one half in 3 months, and the other half in 6 months; what must be discounted for present payment?

*Ans.* 68£ 9s. 10d.

14. The interest on a certain note, for 1 year 9 months, was 49£ 17s. 6d., what was the principal?

*Ans.* 475£.

15. What principal, at 5 per cent., in 16 months 24 days, will gain 35£?

*Ans.* 500£.

16. If I pay 15£ 10s. interest for the use of 500£, 9 months and 9 days, what is the rate per cent?

17. If I buy candles at \$167 per lb., and sell them at 20 cents, what shall I gain in laying out \$100?

*Ans.* \$19.76.

18. Bought hats at 4s. apiece, and sold them again at 4s. 9d.; what is the profit in laying out 100£?

*Ans.* 18£ 15s.

19. Bought 37 gallons of brandy, at \$1.10 per gallon and sold it for \$40; what was gained or lost per cent?

20. At 4s. 6d. profit on 1£, how much is gained in laying out 100£, that is, how much per cent?

*Ans.* 22£ 10s.

21. Bought cloth at \$4.48 per yard; how must I sell it to gain 12½ per cent?

*Ans.* \$5.04.

— at 76  
ent 1

2£ 6s. 5½d.  
liters at 62½  
9s. 2½d.1

5£ 18s. 3d.  
nk stock, at  
Ans. 900£.

k. at 93 per  
£ 9s. 11½d.  
9 months 18

Ans. 400£.  
payable in 4  
per cent 1

thousandths  
nd the deci-  
£ 10s. 1½d.

eady money,  
onths; what  
7£ 1s. 7½d.

he half in 3  
must be dis-  
8£ 9s. 10d.

r 9 months,  
Ans. 475£.  
ths 24 days,

Ans. 500£.  
of 500£, 9

sell them at  
?  
Ans. 319.76.

again at 4s.  
s. 18£ 15s.

per gallon  
er cent?  
ined in lay-

s. 22£ 10s.  
must I sell it  
Ans. 35.04.

22. Bought a barrel of powder for 4£; for how much must it be sold to lose 10 per cent? *Ans. 3£ 12s.*

23. Bought cloth at 15s. per yard, which not proving so good as I expected, I am content to lose 17½ per cent; how must I sell it per yard? *Ans. 12s. 4½d.*

24. Bought 50 gallons of brandy, at 92 cents per gallon, but by accident 10 gallons leaked out; at what rate must I sell the remainder per gallon, to gain upon the whole cost at the rate of 10 per cent? *Ans. \$1.265 per gal.*

25. A merchant bought 10 tons of iron for 3950; the freight and duties came to \$145, and his own charges to 225; how must he sell it per lb. to gain 20 per cent by it? *Ans. 6 cents per lb.*

### EQUATION OF PAYMENTS.

786. Equation of payments is the method of finding the mean time for the payment of several debts due at different times.

1. In how many months will 1£ gain as much as 5 pounds will gain in 6 months?

2. In how many months will 1£ gain as much as 40£ will gain in 15 months? *Ans. 600.*

3. In how many months will the use of 5£ be worth as much as the use of 1£ for 40 months?

4. Borrowed of a friend 1£ for 20 months; afterwards lent my friend 4£; how long ought he to keep it to become indemnified for the use of the 1£?

5. I have three notes against a man; one of 12£, due in 3 months; one of 9£, due in 5 months; and the other of 6£, due in 10 months; the man wishes to pay the whole at once; in what time ought he to pay it?

12£ for 3 months is the same as 1£ for 36 months, and

9£ for 5 months is the same as 1£ for 45 months, and

6£ for 10 months is the same as 1£ for 60 months.

keep 27 pounds  $\frac{1}{4}$  part as long; that is,  $\frac{1}{4} \times 3 = 3$  months 6+ days, Answer.

Hence,—To find the mean time for several payments,—  
**RULE:** Multiply each sum by its time of payment, and divide the sum of the *products* by the sum of the *payments*, and the quotient will be the answer.

*Note.* This rule is founded on the supposition, that what is gained by keeping a debt a certain time *after* it is due, is the same as what is lost by paying it an equal time *before* it is due; but, in the first case, the *gain* is evidently equal to the *interest* on the debt for the given time, while, in the second case, the *loss* is only equal to the *discount* of the debt for that time, which is always *less* than the *interest*; therefore, the rule is not exactly true. The error, however, is so trifling, in most questions that occur in business, as scarce to merit notice.

6. A merchant has owing to him 300£, to be paid as follows: 50£ in 2 months, 100£ in 5 months, and the rest in 8 months; and it is agreed to make one payment of the whole: in what time ought that payment to be?

Ans. 6 months.

7. A owes B 136£, to be paid in 10 months; 96£ to be paid in 7 months; and 260£ to be paid in 4 months; what is the equated time for the payment of the whole?

Ans. 6 months, 7 days+.

8. A owes B 600£, of which 200£ is to be paid at the present time, 200 in 4 months, and 200 in 8 months; what is the equated time for the payment of the whole?

Ans. 4 months.

9. A owes B 300£, to be paid as follows:  $\frac{1}{4}$  in 3 months,  $\frac{1}{4}$  in 4 months, and the rest in 6 months: what is the equated time?

Ans.  $4\frac{1}{2}$  months.

## RATIO;

## THE RELATION OF NUMBERS.

187. 1. What part of 1 gallon is 3 quarts? 1 gallon is 4 quarts, and 3 quarts is  $\frac{3}{4}$  of 4 quarts. Ans.  $\frac{3}{4}$  of a gal.

2. What part of 3 quarts is 1 gallon? 1 gallon, being 4 quarts, is  $\frac{4}{3}$  of 3 quarts; that is, 4 quarts is 1 time 3 quarts and  $\frac{1}{3}$  of another time. Ans.  $\frac{4}{3}=1\frac{1}{3}$ .

3. What part of 5 bushels is 12 bushels?

Finding what part one number is of another is the same as finding what is called the *ratio*, or *relation* of one number to another; thus, the question, What part of 5 bushels is 12 bushels? is the same as What is the ratio of 5 bushels to 12 bushels? The answer is  $\frac{5}{12}=2\frac{1}{3}$ .

*Ratio*, therefore, may be defined, the number of times one number is contained in another; or, the number of times one quantity is contained in another quantity of the same kind.

4. What part of 8 yards is 13 yards? or, What is the ratio of 8 yards to 13 yards?

13 yards is  $\frac{13}{8}$  of 8 yards, expressing the division *fractionally*. If now we perform the division, we have for the ratio  $1\frac{5}{8}$ ; that is, 13 yards is 1 time 8 yards, and  $\frac{5}{8}$  of another time.

We have seen, (§ 15, sign,) that division may be expressed *fractionally*. So also the *ratio* of one number to another, or the part one number is of another, may be expressed *fractionally*, to do which, make the number which is called the *part*, whether it be the larger or the smaller number, the *numerator* of a fraction, under which write the other number for a denominator. When the question is, What is the ratio, &c.? the number *last* named is the *part*; consequently it must be made the *numerator* of the fraction, and the number *first* named the *denominator*.

5. What part of 12 pounds is 11 pounds? or, 11 pounds is what part of 12 pounds? 11 is the number which expresses the *part*. To put this question in the other form, viz. What is the ratio, &c. let that number, which expresses the *part*, be the number *last* named; thus, What is the ratio of 12 pounds to 11 pounds? Ans.  $\frac{11}{12}$ .

6. What part of 1£ is 2s. 6d.? or, What is the ratio of 1£ to 2s. 6d.?

1£ = 240 pence, and 2s. 6d. = 30 pence; hence,  $\frac{240}{30}=8$  is the Answer.

7. What part of 13s. 6d. is 1£ 10s. or, What is the ratio of 13s. 6d. to 1£ 10s.? Ans.  $\frac{29}{30}$ .

8. What is the ratio of 3 to 5? — of 5 to 3? — of 7



to 19? — of 19 to 77 — of 15 to 90? — of 90 to 15?  
 — of 84 to 160; — of 160 to 84? — of 615 to 1107?  
 — of 1107 to 615?      Ans. to the last  $\frac{1}{2}$ .

## PROPORTION;

OR

### THE RULE OF THREE.

¶ 88. 1. If a piece of cloth, 4 yards long, cost 12*£*, what will be the cost of a piece of the same cloth 7 yards long?

Had this piece contained twice the number of yards of the first piece, it is evident the price would have been twice as much; had it contained 3 times the number of yards, the price would have been 3 times as much; or had it contained only half the number of yards, the price would have been only half as much; that is, the cost of 7 yards will be such part of 12 pounds as 7 yards is part of 4 yards. 7 yards is  $\frac{7}{4}$  of 4 yards; consequently, the price of 7 yards must be  $\frac{7}{4}$  of the price of 4 yards, or  $\frac{7}{4}$  of 12 pounds,  $\frac{7}{4}$  of 12 pounds, that is,  $12 \times \frac{7}{4} = 21$  pounds, *answer*.

2. If a horse travel 30 miles in 6 hours, how many miles will he travel in 11 hours at that rate?

11 hours is  $\frac{11}{6}$  of 6 hours, that is, 11 hours is 1 time 6 hours, and  $\frac{5}{6}$  of another time; consequently, he will travel, in 11 hours, 1 time 30 miles, and  $\frac{5}{6}$  of another time, that is, the ratio between the distances will be equal to the ratio between the times.

$\frac{11}{6}$  of 30 miles, that is,  $30 \times \frac{11}{6} = 55$  miles. If, then, no error has been committed, 55 miles must be  $\frac{11}{6}$  of 30 miles. This is actually the case; for  $\frac{11}{6} = \frac{55}{30}$ .

Ans. 55 miles.

Quantities which have the same ratio between them are said to be *proportional*. Thus, these four quantities,

hours, hours, miles, miles,

6, 11, 30, 55,

written in this order, being such, that the second contains the first as many times as the fourth contains the third, that



is, the ratio between the third and fourth being equal to the ratio between the first and second, form what is called a proportion,

It follows, therefore, that *proportion is a combination of two equal ratios*. Ratio exists between two numbers; but proportion requires at least three.

To denote that there is a proportion between the numbers 6, 11, 30, and 55, they are written thus :—

$$6 : 11 :: 30 : 55$$

which is read, 6 is to 11 as 30 is to 55; that is, 6 is the same part of 11 that 30 is of 55; or, 6 is contained in 11 as many times as 30 is contained in 55; or, lastly, the ratio or relation of 11 to 6 is the same as that of 55 to 30.

89. The first term of a ratio, or relation, is called the *antecedent*, and the second the *consequent*. In a proportion there are two antecedents, and two consequents, viz. the antecedent of the first ratio, and that of the second; the consequent of the first ratio, and that of the second. In the proportion  $6 : 11 :: 30 : 55$ , the antecedents are 6, 30; the consequents, 11, 55.

The consequent, as we have already seen, is taken for the numerator, and the antecedent for the denominator of the fraction, which expresses the ratio or relation. Thus, the first ratio is  $\frac{11}{6}$ , the second  $\frac{55}{30} = \frac{11}{6}$ ; and that these two ratios are equal, we know, because the fractions are equal.

The two fractions  $\frac{11}{6}$  and  $\frac{55}{30}$  being equal, it follows that, by reducing them to a common denominator, the numerator of the one will become equal to the numerator of the other, and, consequently, that 11 multiplied by 30 will give the same product as 55 multiplied by 6. This is actually the case; for  $11 \times 30 = 330$ , and  $55 \times 6 = 330$ . Hence it follows, — If four numbers be in proportion, the product of the first and last, or of the two extremes, is equal to the product of the second and third, or of the two means.

Hence it will be easy, having three terms in a proportion given, to find the fourth. Take the last example. Knowing that the distances travelled are in proportion to the times or hours occupied in travelling, we write the proportion thus :

$$\begin{array}{cccc} \text{hours.} & \text{hours.} & \text{miles.} & \text{miles.} \\ 6 : 11 :: 30 : & \end{array}$$

Now, since the product of the extremes is equal to the product of the means, we multiply together the two means, 11, and 30, which makes 330, and, dividing this product by the *known* extreme, 6, we obtain for the result 55, that is 55 miles, which is the other extreme, or term, sought.

3. At 54£ for 36 barrels of flour, how many barrels may be purchased for 186£?

In this question, the unknown quantity is the number of barrels bought for 186£, which ought to contain the 36 barrels as many times as 186£ contains 54£; we thus get the following proportion:

pounds. pounds. barrels. barrels.

54 : 186 :: 36 : ....

36

1116

558

54)6696(124 barrels, the Ans.

54

129

108

216

216

The product 6696, of the two means, divided by 54, the known extreme, gives 124 barrels for the other extreme, which is the term sought, or Answer.

Any three terms of a proportion being given, the operation by which we find the fourth is called the *Rule of Three*. A just solution of the question will sometimes require, that the order of the terms of proportion be changed. This may be done, provided the terms be so placed, that the product of the extremes shall be equal to that of the means.

4. If 3 men perform a certain piece of work in 10 days, how long will it take 6 men to do the same?

The number of days in which 6 men will do the work, being the term sought, the known term of the same kind, viz. 10 days, is made the third term. The two remaining terms are 3 men and 6 men, the ratio of which is  $\frac{3}{6}$ . But the more\* men there are employed in the work, the less time will

\* The rule of three has sometimes been divided into *direct* and *inverse*, a distinction which is totally useless. It may not however be amiss to explain, in this place, in what this distinction consists.

The *Rule of Three Direct* is when more requires more, or less re

to the  
means,  
product  
55, that  
ought.  
rels may

number of  
the 36 bar-  
get the

et 6606,  
eans, di-  
54, the  
ne, gives  
for the  
, which  
sought,

e opera-  
of Three.  
ire, that  
This may  
product

10 days.

work, be-  
hind, viz.  
ng terms  
But the  
time will

cc. and in-  
pwer be  
ists.  
or less re

be required to do it ; consequently, the days will be *less* in proportion as the number of men is *greater*. There is still a proportion in this case, but the order of the terms is inverted ; for the number of men in the second set, being two times that in the first, will require only one half the time. The first number of days, therefore, ought to contain the second as many times as the second number of men contains the first. This order of the terms, being the reverse of that assigned to them in announcing the question, we say, that the number of men is in the *inverse ratio* of the number of days. With a view, therefore, to a just solution of the question, we reverse the order of the two first terms, (in doing which we invert the ratio,) and, instead of writing the proportion, 3 men : 6 men, ( $\frac{3}{6}$ ) we write it, 6 men : 3 men, ( $\frac{6}{3}$ ) that is,

men. men. days. days.  
6 : 3 :: 10 : ...

*Note.* We invert the ratio when we reverse the order of the terms in the proportion, because then the antecedent takes the place of the consequent, and the consequent that of the antecedent ; consequently, the terms of the fraction which express the ratio are inverted ; hence the ratio is inverted. Thus, the ratio expressed by  $\frac{3}{6}=2$ , being inverted, is  $\frac{6}{3}=4$ .

Having stated the proportion as above, we divide the product of the means, ( $10 \times 3=30$ , by) the known extreme, 6, which gives 5, that is, 5 days, for the other extreme, or term sought.

*Ans.* 5 days.

From the examples and illustrations now given, we deduce the following general

quires *less*, as in this example:—If 3 men dig a trench 48 feet long in a certain time, how many feet will 12 men dig in the same time? Here it is obvious, that the *more* men there are employed, the *more* work will be done ; and therefore, in this instance, *more* requires *more*. Again:—If 6 men dig 48 feet in a given time, how much will 3 men dig in the same time? Here *less* requires *less*, for the *less* men there are employed, the *less* work will be done.

The *Rule of Three Inverse* is when *more* requires *less*, or *less* requires *more*, as in this example:—If 6 men dig a certain quantity of trench in 14 hours, how many hours will it require 12 men to dig the same quantity? Here *more* requires *less* ; that is, 12 men being *more* than 6, will require *less* time. Again:—If 6 men perform a piece of work in 7 days, how long will 3 men be in performing the same work? Here *less* requires *more* ; for the number of men, being *less* will require *more* time.

## RULE.

Of the three given numbers, make that the third term which is of the same kind with the answer sought. Then consider, from the nature of the question, whether the answer will be greater or less than this term. If the answer is to be greater, place the greater of the two remaining numbers for the second term, and the less number for the first term; but if it is to be less, place the less of the two remaining numbers for the second term, and the greater for the first; and, in either case, multiply the second and third terms together, and divide the product by the first for the answer, which will always be of the same denomination as the third term.

*Note 1.* If the first and second terms contain *different* denominations, they must both be reduced to the *same* denomination.

If 8 yards of cloth cost 1£ 4s. what will 364 qrs. cost?

$$\begin{array}{rcl} \text{yds.} & \text{qrs.} & \\ 8 & : & 364 :: 1£\ 4s. \end{array}$$

Reduce 8 yards and 364 quarters to the same denomination, by dividing the 364 quarters by 4, which will bring it into yards.  $364 \div 4 = 91$ .

$$\begin{array}{rcl} \text{yds.} & \text{yds.} & \\ 8 & : & 91 :: 1£\ 4s. \end{array}$$

*Note 2.* If the third term be a *compound* number, it must either be reduced to *integers of the lowest denomination*, or the low denominations must be reduced to a *fraction of the highest denomination* contained in it.

$$\begin{array}{rcl} \text{yards.} & \text{yards.} & \\ 8 & : & 91 :: 1£\ 4s. \end{array}$$

20

24 s.

Now multiply the 24s. by 91, and divide the product by 8; the answer will be shillings, which can be reduced to pounds.

Or, the 4s. can be reduced to the fraction of a pound;  $4s. \div 20$ , that is,  $\frac{1}{5} = \frac{1}{5}$  of a pound; so  $1£\ 4s. = 1\frac{1}{5}£$ .

Or, we can reduce the 4s. to the decimal of a pound;  $20 \overline{)40}$  which, annexed to the 1£, is equal to  $1.2£$ .

2

The first method is most usually practised.

*Note 3.* The same rule is applicable, whether the given quantities be integral, fractional, or decimal.

### EXAMPLES FOR PRACTICE.

5. If 6 horses consume 21 bushels of oats in 3 weeks, how many bushels will serve 20 horses the same time ?

*Ans.* 70 bushels.

6. *The above question reversed.* If 20 horses consume 70 bushels of oats in 3 weeks, how many bushels will serve 6 horses the same time ?

*Ans.* 21 bushels.

7. If 365 men consume 75 barrels of provisions in 9 months, how much will 500 men consume in the same time ?

*Ans.* 102 $\frac{2}{3}$  barrels.

8. If 500 men consume 102 $\frac{2}{3}$  barrels of provisions in 9 months, how much will 365 men consume in the same time ?

*Ans.* 75 barrels.

9. A goldsmith sold a tankard for 10£ 12s. at the rate of 5s. 4d. per ounce ; I demand the weight of it.

*Ans.* 39 oz. 15 pwt.

10. If the moon move 13° 10' 35" in 1 day, in what time does it perform one revolution ?

*Ans.* 27 days, 7 h. 43 m.

11. If a person, whose rent is 33£, pay 3£ 2s. parish taxes, how much should a person pay whose rent is 97£ ?

*Ans.* 9£ 2s. 2 $\frac{1}{2}$ d.

12. If I buy 7lbs. of sugar for 3s. 9d., how many pounds can I buy for 1£ 10s.

*Ans.* 56 lbs.

13. If 2 lbs. of sugar cost 1s. 3d., what will 100 lbs. of coffee cost, if 8 lbs. of sugar are worth 5 lbs. of coffee ?

*Ans.* 5£.

14. If I give 6£ for the use of 100£ for 12 months, what must I give for the use of 983£ the same time ?

*Ans.* 58 $\frac{28}{100}$ £.

15. There is a cistern which has 4 pipes ; the first will fill it in 10 minutes, the second in 20 minutes, the third in 40 minutes, the fourth in 80 minutes ; in what time will all four, running together, fill it ?

$\frac{1}{10} + \frac{1}{20} + \frac{1}{40} + \frac{1}{80} = \frac{4}{80}$  cistern in 1 minute.

*Ans.* 5 $\frac{1}{4}$  minutes.

16. If a family of 10 persons spend 3 bushels of malt in a month, how many bushels will serve them when there are 30 in the family ?

*Ans.* 9 bushels.

*Note.* The rule of proportion, although of frequent use, is not of indispensable necessity; for all questions under it may be solved on general principles, without the formality of a proportion; that is, by *analysis*, as already shown, T 62, ex. 1. Thus, in the above example,—If 10 persons spend 3 bushels, 1 person, in the same time, would spend  $\frac{3}{10}$  of 3 bushels, that is,  $\frac{9}{10}$  of a bushel; and 30 persons would spend 30 times as much, that is,  $\frac{27}{10} = 2\frac{7}{10} = 2$  bushels, as before.

17. If a staff, 5 ft. 8 in. in length, cast a shadow of 6 feet, how high is that steeple whose shadow measures 153 feet?

*Ans.* 144 $\frac{1}{2}$  feet.

18. *The same by analysis.* If 6 ft. shadow require a staff of 5 ft. 8 in. = 68 in., 1 ft. shadow will require a staff of  $\frac{1}{6}$  of 68 in. or  $11\frac{1}{3}$  in.; then 153 ft. shadow will require 153 times as much; that is,  $11\frac{1}{3} \times 153 = 1734$  in. = 144 $\frac{1}{2}$  ft., as before.

19. If 3£ sterling be equal to 3 $\frac{1}{4}$ £ Halifax, how much Halifax is equal to 1000£ sterling?

*Ans.* 1111£ 2s. 2 $\frac{3}{4}$ d.

20. If 1111£ 2s. 2 $\frac{3}{4}$ d. Halifax, be equal to 1000£ sterling, how much sterling is equal to 3 $\frac{1}{4}$ £ Halifax? *Ans.* 3£.

21. If 1000£ sterling be equal to 1111£ 2s. 2 $\frac{3}{4}$ d Halifax, how much Halifax is equal to 3£ sterling? *Ans.* 3 $\frac{1}{4}$ £.

22. If 3£ sterling be equal to 3 $\frac{1}{4}$ £ Halifax, how much sterling is equal to 1111£ 2s. 2 $\frac{3}{4}$ d. Halifax.

*Ans.* 1000£.

23. Suppose 2000 soldiers had been supplied with bread sufficient to last them 12 weeks, allowing each man 14 ounces a day; but, on examination, they find 105 barrels, containing 200 lbs. each, wholly spoiled; what must the allowance be to each man, that the remainder may last them the same time?

*Ans.* 12 oz. a day.

24. Suppose 2000 soldiers were put to an allowance of 12 oz. of bread per day for 12 weeks, having a seventh part of their bread spoiled; what was the whole weight of their bread, good and bad, and how much was spoiled?

*Ans.* { The whole weight, 147000 lbs.  
Spoiled, - - - 21000 lbs.

25. — 2000 soldiers, having lost 105 barrels of bread, weighing 200 lbs. each, were obliged to subsist on 12 oz. a day for 12 weeks; had none been lost, they might have

quent use,  
ns under it  
e formality  
own, £ 62.  
sons spend  
pend  $\frac{1}{10}$  of  
sons would  
as before.

w of 6 feet,  
s 153 feet ?  
144 feet.  
quire a staff  
e a staff of  
require 153  
1734 in. =

how much  
£ 2s. 2 $\frac{3}{4}$ d.  
1000£ ster.  
Ans. 3£.  
23d Hali-  
Ans. 3 $\frac{1}{4}$ £.  
how much

Ans. 1000£.  
with bread  
ch man 14  
105 barrels,  
at must the  
ay last them  
2 oz. a day.  
allowance of  
eventh part  
ight of their  
d ?  
147000 lbs.  
21000 lbs.  
els of bread,  
on 12 oz. a  
might have

had 14 oz. a day ; what was the whole weight, including what was lost, and how much had they to subsist on ?

Ans. { Whole weight, 147000 lbs.  
Left, to subsist on, 126000 lbs.

26. — 2000 soldiers, after losing one seventh part of their bread, had each 12 oz. a day for 12 weeks ; what was the whole weight of their bread, including that lost, and how much might they have had per day, each man, if none had been lost ?

Ans. { Whole weight, 147000 lbs.  
Loss, - - 21000 lbs.  
14 oz. per day, had none been lost.

27. There was a certain building raised in 8 months by 120 workmen ; but, the same being demolished, it is required to be built in 2 months ; I demand how many men must be employed about it.

Ans. 480 men.

28. There is a cistern having a pipe which will empty it in 10 hours, how many pipes of the same capacity will empty it in 24 minutes ?

Ans. 25 pipes.

29. A garrison of 1200 men has provisions for 9 months, at the rate of 14 oz. per day ; how long will the provisions last, at the same allowance, if the garrison be reinforced by 400 men ?

Ans. 6 $\frac{1}{2}$  months.

30. If a piece of land, 40 rods in length and 4 in breadth, make an acre, how wide must it be when it is but 25 rods long ?

Ans. 6 $\frac{1}{2}$  rods.

31. If a man perform a journey in 15 days when the days are 12 hours long, in how many will he do it when the days are but 10 hours long ?

Ans. 18 days.

32. If a field will feed 6 cows 91 days, how long will it feed 21 cows ?

Ans. 26 days.

33. Lent a friend 292£ for 6 months ; some time after, he lent me 806£ ; how long may I keep it to balance the favor ?

Ans. 2 months 5 + days.

34. If 30 men can perform a piece of work in 11 days, how many men will accomplish another piece of work, 4 times as big, in a fifth part of the time ?

Ans. 600 men.

35. If  $\frac{1}{4}$  lb. of sugar cost  $\frac{1}{5}$  of a shilling, what will  $\frac{3}{4}$  of a lb. cost ?

Ans. 4d. 3 $\frac{1}{4}$ q.

Note. See ¶ 62, ex. 1, where the above question is solved by analysis. The eleven following are the next succeeding examples in the same ¶.



36. If 7 lbs. of sugar cost  $\frac{1}{2}$  of 5s., what cost 12 lbs.  
*Ans. 6s.*
37. If  $6\frac{1}{2}$  yds. of cloth cost 3£, what cost 9½ yds ?  
*Ans. 4£ 5s. 4½d.*
38. If 2 oz. of silver cost 11s. 2½d. what costs  $\frac{1}{2}$  oz. ?  
*Ans. 4s. 2½d.*
39. If  $\frac{1}{2}$  oz. cost 4½s., what costs 1 oz. ? *Ans. 6s. 5d.*
40. If  $\frac{1}{3}$  lb. less by  $\frac{1}{4}$  lb. cost 13½d., what cost 14 lbs.  
 less by  $\frac{1}{2}$  of 2 lbs. ? *Ans. 4£ 9s. 9½d.*
41. If  $\frac{3}{4}$  yd. cost  $\frac{1}{2}$ £, what will 40½ yds. cost ?  
*Ans. 59£ 1s. 2½d.*
42. If  $\frac{1}{16}$  of a ship cost 251£, what is  $\frac{3}{4}$  of her worth ?  
*Ans. 58£ 15s. 8½d.*
43. At 3½£ per cwt. what will 9½ lbs. cost ?  
*Ans. 6s. 3½d.*
44. A merchant, owning  $\frac{1}{2}$  of a vessel, sold  $\frac{1}{3}$  of his share  
 for 957£ ; what was the vessel worth ? *Ans. 1794£ 7s. 6d.*
45. If  $\frac{1}{2}$  yd. cost  $\frac{1}{2}$ £, what will  $\frac{2}{5}$  of an ell English  
 cost ? *Ans. 17s. 1d. 2½q.*
46. A merchant bought a number of bales of velvet, each  
 containing 129½ yds., at the rate of 7£ for 5 yds., and sold  
 them out at the rate of 11£ for 7 yds., and gained 200£ by  
 the bargain ; how many bales were there ? *Ans. 9 bales.*
47. At 9£ for 6 barrels of flour, what must be paid for  
 178 barrels ? *Ans. 267£.*
48. At 9s. 6d. for 3 cwt. of hay, how much is that per  
 ton ? *Ans. 3£ 3s. 4d.*
49. If 2½ lbs. of tobacco cost 75 cents, how much will  
 185 lbs. cost ? *Ans. 85.55*
50. What is the value of '15 of a hogshead of lime, at  
 11s. 11½d per hhd. ? *Ans. 1s. 9½d.*
51. If '15 of a hhd. of lime cost 1s. 9½d., what is it per  
 hhd. ? *Ans. 11s. 11½d.*

### COMPOUND PROPORTION.

T 90. It frequently happens, that the relation of the  
 quantity required, to the given quantity of the same kind,  
 depends upon several circumstances combined together ; it  
 is then called *Compound Proportion or Double Rule of Three.*

1. If a man travel 273 miles in 13 days, travelling only

This  
 W  
 answ

7 hours in a day, how many miles will he travel in 12 days, if he travel 10 hours in a day ?

This question may be solved several ways. First by analysis :—

If we know how many miles the man travelled in 1 hour it is plain, we might take this number 10 times, which would be the number of miles he would travel in 10 hours, or in 1 of these long days, and this again, taken 12 times, would be the number of miles he would travel in 12 days, travelling 10 hours each day.

If he travel 273 miles in 13 days, he travels  $\frac{1}{13}$  of 273 miles ; that is  $21\frac{1}{3}$  miles, in 1 day of 7 hours ; and  $\frac{1}{7}$  of  $21\frac{1}{3}$  miles is  $3\frac{1}{3}$  miles, the distance he travels in 1 hour : then, 10 times  $3\frac{1}{3}$  =  $33\frac{1}{3}$  miles, the distance he travels in 10 hours ; and 12 times  $33\frac{1}{3}$  =  $400$  = 360 miles, the distance he travels in 12 days, travelling 10 hours each day.

Ans. 360 miles.

But the object is to show how the question may be solved by proportion :—

First ; it is to be regarded, that the number of miles travelled over depends upon two circumstances, viz. the number of days the man travels, and the number of hours he travels each day.

We will not at first consider this latter circumstance, but suppose the number of hours to be the same in each case : the question then will be,—If a man travel 273 miles in 13 days, how many miles will he travel in 12 days ? This will furnish the following proportion :—

13 days : 12 days :: 273 miles : ..... miles

which gives for the fourth term, or answer, 252 miles.

Now, taking into consideration the other circumstance, or that of the hours, we must say,—If a man, travelling 7 hours a day for a certain number of days, travels 252 miles, how far will he travel in the same time, if he travel 10 hours in a day ?


This will lead to the following proportion :—



7 hours : 10 hours :: 252 miles : ..... miles.

This gives for the fourth term, or answer, 360 miles.




We see, then, that 273 miles has to the fourth term, or answer, the same proportion that 13 days has to 12 days,







# IMAGE EVALUATION TEST TARGET (MT-3)



Photographic  
Sciences  
Corporation

23 WEST MAIN STREET  
WEBSTER, N.Y. 14580  
(716) 872-4503

18 20 22 25  
E E E E E

1.0 0.1  
E E E E E

and that 7 hours has to 10 hours. Stating this in the form of a proportion, we have.

$$\begin{array}{l} 13 \text{ days} : 12 \text{ days} \\ 7 \text{ hours} : 10 \text{ hours} \end{array} \} :: 273 \text{ miles} : \dots\dots \text{miles}$$

by which it appears, that 273 is to be multiplied by both 12 and 10; that is, 273 is to be multiplied by the product of  $12 \times 10$ , and divided by the product of  $13 \times 7$ , which, being done, gives 360 miles for the fourth term, or answer, as before.

In the same manner, any question relating to compound proportion, however complicated, may be stated and solved.

2. If 248 men, in 5 days, of 11 hours each, can dig a trench 230 yards long, 3 wide, and 2 deep, in how many days, of 9 hours each, will 24 men dig a trench 420 yards long, 5 wide and 3 deep?

Here the number of days, in which the proposed work can be done, depends on *five circumstances*, viz. the number of men employed, the number of hours they work each day, the length, breadth, and depth of the trench. We will consider the question in relation to each of these circumstances, in the order in which they have been named:—

1st. *The number of men employed.* Were all the circumstances in the two cases alike, except the number of men and the number of days, the question would consist only in finding in how many days 24 men would perform the work which 248 men had done in 5 days; we should then have

$$24 \text{ men} : 248 \text{ men} :: 5 \text{ days} : \dots\dots \text{days.}$$

2d. *Hours in a day.* But the first laborers worked 11 hours in a day, whereas the others worked only 9; less hours will require *more* days, which will give

$$9 \text{ hours} : 11 \text{ hours} :: 5 \text{ days} : \dots\dots \text{days.}$$

3d. *Length of the ditches.* The ditches being of unequal length, as many more days will be necessary as the second is longer than the first; hence we shall have

$$230 \text{ length} : 420 \text{ length} :: 5 \text{ days} : \dots\dots \text{days.}$$

4th. *Widths.* Taking into consideration the widths, which are different, we have

$$3 \text{ wide} : 5 \text{ wide} :: 5 \text{ days} : \dots\dots \text{days.}$$

5th. *Depths.* Lastly, the depths being different, we have

$$2 \text{ deep} : 3 \text{ deep} :: 5 \text{ days} : \dots\dots \text{days.}$$



It would seem, therefore, that 5 days has to the fourth term, or answer, the same proportion

that 24 men has to 248 men, whose ratio is  $\frac{24}{248}$ ,  
 that 9 hours has to 11 hours, the ratio of which is  $\frac{9}{11}$ ,  
 that 230 length has to 420 length, - - - - -  $\frac{230}{420}$ ,  
 that 3 width has to 5 width, - - - - -  $\frac{3}{5}$ ,  
 that 2 depth has to 3 depth, - - - - -  $\frac{2}{3}$ ;  
 all which stated in form of a proportion, we have

Men,	24 : 248	} common term. :: 5 days : ..... days.
Hours,	9 : 11	
Length, 230 : 420		
Width,	3 : 5	
Depth,	2 : 3	

7 91. The continued product of all the second terms  $248 \times 11 \times 420 \times 5 \times 3$ , multiplied by the third term, 5 days, and this product divided by the continued product of the first terms,  $24 \times 9 \times 230 \times 3 \times 2$ , gives  $288\frac{11}{20}$  days for the fourth term, or answer.  $288\frac{11}{20}$ .

But the first and second terms are the fractions  $\frac{24}{248}$ ,  $\frac{9}{11}$ ,  $\frac{230}{420}$ ,  $\frac{3}{5}$  and  $\frac{2}{3}$ , which express the ratios of the men, and of the hours, of the lengths, widths and depths of the two ditches. Hence it follows, that the ratio of the number of days given to the number of days sought, is equal to the product of all the ratios, which result from a comparison of the terms relating to each circumstance of the question.

The product of all the ratios is found by multiplying together the fractions which express them, thus,  
 $\frac{248 \times 11 \times 420 \times 5 \times 3}{24 \times 9 \times 230 \times 3 \times 2} = \frac{17186400}{298080}$ , and this fraction,  $\frac{17186400}{298080}$ , represents the ratio of the quantity required to the given quantity of the same kind. A ratio resulting in this manner, from the multiplication of several ratios, is called a *compound ratio*.

From the examples and illustrations now given we deduce the following general

**RULE**

for solving questions in compound proportion, or double rule of three, viz.—Make that number which is of the same kind with the required answer, the third term; and, of the remaining numbers, take away two that are of the same

kind, and arrange them according to the directions given in simple proportion; then, any other two of the same kind, and so on till all are used.

Lastly, multiply the third term by the continued product of the second terms, and divide the result by the continued product of the first terms, and the quotient will be the fourth term, or answer required.

### EXAMPLES FOR PRACTICE.

1. If 6 men build a wall 20 feet long, 6 ft. high, and 4 ft. thick in 16 days, in what time will 24 men build one 200 ft. long, 8 ft. high, and 6 ft thick? *Ans.* 80 days.

2. If the freight of 9 hhds. of sugar, each weighing 12 cwt. 20 leagues, cost 16£, what must be paid for the freight of 50 tierces, each weighing  $2\frac{1}{2}$  cwt. 100 leagues? *Ans.* 92£ 11s. 10½d.

3. If 56 lb. of bread be sufficient for 7 men 14 days, how much bread will serve 21 men 3 days? *Ans.* 36 lbs.

*The same by analysis.* If 7 men consume 56 lbs. of bread, 1 man, in the same time, would consume  $\frac{1}{7}$  of 56 lbs. = 8 lbs.; and if he consume 8 lbs. in 14 days, he would consume  $\frac{1}{14}$  of 8 =  $\frac{4}{7}$  lb. in 1 day. 21 men would consume 21 times so much as 1 man; that is, 21 times  $\frac{4}{7}$  = 12 lbs. in 1 day, and in 3 days the would consume 3 times as much; that is,  $3 \times 12$  = equal 36 lbs. as before.

*Ans.* 36 lbs.

*Note.* Having wrought the following examples by the rule of proportion, let the pupil be required to do the same by analysis.

4. If 4 reapers receive 2£ 15s. 2½d. for 3 days' work, how many men may be hired 16 days for 25£ 15s. 2½d.? *Ans.* 7 men.

5. If 7 oz. 5 pwt. of bread be bought for 4½d. when corn is 4s. 2d. per bushel, what weight of it may be bought for 1s. 2d. when the price per bushel is 5s. 6d.? *Ans.* 1 lb. 4 oz. 3½ pwt.

6. If 100£ gain 6£ in 1 year, what will 400£ gain in 9 months? *Note.* This and the three following examples reciprocally prove each other.

7. If 100£ gain 6£ in 1 year, in what time will 400£ gain 18£?

8. If 400£ gain 18£ in 9 months, what is the rate per cent per annum?

9. What principal, at 6 per cent per annum, will gain 18£ in 9 months?

10. A usurer put out \$75 at interest, and at the end of 8 months, received, for principal and interest, \$79; find demand at what rate per cent, he received interest.

Ans. 8 per cent.

11. If 3 men receive 8 $\frac{1}{10}$ £ for 19 $\frac{1}{2}$  days' work, how much must 20 men receive for 100 $\frac{1}{2}$  days?

Ans. 305£ 0s. 8d.

## SUPPLEMENT TO THE SINGLE RULE OF THREE.

### QUESTIONS.

1. What is proportion? 2. How many numbers are required to form a ratio? 3. How many to form a proportion? 4. What is the first term of a ratio called? 5. — the second term? 6. Which is taken for the numerator, and which for the denominator of the fraction expressing the ratio? 7. How may it be known when 4 numbers are in proportion? 8. Having three terms in a proportion given, how may the fourth term be found? 9. What is the operation, by which the fourth term is found, called? 10. How does a ratio become inverted? 11. What is the rule in proportion? 12. In what denomination will the 4th term or answer, be found? 13. If the first and second terms contain different denominations, what is to be done? 14. What is compound proportion, or double rule of three? 15. Rule?

### EXERCISES.

1. If I buy 76 yards of cloth for 28£ 5s. 10d.  $\frac{1}{2}$  qrs. what does it cost per ell English? Ans. 9s. 3 $\frac{1}{2}$ d.

2. Bought 4 pieces of Holland, each containing 24 ells English for 24£; how much was that per yard? Ans. 4s.

3. A garrison had provision for 8 months, at the rate of 15 ounces to each person per day; how much must be allowed per day, in order that the provision may last 9 $\frac{1}{2}$  months? Ans. 12 $\frac{1}{2}$  oz.

4. How much land at 12s. 6d. per acre, must be given in exchange for 360 acres, at 18s. 9d. per acre? Ans. 540 acres.

5. Borrowed 185 quarters of corn when the price was 19s. ; how much must I pay when the price is 17s. 4d. ?

Ans. 202½.

6. A person owning  $\frac{1}{4}$  of a coal mine, sells  $\frac{1}{4}$  of his share for 171£ ; what is the whole mine worth ?

Ans. 380£.

7. If  $\frac{1}{4}$  of a gallon cost  $\frac{1}{4}$  of a pound, what costs  $\frac{1}{4}$  of a tun ?

Ans. 140£.

8. At 1½£ per cwt. what cost 3½ lbs. ?

Ans. 10½d.

9. If 4½ cwt. can be carried 36 miles for 35 shillings, how many pounds can be carried 20 miles for the same money ?

Ans. 907½ lbs.

10. If the sun appears to move from east to west 360 degrees in 24 hours, how much is that in each hour ? — in each minute ? — in each second ?

Ans. to the last, 15" of a deg.

11. If a family of 9 persons spend 112£ 10s. in 5 months, how much would be sufficient to maintain them 8 months if 5 persons more were added to the family ?

Ans. 280£.

*Note.* Exercises 14th, 15th, 16th, 17th, 18th, 19th and 20th, "*Supplement to Fractions*," afford additional examples in single and double proportion, should more examples be thought necessary.

## FELLOWSHIP.

T 92. 1. Two men own a farm ; the first owns  $\frac{1}{4}$ , and the second owns  $\frac{1}{4}$  of it ; the farm is sold for 40£ ; what is each man's share of the money ?

2. Two men purchase a horse for 20 pounds, of which one pays 5 pounds, and the other 15 pounds ; the horse is sold for 40 pounds ; what is each man's share of the money ?

3. A and B bought a quantity of cotton ; A paid 100 pounds, and B 200 pounds ; they sold it so as to gain 30 pounds ; what were their respective shares of the gain ?

The process of ascertaining the respective gains or losses of individuals, engaged in joint trade, is called the *Rule of Fellowship*.

The money, or value of the articles employed in trade, is called the *Capital*, or *Stock* ; the gain or loss to be shared is called the *Dividend*.

rice was  
4d. 1

. 2024½.

his share

s. 380£.

ts ½ of a

s. 140£.

ns. 10½d.

ings, how

money ?

907½ lbs.

west 360

ar 1— in

of a deg.

5 months,

8 months

ns. 280£.

19th and

examples

amples be

owns ½, and  
£; what is

of which

the horse is

the money ?

A paid 100

to gain 30

the gain ?

ins or losses

the Rule of

in trade, is

to be shared

It is plain, that *each man's* gain or loss ought to have the same relation to the *whole* gain or loss, as his *share* of the stock does to the *whole* stock.

Hence we have this **RULE** :—As the *whole* stock : to each man's *share* of the stock :: the *whole* gain or loss : his *share* of the gain or loss.

4. Two persons have a joint stock in trade ; A put in 250£, and B 350£ ; they gain 400£ ; what is each man's share of the profit ?

#### OPERATION.

A's stock,	250£	} Then,	
B's stock,	350		
			600 : 250 :: 400 : 166£ 13s. 4d. A's g.
Wholestock, 600£			600 : 350 :: 400 : 233£ 6s. 7d. B's g.

The pupil will perceive, that the process may be contracted by cutting off an equal number of ciphers from the *first* and *second*, or *first* and *third* terms; thus, 6 : 250 :: 4 : 166£ 13s. 4d. &c.

It is obvious, the correctness of the work may be ascertained by finding whether the sums of the *shares* of the gains are equal to the *whole* gain ; thus, 166£ 13s. 4d. + 233£ 6s. 7d. = 400£, the *whole* gain.

5. A, B and C trade in company ; A's capital was 175£, B's 200£, and C's 500£ ; by misfortune they lose 250£ : what loss must each sustain ?

Ans. {	50£,	A's loss.
	57£ 2s. 10½d.	B's loss.
	142£ 17s. 1½d.	C's loss.

6. Divide \$600 among 3 men so that their shares may be to each other as 1, 2, 3, respectively.

Ans. \$100, \$200, and \$300.

7. Two merchants, A and B, loaded a ship with 500 hhds. of rum ; A loaded 350 hhds. and B the rest ; in a storm the seamen were obliged to throw overboard 100 hhds. ; how much must each sustain of the loss ?

Ans. A 70, and B 30 hhds.

8. A and B companied ; A put in 45£, and took out ½ of the gain ; how much did B put in ?

Ans. 30£.

*Notes.* They took out in the same proportion as they put in ; if 3 fifths of the stock is 45£ how much is 2 fifths of it ?

9. A and B companied, and trade with a joint capital of

400£; A receives, for his share of the gain,  $\frac{1}{3}$  as much as B; what was the stock of each?

Ans.  $\begin{cases} 133\text{£ } 6\text{s. } 7\text{d. A's stock.} \\ 266\text{£ } 13\text{s. } 4\text{d. B's stock.} \end{cases}$

10. A bankrupt is indebted to B \$780, to C \$460, and to D \$760; his estate is worth only \$600; how must it be divided?

**Note.** The question evidently involves the principles of fellowship, and may be wrought by it

Ans. B \$234, C \$138, and D \$228.

11. B and C venture equal stocks in trade, and clear 164*l*; by agreement, B was to have 5 per cent of the profits, because he managed the concerns; C was to have but 2 per cent; what was each one's gain? and how much did B receive for his trouble?

Ans. B's gain was  $117\text{£ } 2\text{s. } 10\frac{1}{2}\text{d.}$  and C's  $46\text{£ } 17\text{s. } 1\frac{1}{2}\text{d.}$   
and B received  $70\text{£ } 5\text{s. } 8\frac{1}{2}\text{d.}$  for his trouble.

12. A cotton factory, valued at 12000£, is divided into 100 shares; if the profits amount to 15 per cent. yearly, what will be the profit accruing to 1 share? — to 2 shares? — to 25 shares?      Ans. to the last 450£.

13. In the above-mentioned factory, repairs are to be made which will cost 340£; what will be the tax, on each share, necessary to raise the sum? — on 2 shares? — on 3 shares? — on 10 shares?      Ans. to the last 34£.

14. If a town raise a tax of 1850£, and the whole town be valued at 37000£, what will that be on 1£? What will be the tax of a man whose property is valued at 1780£?

Ans. 1s. on a pound, and 89£ on 1780£.

¶ 93. In assessing taxes, it is necessary to have an inventory of the property, both real and personal, of the whole town, and also of the whole number of the polls; and, as the polls are rated at so much each, we must first take out from the whole tax what the polls amount to; and the remainder is to be assessed on the property. We may then find the tax upon 1£ and make a table containing the taxes on 1, 2, 3, &c. to 10£; then on 20, 30, &c. to 100£; and then on 100, 200, &c. to 1000£. Then knowing the inventory of any individual, it is easy to find the tax upon his property.

15. A certain town, valued at £64580, raises a tax of

much as

A's stock.

B's stock.

60, and to

it be di-

nciples of

D \$228.

and clear

f the prof-

have but

much did

17s. 1½d.

vided into

t. yearly.

— to 2

last 450£.

are to be

x, on each

res? —

last 34£.

hole town

What will

1780£?

on 1780£.

ave an in-

al, of the

the polls?

must first

nt to; and

We may

aining the

. to 100£;

owing the

e tax upon

a tax of

2259£ 18s.; there are 540 polls, which are taxed 3s. each; what is the tax on a pound, and what will be B's tax, whose real estate is valued at 1340£, his personal property at 874£, and who pays for 2 polls.

It will be better in questions relating to the assessment of taxes to use decimals, as we have done in interest. The process will be shorter, and the result will be obtained with exactness. The shillings, therefore, in the given values, will be reduced to the decimal of a pound, and the table will be made out decimally, and the decimal parts in the final answer can be reduced to shillings and pence.

$540 \times '60 (3s.) = 324£$ , amount of the poll taxes, and  $2259.90 (2259£ 18s.) - 324£ = 1935.90$ , to be assessed on property  $64530£ : 1935.90 :: 1£ '03$ ; or,  $\frac{1935.90}{64530} = '03$ , tax on 1£.

TABLE.

Tax on	£	£	Tax on	£	£	Tax on	£	£
1 is	'03		10 is	'30		100 is	3'	
" 2 is	'06		" 20 is	'60		" 200 is	6'	
" 3 is	'09		" 30 is	'90		" 300 is	9'	
" 4 is	'12		" 40 is	1'20		" 400 is	12'	
" 5 is	'15		" 50 is	1'50		" 500 is	15'	
" 6 is	'18		" 60 is	1'80		" 600 is	18'	
" 7 is	'21		" 70 is	2'10		" 700 is	21'	
" 8 is	'24		" 80 is	2'40		" 800 is	24'	
" 9 is	'27		" 90 is	2'70		" 900 is	27'	
						" 1000 is	30'	

Now, to find B's tax, his real estate being 1340£, I find, by the table, that

The tax on	-	-	1000£	-	-	is	-	-	30'£
The tax on	-	-	300	-	-	is	-	-	9'
The tax on	-	-	40	-	-	is	-	-	1'20

Tax on his real estate - - - - - 40'20

In like manner I find the tax on his personal

property to be	-	-	-	-	-	-	-	-	26'22
2 polls at '60 each, are	-	-	-	-	-	-	-	-	1'20

Amount, 67'62

$67.62£ = 67£ 12s. 4½d.$  Ans.

16. What will C's tax amount to, whose inventory is 874 dollars real, and 210 dollars personal property, and who pays for 3 polls? Ans. \$34.32.



17. What will be the tax of a man, paying for 1 poll, whose property is valued at \$34'82? — at \$768? — at \$940? — at \$4657? *Ans.* to the last, \$140'31.

18. Two men paid 2£ 10s. for the use of a pasture 1 month; A kept in 24 cows, and B 16 cows; how much should each pay?

19. Two men hired a pasture for 2£ 10s.; A put in 8 cows 3 months, and B put in 4 cows 4 months; how much should each pay?

T 94. The pasturage of 8 cows for 3 months is the same as 24 cows for 1 month, and the pasturage of 4 cows for 4 months is the same as of 16 cows for 1 month. The shares of A and B, therefore, are 24 to 16, as in the former question. Hence, when *time* is regarded in fellowship, — Multiply each one's stock by the time he continues it in trade, and use the product for his share. This is called *Double Fellowship*. *Ans.* A 1£ 10s. and B 1£.

20. A and B enter into partnership; A puts in 100£ 6 months, and then puts in 50£ more; B puts in 200£ 4 months, and then takes out 80£; at the close of the year, they find that they have gained 95£; what is the profit of each?

*Ans.* { 43£ 14s. 2d. A's share.  
51£ 5s. 9d. B's share.

21. A. with a capital of 500£, began trade Jan. 1, 1826, and meeting with success, took in B as a partner, with a capital of 600£, on the first of March following; 4 months after, they admit C as a partner, who brought 800£ stock; at the close of the year, they find the gain to be 700£; how must it be divided among the partners?

*Ans.* { 250£, A's share,  
250£, B's share.  
200£, C's share.

### QUESTIONS.

1. What is fellowship? 2. What is the rule for operating? 3. When time is regarded in fellowship, what is it called? 4. What is the method of operating in double fellowship? 5. How are taxes assessed? 6. How is fellowship proved?

### ALLIGATION.

T 95. Alligation is the method of mixing two or more simples, of different qualities, so that the composition may be of a mean, or middle quality.

ying for 1 poll,  
at \$768? —  
e last, \$140.31.  
of a pasture 1  
ws; how much

as.; A put in 8  
ths; how much

months is the  
average of 4 cows  
1 month. The  
as in the former  
allowalip.—Mul-  
tiple it in trade,  
is called *Double*  
10% and B 1%.  
puts in 100£ 6  
puts in 200£ 4  
loss of the year,  
at is the profit of  
2% d. A's share.  
9%. B's share.  
ad Jan. 1, 1826,  
partner, with a  
wing; 4 months  
gt 800£ stock;  
in to be 700£;  
y  
50£, A's share,  
50£, B's share.  
00£, C's share.

for operating? 3.  
alled? 4. What is  
How are taxes as-

ing two or more  
nposition may be

When the *quantities* and *prices* of the *simples* are given, to find the *mean price* of the mixture, compounded of them, the process is called *Alligation Medial*.

1. A farmer mixed together 4 bushels of wheat, worth 66 pence per bushel, 3 bushels of rye, worth 32 pence per bushel, and 2 bushels of corn, worth 28 pence per bushel; what is a bushel of the mixture worth?

It is plain, that the cost of the whole, divided by the number of bushels, will give the price of one bushel.

4 bushels,	at 66 pence,	cost	264 pence.
3 "	at 32 "	"	96
2 "	at 28 "	"	56
9 bushels cost			416 pence.

$\frac{416}{9} = 46\frac{2}{3}$  pence, *Ans.*

2. A grocer mixed 5 lbs. of sugar, worth 10 pence per lb. 8 lbs. worth 12 pence, 20 lbs. worth 14 pence; what is a pound of the mixture worth? *Ans.* 12 $\frac{1}{3}$  d.

3. A goldsmith melted together 3 ounces of gold 20 carats fine, and 5 ounces 22 carats fine; what is the fineness of the mixture? *Ans.* 21 $\frac{1}{2}$ .

4. A grocer puts 6 gallons of water into a cask containing 40 gallons of rum, worth 2s. 7d. per gallon; what is a gallon of the mixture worth? *Ans.* 2s. 2 $\frac{1}{3}$  d.

5. On a certain day the mercury was observed to stand in the thermometer as follows; 5 hours of the day, it stood at 64 degrees; 4 hours, at 70 degrees; 2 hours, at 75 degrees, and 3 hours, at 73 degrees; what was the mean temperature for that day?

It is plain this question does not differ, in the mode of its operation, from the former. *Ans.* 69 $\frac{3}{4}$  degrees.

T 96. When the *mean price* or *rate*, and the *prices* or *rates* of the *several simples* are given, to find the *proportions* or *quantities* of each simple, the process is called *Alligation Alternate*; alligation alternate is, therefore, the reverse of alligation medial, and may be proved by it.

1. A man has corn worth 40 pence per bushel, which he wishes to mix with rye worth 50 pence per bushel, so that the mixture may be worth 42 pence per bushel; what proportions, or quantities of each, must he take?

Had the price of the mixture required exceeded the price of the corn, by *just as much* as it fell short of the price of the rye, it is plain, he must have taken *equal quantities* of corn and rye; had the price of the mixture exceeded the price of the corn, by only  $\frac{1}{2}$  as much as it fell short of the price of the rye, the compound would have required 2 times as much corn as rye; and in all cases, the *less* the difference between the price of the mixture and that of one of the simples, the *greater* must be the quantity of *that simple*, in proportion to the other; that is, the quantities of the simples must be *inversely* as the *differences* of their prices from the price of the mixture; therefore, if these differences be mutually exchanged, they will, *directly*, express the *relative quantities* of each simple necessary to form the compound required. In the above example, the price of the mixture is 42 pence, and the price of the corn is 40 pence; consequently, the difference of their prices is 2 pence; the price of the rye is 50 pence, which differs from the price of the mixture by 8 pence. Therefore, by exchanging these differences, we have 8 bushels of corn to 2 bushels of rye, for the proportion required.

*Ans.* 8 bushels of corn to 2 bushels of rye, or in that proportion.

The correctness of this result may now be ascertained by the last rule; thus, the cost of 8 bushels of corn, at 40 pence is 320 pence; and 2 bushels of rye, at 50 pence, is 100 pence; then,  $320 + 100 = 420$ , and 420, divided by the number of bushels,  $(8 + 2,) = 10$ , gives 42 pence for the price of the mixture.

2. A merchant has several kinds of tea; some at 8 shillings, some at 9 shillings, some at 11 shillings, and some at 12 shillings per pound; what proportions of each must he mix, that he may sell the compound, at 10 shillings per pound?

Here we have 4 simples; but it is plain, that what has just been proved of *two* will apply to any number of *pairs*, if in each pair the price of *one* simple is *greater*, and that of the *other less*, than the price of the mixture required. Hence we have this

### RULE.

The mean rate and the several prices being reduced to the same denomination,—*connect with a continued line each*

ed the price  
the price of  
quantities of  
exceeded the  
short of the  
required 2  
the less the  
that of one  
of that simi-  
lities of the  
their prices  
re differen-  
express the  
m the com-  
price of the  
40 pence ;  
pence ; the  
n the price  
nging these  
hels of rye,

or in that  
ertained by  
at 40 pence  
ce, is 100  
y the num-  
the price of

e at 8 shil-  
nd some at  
each must  
illings per

t what has  
r of pairs,  
and that of  
required.

reduced to  
3 line each

price that is LESS than the mean rate with one or more that is GREATER, and each price GREATER than the mean rate with one or more that is LESS.

Write the difference between the MEAN rate, or price, and the price of EACH SIMPLE opposite the price with which it is connected ; (thus the difference of the two prices in each pair will be mutually exchanged ; ) then the sum of the differences, standing against any price, will express the RELATIVE QUANTITY to be taken of that price.

By attentively considering the rule, the pupil will perceive, that there may be as many different ways of mixing the simples, and consequently as many different answers, as there are different ways of linking the several prices.

We will now apply the rule to solve the last question:—

# OPERATIONS.

$$10s. \left\{ \begin{array}{l} 8s. \text{---} -2 \\ 9s. \text{---} -1 \\ 11s. \text{---} -1 \\ 12s. \text{---} -2 \end{array} \right\} \begin{array}{l} lbs. \\ -2 \\ -1 \\ -1 \\ -2 \end{array} \quad Ans. \quad \text{Or,} \quad 10s. \left\{ \begin{array}{l} 8 \text{---} -2+1=3 \\ 9 \text{---} \quad \quad 1 \quad =1 \\ 11 \text{---} \quad \quad -1+2=3 \\ 12 \text{---} \quad \quad -2 \quad =2 \end{array} \right\} \begin{array}{l} \\ \\ \\ \end{array} \quad Ans.$$

Here we set down the prices of the simples, one directly under another, in order, from least to greatest, as this is most convenient, and write the mean rate, (10s.) at the left hand. In the first way of linking, we find, that we may take in the proportion of 2 pounds of the teas at 8 and 12s. to 1 pound at 9 and 11s. In the second way, we find for the answer, 3 pounds at 8 and 11s. to 1 pound at 9 and 12s.

3. What proportions of sugar, at 8 pence, 10 pence, and 14 pence per pound, will compose a mixture worth 12 pence per pound ?

Ans. In the proportion of 2lbs. at 8 and 10 pence to 6lbs. at 14 pence.

Note. As these quantities only express the proportions of each kind, it is plain, that a compound of the same mean price will be formed by taking 3 times, 4 times, one half, or any proportion, of each quantity. Hence,

When the quantity of one simple is given, after finding the proportional quantities, by the above rule, we may say, As the PROPORTIONAL quantity : is to the GIVEN quantity : : so is each of the other PROPORTIONAL quantities : to the REQUIRED quantities of each.

4. If a man wishes to mix 1 gallon of brandy worth 16s. with rum at 9s. per gallon, so that the mixture may be worth 11s. per gallon, how much rum must he use ?

Taking the differences as above, we find the *proportions* to be 2 of brandy to 5 of rum ; consequently, 1 gallon of brandy will require  $2\frac{1}{2}$  gallons of rum ? *Ans.*  $2\frac{1}{2}$  gallons.

5. A grocer has sugars worth 7 pence, 9 pence, and 12 pence per pound, which he would mix so as to form a compound worth 10 pence per pound ; what must be the *proportions* of each kind ?

*Ans.* 2lbs. of the first and second to 4lbs. of the third kind.

6. If he use 1lb. of the first kind, how much must he take of the others ? — if 4lbs. what ? — if 6lbs. what ? — if 10lbs. what ? — if 20lbs. what ?

*Ans.* to the last, 20lbs. of the second, and 40 of third.

7. A merchant has spices at 16d. 20d. and 32d. per pound ; he would mix 5 pounds of the first sort with the others, so as to form a compound worth 24d. per pound ; how much of each sort must he use ?

*Ans.* 5lbs. of the second, and  $7\frac{1}{2}$ lbs. of the third.

8. How many gallons of water, of no value, must be mixed with 60 gallons of rum, worth 48 pence per gallon, to reduce its value to 42 pence per gallon ? *Ans.* 84 gallons.

9. A man would mix 4 bushels of wheat, at 90 pence per bushel, rye at 70 pence, corn at 70 pence, and barley at 30 pence, so as to sell the mixture at 48 pence per bushel ; how much of each may he use ?

10. A goldsmith would mix gold 17 carats fine with some 19, 21, and 24 carats fine, so that the compound may be 22 carats fine ; what proportions of each must he use ?

*Ans.* 2 of the 3 first sorts to 9 of the last.

11. If he use 1oz. of the first kind, how much must he use of the others ? What would be the quantity of the compound ? *Ans.* to last,  $7\frac{1}{2}$  ounces.

12. If he would have the whole compound consist of 15 oz., how much must he use of each kind ? — if of 30oz., how much of each kind ? — if of  $37\frac{1}{2}$ oz., how much ?

*Ans.* to last, 5oz. of the 3 first, and  $22\frac{1}{2}$ oz. of the last.

Hence, when the *quantity* of the compound is given, we may say, *As the sum of the PROPORTIONAL quantities found by the ABOVE RULE, is to the quantity REQUIRED, so is each PROPORTIONAL quantity, found by the rule, to the REQUIRED quantity of EACH.*

worth 18s.  
re may be  
use !

proportions  
1 gallon of  
2½ gallons.  
nce, and 12  
form a com-  
be the pro-

third kind.  
must he take  
what ? —

40 of third.  
per pound ;  
e others, so  
how much

f the third.  
, must be  
per gallon,  
8½ gallons.  
90 pence  
and barley  
e per bush-

with some  
d may be  
e use !

f the last.  
h must he  
ty of the  
½ ounces.  
sist of 15  
of 30oz.,  
much ?

f the last.  
iven, we  
ies found  
so is each  
REQUIR-

13. A man would mix 100 pounds of sugar, some at 8 pence, some at 10 pence, and some at 14 pence per pound, so that the compound may be worth 12 pence per pound ; how much of each kind must he use ?

We find the proportions to be, 2, 2, and 6. Then,  $2+2+6=10$ , and

$$10 : 100 :: \left\{ \begin{array}{l} 2 : 20\text{lbs. at } 8\text{d.} \\ 2 : 20\text{lbs. at } 10\text{d.} \\ 6 : 60\text{lbs. at } 14\text{d.} \end{array} \right\} \text{Ans.}$$

14. How many gallons of water, of no value, must be mixed with brandy at 120 pence per gallon, so as to fill a vessel of 75 gallons, which may be worth 92 pence per gallon ?

Ans.  $17\frac{1}{2}$  gallons of water to  $57\frac{1}{2}$  gallons of brandy.

15. A grocer has currants at 4d., 6d., 9d. and 11d. per lb. ; and he would make a mixture of 240lbs., so that the mixture may be sold at 8d. per lb. ; how many pounds of each sort may he take ?

Ans. 72, 24, 48, and 96lbs., or 48, 48, 72, 72, &c.

Note. This question may have five different answers.

### QUESTIONS.

1. What is alligation ? 2. — medial ? 3. — the rule for operating ? 4. What is alligation alternate ? 5. When the price of the mixture, and the price of the several simples, are given, how do you find the *proportional quantities* of each simple ? 6. When the quantity of one simple is given, how do you find the others ? 7. When the quantity of the whole compound is given, how do you find the quantity of each simple ?

## DUODECIMALS.

T 97. Duodecimals are fractions of a foot. The word is derived from the Latin word *duodecim*, which signifies *twelve*. A foot, instead of being divided *decimally* into *ten* equal parts, is divided *duodecimally* into *twelve* equal parts, called *inches*, or *primes*, marked thus, ('). Again, each of these parts is conceived to be divided into twelve other equal parts, called *seconds*, ("). In like manner, each second is conceived to be divided into twelve equal parts, called *thirds*, ("""); each third into twelve equal parts, called *fourths*, ("""); and so on to any extent.

In this way of dividing a foot, it is obvious, that

1' inch or prime is  $-\ - - - - \frac{1}{12}$  of a foot.  
 1'' second is  $\frac{1}{12}$  of  $\frac{1}{12} - - - = \frac{1}{144}$  of a foot.  
 1''' third is  $\frac{1}{12}$  of  $\frac{1}{12}$  of  $\frac{1}{12} - - = \frac{1}{1728}$  of a foot.  
 1'''' fourth is  $\frac{1}{12}$  of  $\frac{1}{12}$  of  $\frac{1}{12}$  of  $\frac{1}{12} = \frac{1}{20736}$  of a foot.  
 1''''' fifth is  $\frac{1}{12}$  of  $\frac{1}{12}$  of  $\frac{1}{12}$  of  $\frac{1}{12}$  of  $\frac{1}{12} = \frac{1}{248832}$  of a foot, &c.

Duodecimals are added and subtracted in the same manner as compound numbers, 12 of a less denomination making 1 of a greater, as in the following

## TABLE.

12'''' fourths make 1''' third,  
 12''' thirds  $- - -$  1'' second,  
 12'' seconds  $- -$  1' inch or prime,  
 12' inches, or primes, 1 foot.

*Note.* The marks, ', ', ', ', &c. which distinguish the different parts, are called the *indices* of the *parts* or *denominations*.

## MULTIPLICATION OF DUODECIMALS.

Duodecimals are chiefly used in measuring *surfaces* and *solids*.

1. How many square feet in a board 16 feet 7 inches long, and 1 foot 3 inches wide?

*Note.* Length  $\times$  breadth = superficial contents, (T 25.)  
OPERATION.

Length,	16	7'
Breadth,	1	3'
	4	1' 9''
	16	7'

*Ans.* 20 8' 9''

proper place. To multiply 16 feet by 3' is to take  $\frac{3}{12}$  of  $\frac{1}{12}$  of  $\frac{1}{12}$ , that is, 48'; and the 1' which we reserved makes 49' = 4 feet 1'; we therefore set down the 1', and carry forward the 4 feet to its proper place. Then, multiplying the multiplicand by the 1 foot in the multiplier, and adding the two products together, we obtain the *answer*, 20ft, 8' 9''.

The only difficulty that can arise in the multiplication of duodecimals is, in finding of what denomination is the pro-

7 inches, or primes =  $\frac{7}{12}$  of a foot, and 3 inches =  $\frac{3}{12}$  of a foot; consequently, the product of  $7' \times 3' = \frac{21}{144}$  of a foot, that is, 21'' = 1' and 9''; wherefore, we set down the 9'', and reserve the 1' to be carried forward to its

Length  
Breadth

Thick

For  
Writing  
in m  
nomi  
by th



foot.  
foot.  
foot.  
foot.  
foot, &c.

me man-  
ion mak-

me,

guish the  
r denomi-

LS.

faces and

7 inches

(† 25.)

$\frac{7}{2}$  of a  
 $\frac{3}{2}$  of a foot;  
product of 7'  
that is, 21"  
ore, we set  
eserve the  
ward to its  
ke  $\frac{3}{2}$  of  $\frac{1}{2}$   
ed makes  
and carry  
multiplying  
and adding  
20ft, 8' 9".  
plication of  
is the pro-

duct of any two denominations. This may be ascertained as above, and in all cases it will be found to hold true, that the product of any two denominations will always be of the denomination denoted by the sum of their INDICES. Thus, in the above example, the sum of the indices of 7'  $\times$  3' is " ; consequently, the product is 21" ; and thus primes multiplied by primes will produce seconds ; primes multiplied by seconds produce thirds ; fourths multiplied by fifths produce ninths, &c.

It is generally most convenient, in practice, to multiply the multiplicand first by the feet of the multiplier, then by the inches, &c. thus :—

$$\begin{array}{r} \text{ft.} \\ 16 \text{ } 7' \\ 1 \text{ } 3' \\ \hline 16 \text{ } 7' \\ 4 \text{ } 1' \text{ } 9'' \\ \hline 20 \text{ } 8' \text{ } 9'' \end{array}$$

16 ft.  $\times$  1 ft. = 16 ft., and 7'  $\times$  1 ft. = 7'. Then, 16 ft.  $\times$  3' = 48' = 4 ft., and 7'  $\times$  3' = 21" = 1' 9". The two products added together, give for the answer. 20 ft. 8' 9", as before.

2. How many solid feet in a block 15 ft. 8' long, 1 ft. 5' wide, and 1ft. 4' thick ?

OPERATION.

$$\begin{array}{r} \text{ft.} \\ \text{Length, } 15 \text{ } 8' \\ \text{Breadth, } 1 \text{ } 5' \\ \hline 15 \text{ } 8' \\ 6 \text{ } 6' \text{ } 4'' \\ \hline 22 \text{ } 2' \text{ } 4'' \\ \text{Thickness } 1 \text{ } 4' \end{array}$$

$$\begin{array}{r} 22 \text{ } 2' \text{ } 4'' \\ 7 \text{ } 4' \text{ } 9'' \text{ } 4''' \\ \hline \text{Ans. } 29 \text{ } 7' \text{ } 1'' \text{ } 4''' \end{array}$$

The length multiplied by the breadth, and that product by the thickness, gives the solid contents, († 33.)

From these examples we derive the following RULE :— Write down the denominations as compound numbers, and in multiplying remember, that the product of any two denominations will always be of that denomination denoted by the sum of their indices.

### EXAMPLES FOR PRACTICE.

3. How many square feet in a stock of 15 boards, 12 ft. 8' in length, and 13' wide? Ans. 205 ft. 10'.

4. What is the product of 371ft 2' 6" multiplied by 181 ft. 1' 9" ? Ans. 67242 ft. 10' 1" 4''' 6'''.

*Note.* Painting, plastering, paving, and some other kinds of work, are done by the square yard. If the contents in square feet be divided by 9, the quotient, it is evident, will be square yards.

5. A man painted the walls of a room 8 ft. 2' in height, and 72 ft. 4' in compass; (that is, the measure of all its sides;) how many square yards did he paint?

6. How many cord feet of wood in a load 8 feet long, 4 feet wide, and 3 feet 6 inches high?

7. In a pile of wood 176 ft. in length, 3 ft. 9' wide, and 4 ft. 3' high, how many cords?

8. How many cord feet of wood in a load 7 feet long, 3 feet wide, and 3 feet 4 inches high? and what will it come to at 2s. per cord foot?

9. How much wood in a load 10 ft. in length, 3 ft. 9' in width, and 4 ft. 8' in height? and what will it cost at \$1.92 per cord?

**¶ 98. Remark.** By some surveyors of wood, dimensions are taken in feet and *decimals* of a foot. For this purpose, make a rule or scale 4 feet long, and divide it into feet and each foot into 10 equal parts. On one end of the rule for 1 foot, let each of these parts be divided into 10 other equal parts. The former division will be 10ths, and the latter 100ths of a foot. Such a rule will be found very convenient for surveyors of wood and of lumber, for painters, joiners, &c.; for the dimensions taken by it being in feet and decimal parts of a foot, the casts will be no other than so many operations in decimal fractions.

10. How many square feet in a hearth stone, which, by a rule, as above described, measures 4'5 feet in length, and 2,6 feet in width ? and what will be its cost, at 75 cents per

ards, 12 ft.  
205 ft. 10'.  
ied by 181  
" 4" 6"  
other kinds  
contents in  
vident, will

in height,  
e of all its

. 5 ft. 8' 8"  
feet long, 4

feet make a  
7 cord feet.  
9' wide, and

1½ cord feet.  
feet long, 3  
will it come

me to 8s. 9d.  
th, 3 ft. 9' in  
cost at \$1.92

ne to \$2.62½.

wood, dimen-  
For this pur-  
de it into feet

d of the rule  
nto 10 other

e, and the lat-  
nd very con-

for painters,  
being in feet

no other than

, which, by a  
length, and  
t 75 cents per

square foot 1      *Ans.* 11'7 feet; and it will cost \$8'775.

11. How many cords in a load of wood 7'5 feet in length,  
3'6 feet in width, and 4'8 feet in height? *Ans.* 1 cord 1½ ft.

12. How many cord feet in a load of wood 10 feet long,  
3'4 feet wide, and 3'5 feet high? *Ans.* 7½.

### QUESTIONS.

1. What are duodecimals? 2. From what is the word *derived*?
3. Into how many parts is a foot usually divided, and what are the parts called? 4. What are the *other* denominations? 5. What is understood by the *indices* of the denominations? 6. In what are duodecimals chiefly used? 7. How are the contents of a *surface* bounded by straight lines found? 8. How are the contents of a *solid* found? 9. How is it known of what denomination is the product of any two denominations? 10. How may a scale or rule be formed for taking dimensions in feet and decimal parts of a foot?

## INVOLUTION.

7 99. Involution, or the raising of powers, is the multiplying any given number into itself continually a certain number of times. The products thus produced are called the *powers* of the given number. The number itself is called the *first* power or *root*. If the *first* power be multiplied by *itself*, the product is called the *second* power or *square*: if the square be multiplied by the first power, the product is called the *third* power, or *cube*, &c. thus,

5 is the root, or 1st power of 5.

$5 \times 5 = 25$  is the 2d power, or square, of 5,  $= 5^2$

$5 \times 5 \times 5 = 125$  is the 3d power, or cube, of 5,  $= 5^3$

$5 \times 5 \times 5 \times 5 = 625$  is the 4th power, or biquadrate, of 5,  $= 5^4$

The number denoting the power is called the *index*, or *exponent*; thus,  $5^4$  denotes that 5 is raised or involved to the 4th power.

1. What is the square, or 2d power of 7? *Ans.* 49.

2. What is the square of 30? *Ans.* 900.

3. What is the square of 4000? *Ans.* 16000000.

4. What is the cube, or 3d power, of 4? *Ans.* 64.

5. What is the cube of 800? *Ans.* 512000000.

6. What is the 4th power of 60? *Ans.* 12960000.

7. What is the square of 1? — of 2? — of 3? —  
of 4? *Ans.* 1, 4, 9, and 16.

8. What is the cube of 11 — of 21 — of 31 — of 41? *Ans.* 1, 8, 27, and 64.

9. What is the square of  $\frac{2}{3}$  1 — of  $\frac{4}{5}$  1 — of  $\frac{7}{8}$  1? *Ans.*  $\frac{4}{9}$ ,  $\frac{16}{25}$ ,  $\frac{49}{64}$ .

10. What is the cube of  $\frac{2}{3}$  1 — of  $\frac{4}{5}$  1 — of  $\frac{7}{8}$  1? *Ans.*  $\frac{8}{27}$ ,  $\frac{64}{125}$ , and  $\frac{343}{512}$ .

11. What is the square of  $\frac{1}{2}$  1 — the 5th power of  $\frac{1}{2}$  1? *Ans.*  $\frac{1}{4}$ , and  $\frac{1}{32}$ .

12. What is the square of 1'5 1 — the cube 1? *Ans.* 2'25, and 3'375.

13. What is the 6th power of 1'2 1? *Ans.* 2'985984.

14. Involve  $2\frac{1}{2}$  to the 4th power.

*Note.* A mixed number, like the above, may be reduced to an improper fraction before involving: thus,  $2\frac{1}{2} = \frac{5}{2}$ ; or it may be reduced to a decimal; thus,  $2\frac{1}{2} = 2.5$ .

15. What is the value of  $7^4$ , that is, the 4th power of 7 1? *Ans.*  $7^4 = 2401$ .

16. How much is  $9^3$  1 —  $6^5$  1 —  $10^4$  1? *Ans.* 729, 7776, 10000.

17. How much is  $2^7$  1 —  $3^6$  1 —  $4^5$  1 —  $5^3$  1 —  $6^5$  1 —  $10^3$  1? *Ans.* to the last, 100000000.

The powers of the nine digits, from the first power to the fifth, may be seen in the following

TABLE.

Roots	or 1st Powers	1	2	3	4	5	6	7	8	9
Squares	or 2d Powers	1	4	9	16	25	36	49	64	81
Cubes	or 3d Powers	1	8	27	64	125	216	343	512	729
Biquadrates	or 4th Powers	1	16	81	256	625	1296	2401	4096	6561
Quintals	or 5th Powers	1	32	243	1024	3125	7776	16807	32768	59049

## EVOLUTION.

100. Evolution, or the *extracting* of roots, is the method of finding the root of any power or number.

The *root*, as we have seen, is that number, which, by a continual multiplication into itself, produces the given power. The *square* root is a number which, being squared, will produce the given number; and the *cube*, or *third* root, is a number, which being cubed or involved to the 3d power will produce the given number: thus, the *square* root of 144

99, 100.

37 —  
and 64.

$\frac{1}{4}$ ,  $\frac{1}{8}$ ,  $\frac{1}{16}$ ,  
 $\frac{1}{32}$ ,  $\frac{1}{64}$ ,  
and  $\frac{1}{128}$ .  
power of  $\frac{1}{2}$ ?  
and  $\frac{1}{3}$ .

and 3.375.  
2965984.

be reduced  
 $2\frac{1}{2} = \frac{5}{2}$ ; or

$2\frac{1}{2} = 2.5$   
power of 7?  
Ans. 2401.

776, 10000.  
—  $5^3$  ? —

100000000.  
power to the

71	81	9
91	64	81
31	512	729
01	4096	6561
07	32768	59049

pts, is the me-  
ber.

, which, by a  
ne given pow-  
squared, will  
third root, is a  
the 3d power  
are root of 14

is 12, because  $12^2=144$ ; and the *cube root* of 343 is 7, be-  
cause  $7^3$ , that is,  $7 \times 7 \times 7=343$ ; and so of other numbers.

Although there is no number which will not produce a  
perfect power by involution, yet there are many numbers of  
which *precise roots* can never be obtained. But by the help  
of *decimals*, we can approximate, or approach, towards the  
root to any assigned degree of exactness. Numbers, whose  
precise roots cannot be obtained, are called *surd numbers*,  
and those whose roots can be exactly obtained, are called  
*rational numbers*.

The square root is indicated by this character  $\sqrt{\quad}$  placed  
before the number; the other roots by the same character  
with the index of the root placed over it. Thus, the square  
root of 16 is expressed  $\sqrt{16}$ ; and the cube root of 27 is  
expressed  $\sqrt[3]{27}$ ; and the 5th root of 7776,  $\sqrt[5]{7776}$ .

When the power is expressed by several numbers, with  
the sign + or — between them, a line, or *vinculum*, is drawn  
from the top of the sign over all the parts of it; thus, the  
square root of  $21-5$  is  $\sqrt{21-5}$ , &c.

## EXTRACTION OF THE SQUARE ROOT.

¶ 101. To extract the square root of any number is to  
find a number, which, being multiplied into itself, shall pro-  
duce the given number.

1. Supposing a man has 625 yards of carpeting, a yard  
wide, what is the length of one side of a square room, the  
floor of which the carpeting will cover? that is, what is  
one side of a square, which contains 625 square yards?

We have seen, (¶ 32,) that the contents of a square sur-  
face is found by multiplying the length of one side into it-  
self, that is, by raising it to the second power; and hence,  
having the contents, (625) given, we must extract its *square*  
root to find one side of the room.

This we must do by a sort of trial: and,

1st. We will endeavor to ascertain how many figures  
there will be in the root. This we can easily do, by point-  
ing off the number, from units, into periods of two figures  
each; for the square of any root always contains just twice

as many, or one figure *less* than twice as many figures, as are in the root; of which truth the pupil may easily satisfy himself by trial. Pointing off the number, we find, that

## OPERATION.


625(2

4

---

225

FIG. 1.



the root will consist of *two* figures, a ten and a unit.

2d. We will now seek for the first figure, that is, for the *tens* of the root, and it is plain that we must extract it from the left hand period 6, (hundreds.) The greatest square in 6 (hundreds) we find, by trial, to be 4, (hundreds,) the root of which is 2, (tens, = 20;) therefore, we set 2 (tens) in the root. The root, it will be recollected, is *one side* of a square. Let us, then, form a square, (A. Fig. 1.) each side of which shall be supposed 2 tens, = 20 yards, expressed by the root now obtained.

The contents of this square are  $20 \times 20 = 400$  yards, now disposed of, and which, consequently, are to be deducted from the whole number of yards, (625,) leaving 225 yards. This deduction is most readily performed by subtracting the square number 4, (hundreds,) or the square of 2, (the figure in the root already found,) from the period 6, (hundreds,) and bringing down the next period by the side of the remainder making 225, as before.

3d. The square A is now to be enlarged by the addition of the 225 remaining yards; and, in order that the figure may retain its *square form*, it is evident, the addition must be made on *two sides*. Now, if the 225 yards be divided by the *length* of the *two sides*, ( $20 + 20 = 40$ ,) the quotient will be the *breadth* of this new addition of 225 yards to the sides *c d* and *b c* of the square A.

But our root already found, = 2 tens, is the length of *one side* of the figure A; we therefore take *double* this root, = 4 tens, for a divisor.

5 yds.

20 yds.

a

which  
add th  
the rig  
whole  
the co  
figure  
our div  
Conseq  
25 yard  
will ex

The  
parts of  
The squ  
The fig


"

"

## OPERATION--CONTINUED.

$$\begin{array}{r}
 625 \overline{)25} \\
 \underline{4} \phantom{00} \\
 45 \overline{)225} \\
 \underline{225} \\
 0
 \end{array}$$

FIG. II.



which is equal to this last quotient, 5; we must, therefore, add this quotient, 5, to the divisor, 40, that is, place it as the right hand of the 4 (tens), making it 45; and then the whole divisor, 45, multiplied by the quotient, 5, will give the contents of the whole addition around the sides of the figure A, which, in this case, being 225 yards, the same as our dividend, we have no remainder, and the work is done. Consequently, Fig. II. represents the floor of a square room, 25 yards on a side, which 625 square yards of carpeting will exactly cover.

The proof may be seen by adding together the several parts of the figure, thus:—

The square A contains 400 yards.

The figure B " 100 "

" " C " 100 "

" " D " 25 "

Proof. 625 "

S

Or we may prove it by involution, thus:—  
 $25 \times 25 = 625$ , as before.

The divisor, 4, (tens) is in reality 40, and we are to seek how many times 40 is contained in 225, or, which is the same thing, we may seek how many times 4 (tens) is contained in 22, (tens,) rejecting the right hand figure of the dividend, because we have rejected the cipher in the divisor. We find our quotient, that is, the breadth of the addition, to be 5 yards; but, if we look at Fig. II., we shall perceive that this addition of 5 yds. to the two sides does not complete the square; for there is still wanting, in the corner D, a small square, each side of



From this example and illustration we derive the following general

### RULE

*For the extraction of the Square Root.*

I. Point off the given number into periods of two figures each, by putting a dot over the units, another over the hundreds, and so on. These dots show the number of figures of which the root will consist.

II. Find the greatest square number in the left hand period, and write its root as a quotient in division. Subtract the square number from the left hand period, and to the remainder bring down the next period for a dividend.

III. Double the root already found for a divisor; seek how many times the divisor is contained in the dividend, excepting the right hand figure, and place the result in the root, and also at the right hand of the divisor; multiply the divisor, thus augmented, by the last figure of the root, and subtract the product from the dividend; to the remainder bring down the next period for a new dividend.

IV. Double the root already found for a new divisor, and continue the operation as before, until all the periods are brought down.

*Note 1.* If we double the right hand figure of the last divisor, we shall have the double of the root.

*Note 2.* As the value of figures, whether integers or decimals, is determined by their distance from the place of units, so we must always begin at unit's place to point off the given number, and, if it be a mixed number, we must point it off both ways from units, and if there be a deficiency in any period of decimals, it may be supplied by a cipher. It is plain, the root must always consist of so many integers and decimals as there are periods belonging to each in the given number.

### EXAMPLES FOR PRACTICE.

3. What is the square root of 10342656?

Y 101.  
follow

o figures  
over the  
r of fig-  
  
hand pe-  
Subtract  
o the re-  
l.  
or; seek  
dividend,  
ult in the  
ultiply the  
root, and  
remainder  
  
w divisor,  
he periods  
  
f the last  
  
ntegers or  
e place of  
point off  
we must  
a deficien-  
a cipher.  
y integers  
ach in the

OPERATION.

$$\begin{array}{r} 10342656 \text{ (3216, Ans.} \\ 9 \\ \hline 62 \text{ ) } 134 \\ 124 \\ \hline 641 \text{ ) } 1026 \\ 641 \\ \hline 6426 \text{ ) } 38556 \\ 38556 \\ \hline \end{array}$$

3. What is the square root of 43264 ?

OPERATION.

$$\begin{array}{r} 43264 \text{ (208, Ans.} \\ 4 \\ \hline 408 \text{ ) } 3264 \\ 3264 \\ \hline \end{array}$$

- |  |                |
|--|----------------|
| 4. What is the square root of 998001 ?         | Ans. 999.      |
| 5. What is the square root of 234'09 ?         | Ans. 15'3.     |
| 6. What is the square root of 964'5192360241 ? | Ans. 31'05671. |
| 7. What is the square root of '001296 ?        | Ans. '036.     |
| 8. What is the square root of '2916 ?          | Ans. '54.      |
| 9. What is the square root of 36372961 ?       | Ans. 6031.     |
| 10. What is the square root of 164 ?           | Ans. 12'8+.    |

Y 102. In this last example, as there was a remainder, after bringing down all the figures, we continued the operation to decimals, by annexing two ciphers for a new period, and thus we may continue the operation to any assigned degree of exactness; but the pupil will readily perceive, that he can never in this manner, obtain the *precise* root; for the last figure in each *dividend* will always be a cipher, and the last figure in each *divisor* is the same as the last *quotient* figure; but no one of the nine digits, multiplied into itself,

produces a number ending with a *cipher*; therefore, whatever be the quotient figure, there, will still be a remainder.

11. What is the square root of 3? *Ans.* 1.73+.  
 12. What is the square root of 10? *Ans.* 3.16+.  
 13. What is the square root of 18421 *Ans.* 1357+.  
 14. What is the square root of  $\frac{1}{4}$ ?

*Note.* We have seen, (¶ 90, ex. 9.) that fractions are squared by squaring both the numerator and the denominator. Hence it follows, that the square root of a fraction is found by extracting the root of the numerator and of the denominator. The root of 4 is 2, and the root of 9 is 3.

- Ans.*  $\frac{1}{2}$ .  
 15. What is the square root of  $\frac{1}{16}$ ? *Ans.*  $\frac{1}{4}$ .  
 16. What is the square root of  $\frac{1}{100}$ ? *Ans.*  $\frac{1}{10}$ .  
 17. What is the square root of  $\frac{81}{144}$ ? *Ans.*  $\frac{9}{12} = \frac{3}{4}$ .  
 18. What is the square root of  $20\frac{1}{4}$ ? *Ans.* 4 $\frac{1}{2}$ .

When the numerator and denominator are not exact squares, the fraction may be reduced to a decimal, and the approximate root found, as directed above.

19. What is the square root of  $\frac{3}{4} = .75$ ? *Ans.* .866+.  
 20. What is the square root of  $\frac{9}{16}$ ? *Ans.* .912+.

## SUPPLEMENT TO THE SQUARE ROOT.

### QUESTIONS.

1. What is involution? 2. What is understood by a power? 3. — the first, the second, the third, the fourth power? 4. What is the index, or exponent? 5. How do you involve a number to any required power? 6. What is evolution? 7. What is a root? 8. Can the precise root of all numbers be found? 9. What is a surd number? 10. — a rational? 11. What is it to extract the square root of any number? 12. Why is the given number divided into periods of two figures each? 13. Why do we double the root for a divisor? 14. Why do we, in dividing, reject the right hand figure of the dividend? 15. Why do we place the quotient figure to the right hand of the divisor? 16. How may we prove the work? 17. Why do we point off mixed numbers both ways from units? 18. When there is a remainder, how may we continue the operation? 19. Why can we never obtain the precise root of surd numbers? 20. How do we extract the square root of vulgar fractions?

### EXERCISES.

1. A general has 4006 men; how many must he place

c, what-  
remainder.

1.73+.  
3.16+.  
3.57+.

ions are  
nomina-  
action in  
the de-  
s.

Ans. 7.  
Ans. 5.  
Ans. 10.  
8 = 3.  
Ans. 41.  
ot exact  
and the

806+.  
912+.

OT.

er? 3.  
What is  
to any  
ot? 8.  
a surd  
square  
periods  
visor?  
o divi-  
and of  
do we  
here is  
an we  
ve ex-

in rank and file to form them into a square 1 *Ans. 64.*

2 If a square field contains 2025 square rods, how many rods does it measure on each side? *Ans. 45.*

3. How many trees in each row of a square orchard containing 5025 trees? *Ans. 75.*

4. There is a circle whose area, or superficial contents, is 5184 feet; what will be the length of the side of a square a equal area?  $\sqrt{5184}=72$  feet, *Ans.*

5. A has two fields, one containing 40 acres, and the other containing 50 acres, for which B offers him a square field containing the same number of acres as both of these; how many rods must each side of this field measure?

*Ans. 120 rods.*

6. If a certain square field measure, 20 rods on each side, how much will the side of a square field measure, containing 4 times as much?  $\sqrt{20 \times 20 \times 4}=40$  rods' *Ans.*

7. If the side of a square be 5 feet, what will be the side of one 4 times as large? — 9 times as large? — 16 times as large? — 25 times as large? — 36 times as large? *Ans. 10ft. 15ft. 20ft. 25ft. and 30ft.*

8. It is required to lay out 288 rods of land in the form of a parallelogram, which shall be twice as many rods in length as it is in width.

*Note.* If the field be divided in the middle, it will form two equal squares. *Ans. 24 rods long, and 12 rods wide.*

9. I would set out, at equal distances, 784 apple trees, so that my orchard may be 4 times as long as it is broad; how many rows of trees must I have, and how many trees in each row? *Ans. 14 rows, and 56 in each row.*

10. There is an oblong piece of land, containing 192 square rods, of which the width is  $\frac{3}{4}$  as much as the length; required its dimensions. *Ans. 16 by 12.*

11. There is a circle, whose diameter is 4 inches; what is the diameter of a circle, 9 times as large?

*Note.* The areas or contents of circles are in proportion to the squares of their diameters, or of their circumferences. Therefore, to find the diameter required, square the given diameter, multiply the square by the given ratio, and the square root of the product will be the diameter required.

$\sqrt{4 \times 4 \times 9}=12$  inches, *Ans.*

12. There are two circular ponds in a gentleman's pleas-


ure ground; the diameter of the less is 100 feet, and the greater is 3 times as large; what is its diameter?

*Ans.* 173'2+ ft.

13. If the diameter of a circle be 12 inches, what is the diameter of one  $\frac{1}{4}$  as large?

*Ans.* 6 inches.

¶ 103. 14. A carpenter has a large wooden square; one part of it is 4 feet long, and the other part 3 feet long; what is the length of a pole that will just reach from one end to the other?



*Note.* A figure of 3 sides is called a triangle, and if one of the corners be a square corner, or right angle, like the angle at B in the annexed figure, it is called a right-angled triangle, of which the square of the longest side A C, (called the hypotenuse, is equal to the sum

of the squares of the other two sides, A B and B C.

$4^2=16$ , and  $3^2=9$ ; then,  $\sqrt{9+16}=5$  feet, *Ans.*

15. If, from the corner of a square room, 6 feet be measured off one way, and 8 feet the other way, along the sides of the room, what will be the length of a pole reaching from point to point?

*Ans.* 10 feet.

16. A wall is 32 feet high, and a ditch before it is 24 feet wide; what is the length of a ladder that will reach from the top of the wall to the opposite side of the ditch?

*Ans.* 40 feet.

17. If the ladder be 40 feet, and the wall 32 feet, what is the width of the ditch?

*Ans.* 24 feet.

18. The ladder and ditch given, required the wall.

*Ans.* 32 feet.

19. The distance between the lower ends of two equal rafters is 32 feet, and the height of the ridge, above the beam on which they stand, is 12 feet; required the length of each rafter.

*Ans.* 20 feet.

20. There is a building 30 feet in length and 22 feet in width, and the eaves project beyond the wall 1 foot on every side; the roof terminates in a point at the centre of the

buildi  
is 10  
is the  
eaves  
middl  
one e  
*An*  
22'36  
21.  
what

22.  
my ro  
rods a

23.  
tance

F  
¶ 1  
of the  
length  
length  
consec  
power  
Her  
ber of  
of wh  
numbe  
1.  
each s  
2.  
on ea  
3.  
contai  
No  
4.  
feet!  
feet!

and the  
2+ ft.  
it is the  
inches.  
e; one  
t long;  
om one

re of 3  
triangle,  
corners  
or right  
gle at B  
figure, it  
t-angled  
ich the  
gest side  
hypot-  
the sum

et, Ans.  
be meas-  
the sides  
reaching  
. 10 feet.  
it is 24  
will reach  
he ditch?  
s. 40 feet.  
et, what  
s. 24 feet.  
all.  
s. 32 feet.  
two equal  
above the  
he length  
s. 20 feet.  
22 feet in  
ot on ex-  
tre of the

building, and is there supported by a post, the top of which is 10 feet above the beams on which the rafters rest; what is the distance from the foot of the post to the corners of the eaves? and what is the length of a rafter reaching to the middle of one side? — a rafter reaching to the middle of one end? and a rafter reaching to the corners of the eaves?

*Answers*, in order, 20ft.; 15'62+ft.; 18'86+ft.; and 22'36+ft.

21. There is a field 800 rods long and 600 rods wide; what is the distance between two opposite corners?

*Ans.* 1000 rods.

22. There is a square field containing 90 acres; how many rods in length is each side of the field? and how many rods apart are the opposite corners?

*Answers*, 120 rods; and 169'7+rods.

23. There is a square field containing 10 acres; what distance is the centre from each corner? *Ans.* 28'28+ rods.

## EXTRACTION OF THE CUBE ROOT.

¶ 104. A solid body, having six equal sides, and each of the sides an exact square, is a CUBE, and the measure in length of one of its sides is the root of that cube; for the length, breadth, and thickness of such a body are all alike; consequently, the length of one side, raised to the third power, gives the solid contents. See ¶ 33.

Hence it follows, that extracting the cube root of any number of feet is finding the length of one side of a cubic body, of which the whole contents will be equal to the given number of feet.

1. What are the solid contents of a cubic block, of which each side measures 2 feet? *Ans.*  $2^3=2 \times 2 \times 2=8$  feet.

2. How many solid feet in a cubic block, measuring 5 feet on each side? *Ans.*  $5^3=125$  feet.

3. How many feet in length is each side of a cubic block, containing 125 solid feet? *Ans.*  $\sqrt[3]{125}=5$  feet.

*Note.* The root may be found by trial.

4. What is the side of a cubic block containing 64 solid feet? — 27 solid feet? — 216 solid feet? — 512 solid feet? *Ans.* 4ft.; 3 ft.; 6 ft.; and 8 ft.

5. Supposing a man has 13824 feet of timber, in separate blocks of 1 cubic foot each; he wishes to pile them up in a cubic pile; what will be the length of each side of such a pile?


It is evident, the answer is found by extracting the cube root of 13824; but this number is so large, that we cannot so easily find the root by trial as in the former examples; We will endeavor, however, to do it by a *sort of trial*; and,

1st. We will try to ascertain the number of figures, of which the root will consist. This we may do by pointing the number off into periods of *three* figures each (T 101, ex. 1.)

OPERATION.

$$\begin{array}{r} 13824(2 \\ 8 \\ \hline 5824 \end{array}$$

Fig. I.



feet, which are now disposed of, and which, consequently, are to be deducted from the whole number of feet, 13824. 8000 taken from 13824 leave 5824 feet. This deduction is most readily performed by subtracting the cubic number, 8, or the cube of 2, (the figure of the root already found,) from the period 13, (thousands,) and bringing down the next period by the side of the remainder, making 5824, as before.

2d. The cubic pile A D is now to be enlarged by the addition of 5824 solid feet, and, in order to preserve the cubic form of the pile, the addition must be made on one half of its sides, that is, on 3 sides a, b, and c. Now, if the 5824

Pointing off, we see, the root will consist of 2 figures, a *ten* and a *unit*. Let us, then seek for the first figure, or tens of the root, which must be extracted from the left hand period, 13 (thousands.) The greatest cube in 13, (thousands) we find *by trial*, or by the *table of powers*, to be 8, (thousands,) the root of which is 2, (tens;) therefore, we place 2 (tens) in the root. The root, it will be recollected, is one side of a cube. Let us, then, form a cube, (Fig. I.) each side of which shall be supposed 20 feet, expressed by the root now obtained.

The contents of this cube are  $20 \times 20 \times 20 = 8000$  solid

solid  
sides,  
tient v  
the sid  
the le  
root, 2  
side, a  
400 X  
conver  
tiplied  
for the

OPER

Divis.

20



de, whic  
ens.] 3  
me in c




solid feet be divided by the square contents of these 3 equal sides, that is, by 3 times,  $[20 \times 20 = 400] = 1200$ , the quotient will be the thickness of the addition made to each of the sides  $a, b, c$ . But the root, 2, [tens,] already found, is the length of one of these sides; we therefore square the root, 2, [tens,]  $= 20 \times 20 = 400$ , for the square contents of one side, and multiply the product by 3, the number of sides,  $400 \times 3 = 1200$ , or, which is the same in effect, and more convenient in practice, we may square the 2, (tens,) and multiply the product by 300, thus,  $2 \times 2 = 4$ , and  $4 \times 300 = 1200$ , for the divisor, as before.

OPERATION—CONTINUED.

$$\begin{array}{r}
 13824 \text{ (24 Root.} \\
 8 \\
 \hline
 \text{Divis. } 1200) 5824 \text{ Dividend.} \\
 \underline{4800} \\
 960 \\
 \underline{64} \\
 5824 \\
 \underline{0000}
 \end{array}$$

FIG. II.



The divisor, 1200, is contained in the dividend 4 times; consequently, 4 feet is the thickness of the addition made to each of the three sides,  $a, b, c$ , and  $4 \times 1200 = 4800$ , is the solid feet contained in these additions; but if we look at Fig. II., we shall perceive, that this addition to the 3 sides does not complete the cube; for there are deficiencies in the 3 corners  $n, n, n$ . Now the length of each of these deficiencies is the same as the length of each side, that is, 2 [tens]  $= 20$ , and their width and thickness are each equal to the last quotient figure, [4]; their contents, therefore, or the number of feet required to fill these deficiencies, will be found by multiplying the square of the last quotient figure,  $[4^2] = 16$ , by the length of all the deficiencies, that is, by 3 times the length of each side, which is expressed by the former quotient figure, 2 [tens,] 3 times 2 [tens] are 6 [tens]  $= 60$ ; or what is the same in effect, and more convenient in practice, we may

, in sepa-  
e them up  
h side of  
  
g the cube  
we cannot  
examples;  
trial; and,  
figures, of  
y pointing  
01, ex. 1.)  
e see, the  
2 figures,  
et us, then  
t figure, or  
which must  
m the left  
thousands.)  
be in 13,  
d by trial,  
owers, to be  
the root of  
) therefore,  
in the root.  
e recollect-  
a cube. Let  
be, (Fig. I.)  
ch shall be  
expressed  
y obtained.  
his cube are  
8000 solid  
nsequently,  
feet, 13824.  
eduction is  
number, 8,  
ound,) from  
he next pe-  
as before.  
d by the ad-  
ve the cubic  
one half of  
if the 5824

multiply the quotient figure, 2, [tens,] by 30, thus,  $2 \times 30 = 60$ , as before; then,  $60 \times 16 = 960$ , contents of the three deficiencies,  $n, n, n$ .

FIG. III.





FIG. IV.



number; or it may be proved by adding together the contents of all the several parts, thus,

Feet.

8000=contents of Fig. I.

4800=addition to the sides  $a, b$ , and  $c$ , Fig. 1.

960=addition to fill the deficiencies  $n, n, n$ , Fig. II.

64=addition to fill the corner  $e, e, e$ , Fig. IV.

13824=contents of the whole pile, Fig. IV. 24 feet each side.

From the foregoing example and illustration we derive the following

Looking at Fig. III., we perceive there is still a deficiency in the corner where the last blocks meet. This deficiency is a cube, each side of which is equal to the last quotient figure, 4. The cube of 4, therefore,  $[4 \times 4 \times 4 = 64,]$  will be the solid contents of this corner, which, in Fig. IV. is seen filled.

Now, the sum of these several additions, viz.  $4800 + 960 + 64 = 5824$ , will make the subtrahend, which, subtracted from the dividend, leaves no remainder, and the work is done.

Fig. IV. shows the pile which 13824 solid blocks of one foot each would make, when laid together, and the root 24, shows the length of one side of the pile. The correctness of the work may be ascertained by cubing the side now found,  $24^3$ , thus,  $24 \times 24 \times 24 = 13824$ , the given

I.  
each  
figure

II.  
put it

III.  
and t  
this t

IV.  
it the

V.  
divid  
the d  
under

VI.  
forme  
and T  
cube  
trahen

VI.  
the r  
dend,  
whol

No  
the d  
perio

No  
the d  
root.

No  
we c  
ure,  
hend  
small

# RULE

*For extracting the Cube Root.*

I. Separate the given number into periods of three figures each by putting a point over the unit figure, and every third figure beyond the place of units.

II. Find the greatest cube in the left hand period, and put its root in the quotient.

III. Subtract the cube thus found from the said period, and to the remainder bring down the next period, and call this the *dividend*.

IV. Multiply the square of the quotient by 300, calling it the divisor.

V. Seek how many times the divisor may be had in the dividend, and place the result in the root ; then multiply the divisor by this quotient figure, and write the product under the dividend.

VI. Multiply the square of this quotient figure by the former figure or figures of the root, and this product by 30, and place the product under the last ; under all write the cube of this quotient figure, and call their amount the *subtrahend*.

VII. Subtract the subtrahend from the dividend, and to the remainder bring down the next period for a new dividend, with which proceed as before ; and so on, till the whole is finished.

*Note 1.* If it happens that the divisor is not contained in the dividend, a cipher must be put in the root, and the next period brought down for a dividend.

*Note 2.* The same rule must be observed for continuing the operation, and pointing off for decimals, as in the square root.

*Note 3.* The pupil will perceive that the number which we call the *divisor*, when multiplied by the last quotient figure, does not produce so large a number as the real subtrahend ; hence, the figure in the root must frequently be smaller than the quotient figure.

## EXAMPLES FOR PRACTICE.

6. What is the cube root of 1860867 ?

## OPERATION.

1860867 (123 Ans.

1

 $1^2 \times 300 = 300$  860 first Dividend.

$$\begin{array}{r} 600 \\ 2^2 \times 1 \times 30 = 120 \\ 2^3 = 8 \end{array}$$

728 first Subtrahend.

 $12^2 \times 300 = 43200$  132867 second Dividend.

$$\begin{array}{r} 129600 \\ 3^2 \times 12 \times 30 = 3240 \\ 3^3 = 27 \end{array}$$

132867 second Subtrahend.

000000

7. What is the cube root of 373248 ? Ans. 72.  
 8. What is the cube root of 21024576 ? Ans. 276.  
 9. What is the cube root of 84604519 ? Ans. 439.  
 10. What is the cube root of 000343 ? Ans. 07  
 11. What is the cube root of 2 ? Ans. 1.25+.  
 12. What is the cube root of  $\frac{8}{27}$  ? Ans.  $\frac{2}{3}$ .  
*Note.* See ¶ 99, ex. 10, and ¶ 102, ex. 14.  
 13. What is the cube root of  $\frac{125}{1000}$  ? Ans.  $\frac{5}{10}$ .  
 14. What is the cube root of  $\frac{343}{1000}$  ? Ans.  $\frac{7}{10}$ .  
 15. What is the cube root of  $\frac{1}{500}$  ? Ans.  $\frac{1}{125}$ .  
 16. What is the cube root of  $\frac{1}{125}$  ? Ans.  $\frac{1}{5}$ .

## SUPPLEMENT TO THE CUBE ROOT.

## QUESTIONS.

1. What is a cube? 2. What is understood by the cube root? 3. What is it to extract the cube root? 4. Why is the square of the quotient multiplied by 300 for a divisor? 5. Why, in finding the subtrahend, do we multiply the square of the last quotient figure by 30 times the former figure of the root? 6. Why do we cube the quotient figure? 7. How do we prove the operation?

EXERCISES.

1. What is the side of a cubical mound, equal to one 288 feet long, 216 feet broad, and 48 feet high ? *Ans.* 144 ft.

2. There is a cubic box, one side of which is 2 feet ; how many solid feet does it contain ? *Ans.* 8 feet.

3. How many cubic feet in one 8 times as large ? and what would be the length of one side ?

*Ans.* 64 solid feet, and one side is 4 feet.

4. There is a cubical box, one side of which is 5 feet ; what would be the side of one containing 27 times as much ? — 64 times as much ? — 125 times as much ?

*Ans.* 15, 20, and 25 feet.

5. There is a cubical box, measuring 1 foot on each side ; what is the side of a box 8 times as large ? — 27 times ? — 64 times ?

*Ans.* 2, 3, and 4 feet.

¶ 105. Hence we see, that the *sides* of cubes are as the *cube roots* of their *solid contents*, and, consequently, their *contents* are as the *cubes* of their *sides*. The same proportion is true of the *similar sides*, or of the *diameters* of all solid figures of similar forms.

6. If a ball, weighing 4 pounds, be 3 inches in diameter, what will be the diameter of a ball of the same metal, weighing 32 pounds ?  $4 : 32 :: 3^3 : 6^3$  *Ans.* 6 inches.

7. If a ball, 6 inches in diameter, weigh 32 pounds, what will be the weight of a ball 3 inches in diameter ? *Ans.* 4lbs.

8. If a globe of silver, 1 inch in diameter, be worth \$6, what is the value of a globe 1 foot in diameter ?

*Ans.* \$10368.

9. There are two globes ; one of them is 1 foot in diameter, and the other 40 feet in diameter ; how many of the smaller globes would it take to make 1 of the larger ?

*Ans.* 64000.

10. If the diameter of the sun is 112 times as much as the diameter of the earth, how many globes like the earth would it take to make one as large as the sun ? *Ans.* 1404928.

11. If the planet Saturn is 1000 times as large as the earth, and the earth is 7900 miles in diameter, what is the diameter of Saturn ? *Ans.* 79000 miles.

12. There are two planets of equal density ; the diameter of the less is to that of the larger as 2 to 9 ; what is the ratio of their solidities ? *Ans.*  $\frac{8}{729}$  ; or, as 8 to 729.

*Note.* The roots of most powers may be found by the square and cube root only. Thus, the biquadrate, or 4th root, is the square root of the square root; the 6th root is the cube root of the square root; the 8th root is the square root of the 4th root; the 9th root is the cube root of the cube root, &c. Those roots, viz. the 5th, 7th, 11th, &c., which are not resolvable by the square and cube roots, seldom occur, and, when they do, the work is most easily performed by logarithms; for, if the logarithm of any number be divided by the index of the root, the quotient will be the logarithm of the root itself.

### ARITHMETICAL PROGRESSION.

¶ 106. Any rank or series of numbers, more than two, increasing or decreasing by a constant difference, is called an *Arithmetical Series*, or *Progression*.

When the numbers are formed by a continual addition of the common difference, they form an *ascending series*; but when they are formed by a continual subtraction of the common difference, they form a *descending series*.

Thus, { 3, 5, 7, 9, 11, 13, 15, &c. is an *ascending series*.  
 { 15, 13, 11, 9, 7, 5, 3, &c. is a *descending series*.

The numbers which form the series are called the *terms* of the series. The *first* and *last* terms are the *extremes*, and the other terms are called the *means*.

There are five things in arithmetical progression, any three of which being given, the other two may be found:—

- 1st. The *first* term.
- 2d. The *last* term.
- 3d. The *number* of terms.
- 4th. The *common difference*.
- 5th. The *sum* of all the terms.

1. A man bought 100 yards of cloth, giving 4 pence for the first yard, 7 pence for the second, 10 pence for the third, and so on, with a common difference of 3 pence; what was the cost of the last yard?

As the *common difference*, 3, is added to every yard except the last, it is plain the *last* yard must be  $99 \times 3 = 297$  pence, more than the *first* yard. *Ans.* 301 pence.

by the  
th root,  
t is the  
are root  
the cube  
, which  
dom oc-  
performed  
er be di-  
the log-

Hence, when the first term, the common difference, and the number of terms, are given, to find the last term,—Multiply the number of terms, less 1, by the common difference, and add the first term to the product for the last term.

2. If the first term be 4, the common difference 3, and the number of terms 100, what is the last term? *Ans.* 301.

3. There are, in a certain triangular field, 41 rows of corn; the first row, in 1 corner, is a single hill, the second contains 3 hills, and so on, with a common difference of 2; what is the number of hills in the last row? *Ans.* 81 hills.

4. A man puts out 1£, at 6 per cent simple interest, which, in 1 year, amounts to  $1\frac{3}{50}$ £, in 2 years to  $1\frac{6}{25}$ £, and so on, in arithmetical progression, with a common difference of  $\frac{3}{50}$ £; what would be the amount in 40 years?

*Ans.*  $3\frac{28}{50}$ £.

han two.  
is called  
ddition of  
ries; but  
the com-

Hence we see, that the yearly amounts of any sum, at simple interest, form an arithmetical series, of which the principal is the first term, the last amount is the last term, the yearly interest is the common difference, and the number of years is 1 less than the number of terms.

ng series-  
ng series.  
the terms  
emes, and

5. A man bought 100 yards of cloth in arithmetical progression; for the first yard he gave 4 pence, and for the last 301 pence; what was the common increase of the price on each succeeding yard?

any three  
:—

This question is the reverse of example 1; therefore,  $301 - 4 = 297$ , and  $297 \div 99 = 3$ , common difference.

Hence, when the extremes and number of terms are given, to find the common difference,—Divide the difference of the extremes by the number of terms, less 1, and the quotient will be the common difference.

6. If the extremes be 5 and 605, and the number of terms 151, what is the common difference? *Ans.* 4.

7. If a man puts out 1£, at simple interest, for 40 years, and receives, at the end of the time,  $3\frac{2}{3}$ £, what is the rate?

If the extremes be 1 and  $3\frac{2}{3}$ , and the number of terms 41, what is the common difference? *Ans.*  $\frac{2}{30}$ .

pence for  
the third,  
what was

8. A man had 8 sons, whose ages differed alike; the youngest was 10 years old, and the eldest 45; what was the common difference of their ages? *Ans.* 5 years.

rd except  
3, = 297  
01 pence.

9. A man bought 100 yards of cloth in arithmetical se-



ries, he gave 4 pence for the *first* yard, and 301 pence for the *last* yard; what was the average price per yard, and what was the amount of the whole?

Since the price of each succeeding yard increases by a *constant excess*, it is plain, the *average* price is as much *less* than the price of the *last* yard, as it is *greater* than the price of the *first* yard; therefore, one half the sum of the first and last price is the *average price*.

One half of  $4d. + 301d. = 152\frac{1}{2}d.$  = average }  
price; and the price,  $152\frac{1}{2}d. \times 100 = 15250d. =$  } *Ans.*  
 $63\text{£ } 10s. 10d.,$  whole cost.

Hence, when *the extremes and the number of terms* are given, to find the sum of all the terms,—Multiply  $\frac{1}{2}$  the sum of the extremes by the number of terms, and the product will be the answer.

10. If the extremes be 5 and 605, and the number of terms be 151, what is the sum of the series? *Ans.* 46055.

11. What is the sum of the first 100 numbers, in their natural order, that is, 1, 2, 3, 4, &c. ? *Ans.* 5050.

12. How many times does a common clock strike in 12 hours? *Ans.* 78.

13. A man rents a house for 50£, annually, to be paid at the close of each year; what will the rent amount to in 20 years allowing 6 per cent, simple interest, for the use of the money?

The last year's rent will evidently be 50£ without interest, the last but *one* will be the amount of 50£ for 1 year, the last but *two* the amount of 50£ for 2 years, and so on, in arithmetical series, to the first, which will be the amount of 50£ for 19 years = 107£.

If the first term be 50, the last term 107, and the number of terms 20, what is the sum of the series? *Ans.* 1570£.

14. What is the amount of an annual pension of 100£, being in arrears, that is, remaining unpaid, for 40 years, allowing 5 per cent *simple interest*? *Ans.* 7900£.

15. There are, in a certain triangular field, 41 rows of corn; the first row, being in 1 corner, is a single hill, and the last row, on the side opposite, contains 81 hills; how many hills of corn in the field? *Ans.* 1681 hills.

16. If a triangular piece of land, 30 rods in length, be 20 rods wide at one end, and come to a point at the other, what

ence for  
rd, and  
es by a  
uch less  
han the  
n of the

} Ans.

re given,  
m of the  
t will be

number of  
s. 46055.  
in their  
ns. 5050.

ike in 12  
Ans. 78.  
o be paid  
unt to in  
the use of

thout in-  
r 1 year,  
so on, in  
e amount

e number  
s. 1570£.  
of 100£,  
40 years,  
s. 7900£.  
rows of  
hill, and  
ills; how  
81 hills.  
th, be 20  
er, what

number of square rods does it contain? Ans. 300.

17. A debt is to be discharged at 11 several payments, in arithmetical series, the first to be 5£, and the last 75£; what is the whole debt? — the common difference between the several payments?

Ans. whole debt, 440£; common difference, 7£.

18. What is the sum of the series 1, 3, 5, 7, 9, &c. to 1001? Ans. 251001.

Note. By the reverse of the rule under ex. 5, the difference of the extremes 1000, divided by the common difference 2, gives a quotient, which, increased by 1, is the number of terms = 501.

19. What is the sum of the arithmetical series 2, 2½, 3, 3½, 4, 4½, &c., to the 50th term inclusive? Ans. 712½.

20. What is the sum of the decreasing series 30, 29½, 29¼, 29⅛, &c. down to 0?

Note.  $30 \div \frac{1}{8} + 1 = 91$ , number of terms. Ans. 1365.

### QUESTIONS.

1. What is an arithmetical progression? 2. When is the series called ascending? 3. — when descending? 4. What are the numbers, forming the progression, called? 5. What are the first and last terms called? 6. What are the other terms called? 7. When the first term, common difference, and number of terms, are given, how do you find the last term? 8. How may arithmetical progression be applied to simple interest? 9. When the extremes and number of terms are given, how do you find the common difference? 10. — how do you find the sum of all the terms?

### GEOMETRICAL PROGRESSION.

¶ 107. Any series of numbers, continually increasing by a constant multiplier, or decreasing by a constant divisor, is called a *Geometrical Progression*. Thus, 1, 2, 4, 8, 16, &c. is an increasing geometrical series, and 8, 4, 2, 1, ½, ¼, &c. is a decreasing geometrical series.

As in arithmetical, so also in geometrical progression, there are five things, any three of which being given, the other two may be found:—

1st. The first term.

2d. The last term.

3d. The *number of terms*.

4th. The *ratio*.

5th. The *sum of all the terms*.

The *ratio* is the *multiplier*, or *divisor*, by which the series is formed.

1. A man bought a piece of silk, measuring 17 yards, and, by agreement, was to give what the last yard would come to, reckoning 3 pence for the first yard, 6 pence for the second, and so on doubling the price to the last; what did the piece of silk cost him?

$3 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 = 196608$  pence, =  $819\text{ } \text{£} \text{ } 4\text{s.}$  answer.

In examining the process by which the last term (196608) has been obtained, we see, that it is a product, of which the *ratio* (2) is sixteen times a factor, that is, *one time less* than the number of terms. The last term, then, is the *sixteenth power* of the *ratio*, (2,) multiplied by the *first term* (3.)

Now, to raise 2 to the 16th power, we need not produce all the *intermediate powers*; for  $2^4 = 2 \times 2 \times 2 \times 2 = 16$ , is a product of which the *ratio* 2 is 4 times a factor; now, if 16 be multiplied by 16, the product, 256, evidently contains the same factor (2) 4 times + 4 times, = 8 times; and  $256 \times 256 = 65536$ , a product of which the *ratio* (2) is 8 times + 8 times, = 16 times, factor; it is, therefore, the 16th power of 2, and, multiplied by 3, the first term, gives 196608, the last term, as before. Hence,

When the *first term*, *ratio*, and *number of terms*, are given, to find the last term,—

I. Write down a few leading powers of the *ratio* with their *indices* over them.

H. Add together the most *convenient indices*, to make an index *less by one* than the number of the term sought.

III. Multiply together the *powers* belonging to those *indices*, and their product, multiplied by the *first term*, will be the term sought.

2. If the first term be 5, and the ratio 3, what is the 8th term?

Powers of the ratio, with their indices over them.  $\begin{cases} 1 & 2 & 3 & +4 = 7 \\ 3, & 9, & 27, & \times 81 = 2187 \times 5 \\ & & & \text{first term,} = 10935, \text{ Ans.} \end{cases}$

3. A man plants 4 kernels of corn, which, at harvest, produce 32 kernels; these he plants the second year; now, supposing the annual increase to continue 8 fold, what would be the produce of the 16th year, allowing 1000 kernels to a pint?

Ans. 2199023255552 bushels.

4. Supposing a man had put out one penny at compound interest in 1620, what would have been the amount in 1824, allowing it to double once in 12 years?

$2^{17} = 131072$ . Ans. 546 £ 2s. 8d.

5. A man bought 4 yards of cloth, giving 2 pence for the first yard, 6 pence for the second, and so on, in 3 fold ratio: what did the whole cost him?

$2+6+18+54 = 80$  pence. Ans. 80 pence.

In a long series, the process of adding in this manner would be tedious. Let us try, therefore, to devise some shorter method of coming to the same result. If all the terms, *excepting the last*, viz.  $2+6+18$ , be multiplied by the ratio, 3, the product will be the series  $6+18+54$  subtracting the *former* series from the *latter*, we have, for the remainder,  $54-2$ , that is, the *last term*, less the *first term*, which is evidently as many times the first series ( $2+6+18$ ) as is expressed by the ratio, less 1: hence, if we divide the difference of the extremes ( $54-2$ ) by the ratio, less 1, ( $3-1$ ), the quotient will be the sum of all the terms, *excepting the last*, and, adding the last term, we shall have the *whole amount*. Thus,  $54-2=52$ , and  $3-1=2$ ; then,  $52 \div 2 = 26$ , and 54 added, makes 80, Ans. as before.

Hence, when the extremes and ratio are given, to find the sum of the series,—Divide the *difference* of the extremes by the ratio less 1, and the quotient, increased by the greater term, will be the answer.

6. If the extremes be 4 and 131072, and the ratio 8, what is the whole amount of the series?

$$\frac{131072-4}{8-1} + 131072 = 149796 \text{ Answer}$$

7. What is the sum of the descending series 3, 1,  $\frac{1}{3}$ ,  $\frac{1}{9}$ , &c., extended to infinity?

It is evident the last term must become 0, or indefinitely near to nothing; therefore, the extremes are 3 and 0, the ratio 3.

Ans.



be reduced to an improper fraction  $= \frac{3}{2}$ , or to a decimal, 1.5.

$$\frac{\frac{3}{2}^5 - 1}{\frac{3}{2} - 1} \times 1000 = 13187\frac{1}{2}; \text{ or, } \frac{1.5^5 - 1}{1.5 - 1} \times 1000 = 13187.50\text{£} = 13187\text{£ } 10\text{s. Answer.}$$

*Compound Interest by Progression.*

¶ 108. 1. What is the amount of 4£, for 5 years at 6 per cent compound interest?

We have seen, (¶ 86,) that *compound interest* is that, which arises from adding the interest to the principal at the close of each year, and, for the next year, casting the interest on that *amount* and so on. The amount of 1£ for 1 year is 1.06; if the *principal*, therefore, be multiplied by 1.06, the product will be its *amount* for 1 year; this *amount*, multiplied by 1.06, will give the amount (compound interest) for 2 years; and this *second amount*, multiplied by 1.06, will give the amount for 3 years; and so on. Hence, the several *amounts*, arising from any sum at compound interest, form a *geometrical series*, of which the *principal* is the *first term*; the *amount* of 1£ or \$1, &c. at the *given rate* per cent, is the *ratio*; the *time*, in years, is 1 less than the number of terms; and the last amount is the last term.

The last question may be resolved into this: If the first term be 4, the number of terms 6, and the ratio 1.06 what is the last term?

$$1.06^5 = 1.338, \text{ and } 1.338 \times 4 = \text{£}5.352+.$$

Ans. 5£ 7s. 0½d.

*Note 1.* The powers of the amounts of 1£, at 5 and at 6 per cent, may be taken from the table, under ¶ 85. Thus, opposite 5 years, under 6 per cent, you find 1.338, &c.

*Note 2.* The several processes may be conveniently exhibited by the use of letters thus:

Let P represent the Principal.

" R " the Ratio, or the amount of 1£, &c. for 1 yr.

" T " the Time in years.

" A " the Amount.

When two or more letters are *joined together*, like a word, they are to be *multiplied together*. Thus PR. implies, that the *principal* is to be multiplied by the *ratio*. When one letter is placed *above* another, like the index of a power, the

*first* is to be raised to a power, whose index is denoted by the second. Thus  $R^T$  implies, that the ratio is to be raised to a power, whose index shall be equal to the time, that is, the number of years.

2. What is the amount of 40£ for 11 years, at 5 per cent, compound interest?

$$R^T \times P = A; \text{ therefore, } 1.05^{11} \times 40 = 68.4.$$

Ans. 68£ 8s.

3. What is the amount of 6£ for 4 years, at 10 per cent compound interest?

Ans. 8£ 15s. 8d.

4. If the amount of a certain sum for 5 years, at 6 per cent, compound interest, be 5£ 7s. 0½d. what is that sum, or principal?

If the number of terms be 6, the ratio 1.06, and the last term 5.352, what is the first term?

This question is the reverse of the last; therefore,

$$\frac{A}{R^T} = P; \text{ or, } \frac{5.352}{1.338} = 4.$$

Ans. 4£.

5. What principal, at 10 per cent. compound interest, will amount, in 4 years, to 8.7846£?

Ans. 6£.

6. What is the present worth of 68£ 8s. due 11 years hence, discounting at the rate of 5 per cent compound interest?

Ans. 40£.

7. At what rate per cent, will, 6£ amount to 8.7846£ in 4 years?

If the first term be 6, the last term 8.7846, and the number of terms 5, what is the ratio?

$\frac{A}{P} = R^T$ , that is,  $\frac{8.7846}{6} = 1.4641 = \text{the 4th power of the ratio}$ ; and then, by extracting the 4th root, we obtain 1.10 for the ratio.

Ans. 10 per cent.

8. In what time will 6£ amount to 8.7846£, at 10 per cent. compound interest?

$\frac{A}{P} = R^T$ , that is,  $\frac{8.7846}{6} = 1.4641 = 1.10^T$ ; therefore, if we divide 1.4641 by 1.10, and then divide the quotient thence arising by 1.10, and so on, till we obtain a quotient that will not contain 1.10 the number of these divisions will be the number of years.

Ans. 4 years.

9. At 5 per cent compound interest, in what time will 40£ amount to 68£ 8s.?

Ha  
whic  
unde  
find

10  
4£ a

¶ 11  
show  
the a  
An  
year  
the lif  
Wh  
becom

The  
paid,  
have

1. V  
which  
compo

The  
the lar  
last bu  
years,  
will be  
that is,  
the ter

If th  
ratio 1  
Cons

Henc  
are giv  
of 1£,  
of year  
mainder  
the ann

Note.



noted by the  
be raised to  
ne, that is,

, at 5 per

ns. 68£ 8s.

10 per cent

3£ 15s. 8d.

s, at 6 per

that sum,

and the last

fore,

Ans. 4£.

nd interest,

Ans. 6£.

ne 11 years

ound in-

Ans. 40£.

o 87846£

nd the num-

wer of the

obtain 1'10

0 per cent.

, at 10 per

herefore, if

uent thence

otient that

ons will be

ns. 4 years.

t time will

Having found the power of the ratio 1'05, as before, which is 1'71, you may look for this number in the *table*, under the given rate, 5 per cent, and against it you will find the number of years. *Ans.* 11 years.

10. At 6 per cent compound interest, in what time will 4£ amount to 5£ 7s. 0½d. *Ans.* 5 years.

### *Annuities at Compound Interest.*

¶ 109. It may not be amiss, in this place, briefly to show the application of compound interest, in computing the amount and present worth of *annuities*.

An **ANNUITY** is a sum payable at *regular periods*, of one year each, either for a *certain number of years*, or during the *life of the pensioner*, or *forever*.

When annuities, rents, &c. are not paid at the time they become due, they are said to be in *arrears*.

The sum of all the annuities, rents, &c. remaining unpaid, together with the *interest* on each, for the time they have remained due, is called the *amount*.

1. What is the amount of an annual pension of 100£, which has remained unpaid 4 years, allowing 6 per cent compound interest?

The last year's pension will be 100£, without interest; the last but one will be the amount of 100£ for 1 year; the last but two the amount (compound interest) of 100£ for 2 years, and so on; and the sum of these several amounts will be the answer. We have then, a series of amounts, that is, a geometrical series, (¶ 108,) to find the sum of all the terms.

If the first term be 100, the number of terms 4, and the ratio 1'06, what is the sum of all the terms?

Consult the rule, under ¶ 107, ex. 11.

$$\frac{1'06^4 - 1}{'06} \times 100 = 437'45. \quad \text{Ans. } 437£ \ 9s.$$

Hence, when the annuity, the time, and rate per cent, are given, to find the amount,—**RAISE** the ratio (the amount of 1£, &c. for 1 year) to a power denoted by the number of years; from this power subtract 1, then divide the remainder by the ratio less 1, and the quotient, multiplied by the annuity, will be the amount.

*Note.* The powers of the amounts, at 5 and 6 per cent

up to the 24th, may be taken from the table, under ¶ 85.

2. What is the amount of an annuity of 50£, it being in arrears 20 years, allowing 5 per cent compound interest ?

Ans. 1653£ 5s. 9½d.

3. If the annual rent of a house, which is 150£, be in arrears 4 years, what is the amount, allowing 10 per cent compound interest ?

Ans. 696£ 3s.

4. To how much would a salary of 500£ per annum amount in 14 years, the money being improved at 6 per cent compound interest ? — in 10 years ? — in 20 years ? — in 22 years ? — in 24 years ?

Ans. to the last, 25407£ 15s.

¶ 110. If the annuity is paid in advance, or if it be bought at the beginning of the first year, the sum which ought to be given for it is called the *present worth*.

5. What is the present worth of an annual pension of 100£, to continue for 4 years, allowing 6 per cent compound interest ?

The present worth is, evidently, a sum, which, at 6 per cent, compound interest, would, in 4 years, produce an amount, equal to the amount of the annuity in arrears the same time.

By the last rule we find the amount=437'45£, and by the directions under ¶ 108, ex. 4, we find the present worth=346'51£.

Ans. 346£ 10s. 4½d.

Hence, to find the present worth of any annuity,—First find its amount in arrears for the whole time ; this amount, divided by that power of the ratio denoted by the number of years, will give the present worth.

6. What is the present worth of an annual salary of 100£ to continue 20 years, allowing 5 per cent ?

Ans. 1246£ 4s. 4½d.

The operations under this rule being somewhat tedious, we subjoin a

#### TABLE,

Showing the present worth of 1£ or \$1 annuity, at 5 and 6 per cent compound interest, for any number of years, from 1 to 34.

¶ 85.  
it being  
interest ?  
5s. 9 $\frac{1}{2}$ d.  
be in ar-  
per cent  
696£ 3s.  
er annum  
at 6 per  
— in 20

407£ 15s.  
r if it be  
um which  
ension of  
compound

h, at 6 per  
duce an a-  
rrears the

and by the  
t worth=  
2 10s. 4 $\frac{1}{2}$ d.  
ity,—First  
is amount,  
ne number

ry of 100£

£ 4s. 4 $\frac{1}{2}$ d.  
at tedious,

at 5 and  
of years,

Years.	5 per cent.	6 per cent.	Years.	5 per cent.	6 per cent.
1	0·95238	0·94339	18	11·68958	10·8276
2	1·85941	1·83339	19	12·08532	11·15811
3	2·72325	2·67301	20	12·46221	11·46902
4	3·54595	3·4651	21	12·82115	11·76407
5	4·32948	4·21236	22	13·163	12·04158
6	5·07569	4·91732	23	13·48807	12·30338
7	5·78637	5·58238	24	13·79864	12·55035
8	6·46321	6·20979	25	14·09394	12·78335
9	7·10782	6·80169	26	14·37518	13·00316
10	7·72173	7·36008	27	14·64303	13·21053
11	8·30641	7·88687	28	14·89813	13·40616
12	8·86325	8·38384	29	15·14107	13·59072
13	9·39357	8·85268	30	15·37245	13·76483
14	9·89864	9·29498	31	15·59281	13·92908
15	10·37966	9·71225	32	15·80268	14·08398
16	10·83777	10·10589	33	16·00255	14·22917
17	11·27407	10·47726	34	16·1929	14·36613

It is evident that the present worth of 2£ annuity is 2 times as much as that of 1£ ; the present worth of 3£ will be 3 times as much, &c. Hence, to find the present worth of any annuity, at 5 or 6 per cent,—Find, in this table, the present worth of 1£ annuity, and multiply it by the given annuity, and the product will be the present worth.

7. What ready money will purchase an annuity of 150£ to continue 30 years, at 5 per cent, compound interest ?

The present worth of 1£ annuity, by the table, for 30 years, is 15·37245 ; therefore,  $15·37245 \times 150 = 2305·867£ = 2305£ 17s. 4d.$  Answer.

8. What is the present worth of a yearly pension of 40£ to continue 10 years, at 6 per cent compound interest ?—at 5 per cent ? —to continue 15 years ? — 20 years ? — 25 years ? 34 years ? Ans. to the last, 647£ 14s. 3 $\frac{1}{2}$ d.

When annuities do not commence till a certain period of time has elapsed, or till some particular event has taken place, they are said to be in REVERSION.

9. What is the present worth of 100£ annuity, to be continued 4 years, but not to commence till 2 years hence, allowing 6 per cent compound interest ?

The present worth is evidently a sum which, at 6 per cent compound interest, would in 2 years produce an amount

equal to the present worth of the annuity, were it to commence immediately. By the last rule, we find the present worth of the annuity, to commence immediately, to be 346'51£, and, by directions under ¶ 108, ex. 4, we find the present worth of 346'51£ for 2 years to be 308'393£.

Ans. 308£ 7s. 10½d.

Hence, to find the present worth of any annuity taken in reversion, at compound interest,—First, find the present worth, to commence immediately, and this sum, divided by the power of the ratio, denoted by the time in reversion, will give the answer.

10. What ready money will purchase the reversion of a lease of 60£ per annum, to continue 6 years, but not to commence till the end of 3 years, allowing 6 per cent compound interest to the purchaser?

The present worth to commence immediately, we find to be, 295'039, and  $\frac{295'039}{1'06^3} = 247'72$ . Ans. 247£ 14s. 4½d.

It is plain, the same result will be obtained by finding the present worth of the annuity, to commence immediately, and to continue to the end of the time, that is  $3+6=9$  years, and then subtracting from this sum the present worth of the annuity, continuing for the time of the reversion, 3 years. Or, we may find the present worth of 1£ for the two times by the table, and multiply their difference by the given annuity. Thus, by the table,

The whole time, 9 years, = 6'80169

The time in reversion, 3 years, = 2'67301

Difference, = 4'12868  
60

247'72080£

247'72080£ = 247£ 14s. 4½d. Ans.

11. What is the present worth of a lease of 100£ to continue 20 years, but not to commence till the end of 4 years, allowing 5 per cent? — what, if it be 6 years in reversion? — 8 years? — 10 years? — 14 years?

¶ 111. 12. What is the worth of a freehold estate, of which, the yearly rent is 60£, allowing to the purchaser 6 per cent?

In this case, the annuity continues *forever*, and the estate

is evidently worth a sum, of which the yearly interest is equal to the yearly rent of the estate. The principal multiplied by the rate gives the interest; therefore, the interest divided by the rate will give the principal;  $60 \div .06 = 1000$ .

Ans. 1000£.

Hence, to find the present worth of an annuity, continuing forever,—Divide the annuity by the *rate per cent*, and the quotient will be the present worth.

*Note.* The worth will be the same, whether we reckon simple or compound interest; for since a *year's interest of the price is the annuity*, the profits arising from *that price* can neither be *more nor less* than the profits arising from the *annuity*, whether they be employed at simple or compound interest.

13. What is the worth of 100£ annuity, to continue forever, allowing to the purchaser 4 per cent? —allowing 5 per cent? —8 per cent? —10 per cent? —15 per cent? —20 per cent? *Ans. to last, 500£.*

14. Suppose a freehold estate of 60£ *per annum*, to commence 2 years hence, be put on sale; what is its value, allowing the purchaser 6 per cent?

Its present worth is a sum which, at 6 per cent, compound interest, would, in 2 years, produce an amount equal to the worth of the estate *if entered on immediately*.

$$\frac{60}{.06} = 1000£ = \text{the worth, if entered on immediately,}$$

$$\text{and } \frac{1000£}{1.06^2} = 889.996£ = 889£ 19s. 11d. \text{ the present worth.}$$

The same result may be obtained by subtracting from the worth of the estate, to commence immediately, the present worth of the annuity 60, for 2 years, the time of *reversion*. Thus, by the table, the present worth of 1£ for 2 years is  $1.83339 \times 60 = 110.0034 =$  present worth of 60£ for 2 years, and  $1000£ - 110.0034 = 889.9966£ = 889£ 19s. 11d.$  *Ans. as before.*

15. What is the present worth of a perpetual annuity of 100£, to commence 6 years hence, allowing the purchaser 5 per cent compound interest? — what, if 8 years in reversion? — 10 years? — 4 years? — 30 years?

*Ans. to the last, 462£ 15s. 1½d.*

The foregoing examples, in compound interest, have been

confined to *yearly* payments ; if the payments are *half yearly*, we take half the principal or annuity, half the rate per cent, and twice the number of years, and work as before, and so for any other part of a year.

### QUESTIONS.

1. What is a geometrical progression or series? 2. What is the ratio? 3. When the first term, the ratio and the number of terms, are given, how do you find the *last term*? 4. When the extremes and ratio are given, how do you find the *sum of all the terms*? 5. When the first term, the ratio, and the number of terms, are given, how do you find the amount of the series? 6. When the ratio is a fraction, how do you proceed? 7. What is *compound interest*? 8. How does it appear that the *amounts*, arising by compound interest, form a *geometrical series*? 9. What is the *ratio* in compound interest? — the number of *terms*? — the *first term*? the *last term*? 10. When the rate, the time, and the principal, are given, how do you find the *amount*? — 11. When A. R. and T. are given, how do you find P.? 12. When A. P. and T. are given, how do you find R.? 13. When A. P. and R. are given, how do you find T.? 14. What is an *annuity*? 15. When are annuities said to be *in arrears*? 16. What is the *amount*? 17. In a geometrical series, to what is the *amount* of an annuity, equivalent? 18. How do you find the *amount* of an annuity, at compound interest? 19. What is the *present worth* of an annuity? — how computed at compound interest? — how found by the table? 20. What is understood by the term *reversion*? 21. How do you find the present worth of an annuity, taken in *reversion*? — by the table? 22. How do you find the present worth of a *freehold estate*, or a *perpetual annuity*? — the same taken in *reversion*? — by the table?

## PERMUTATION.

¶ 112. Permutation is the method of finding how many different ways the order of any number of things may be varied or changed.

1. Four gentlemen agreed to dine together so long as they could sit, every day, in a different order, or position ; how many days did they dine together ?

Had there been but two of them, *a* and *b*, they could sit only in 2 times 1 ( $1 \times 2 = 2$ ) different positions, thus, *a b*, and *b a*. Had there been *three*, *a b* and *c*, they could sit in  $1 \times 2 \times 3 = 6$  different positions ; for, beginning the order with *a*, there will be 2 positions, viz. *a b c*, and *a c b* ; next beginning with *b*, there will be 2 positions, *b a c*, and *b c a* ; last-

are half year-  
half the rate per  
work as before,

2. What is the  
number of terms,  
when the extremes  
all the terms? 5.  
terms, are given,  
when the ratio is a  
compound interest? 8.  
compound interest.  
the last term?  
given, how do you  
are given, how do  
do you find R.? 14. What is  
in arrears? 16.  
to what is the a-  
find the amount  
the present worth  
interest?  
by the term rever-  
an annuity, taken  
find the present  
— the same

ding how many  
things may be  
ther so long as  
er, or position;

ey could sit only  
thus,  $a b$ , and  $b$   
uld sit in  $1 \times 2$   
e order with  $a$ ,  
 $b$ ; next begin-  
and  $b c a$ ; last-

ly, beginning with  $c$ , we have  $c a b$ , and  $c b a$ , that is, in  
all,  $1 \times 2 \times 3 = 6$  different positions. In the same manner  
if there be four, the different positions will be  $1 \times 2 \times 3 \times$   
 $4 = 24$ . Ans. 24.

Hence, to find the number of different changes or permu-  
tations, of which any number of different things are capa-  
ble,—Multiply continually together, all the terms of the  
natural series of numbers, from 1 up to the given number,  
and the last product will be the answer.

2. How many variations may there be in the position of  
the nine digits? Ans. 362880.

3. A man bought 25 cows, agreeing to pay for them 1  
penny for every different order in which they could all be  
placed; how much did the cows cost him?

Ans. 646300418472124416000000

4. A certain Church has 8 bells; how many changes  
may be rung upon them. Ans. 40320.

### MISCELLANEOUS EXAMPLES.

113. 1.  $4 + 6 \times 7 - 1 = 60$

A line, or *vinculum*, drawn over several numbers, signi-  
fies, that the numbers under it, are to be taken jointly, or  
as one whole number.

2.  $9 - 8 + 4 \times 8 + 4 - 6 =$  how many? Ans. 30.

3.  $7 + 4 - 2 + 3 + 40 \times 5 =$  how many? Ans. 230.

4.  $3 + 6 - 2 \times 4 - 2 =$  how many?  
 $2 \times 2$  Ans.  $3\frac{1}{2}$ .

5. There are 2 numbers; the greater is 25 times 78, and  
their difference is 9 times 15; their sum and product are  
required. Ans. 3765 is their sum, 3539250 their product.

6. What is the difference between thrice five and thirty,  
and thrice thirty-five?  $35 + 3 - 5 \times 3 + 30 = 60$ , Ans.

7. What is the difference between six dozen dozen, and  
half a dozen dozen? Ans. 792.

8. What number divided by 7 will make 6488?

9. What number multiplied by 6 will make 2058?



10. A gentlemen went to sea at 17 years of age ; 8 years after he had a son born, who died at the age of 35 ; after whom the father lived twice 20 years ; how old was the father at his death ?

Ans. 100 years.

11. What number is that, which being multiplied by 15 the product will be  $\frac{3}{4}$  ?

$\frac{3}{4} \div 15 = \frac{1}{20}$ , Ans.

12. What decimal is that, which being multiplied by 15, the product will be .75 ?

$.75 \div 15 = .05$  Ans.

13. What is the decimal equivalent to  $\frac{2}{5}$  ?

Ans. .0285714.

14. What fraction is that, to which if you add  $\frac{3}{5}$ , the sum will be  $\frac{8}{9}$  ?

Ans.  $\frac{1}{9}$ .

15. What number is that, from which if you take  $\frac{3}{8}$ , the remainder will be  $\frac{1}{4}$  ?

Ans.  $\frac{3}{8}$ .

16. What number is that, which being divided by  $\frac{3}{4}$ , the quotient will be 21 ?

Ans. 15 $\frac{3}{4}$ .

17. What number is that, from which if you take  $\frac{3}{4}$  of itself, the remainder will be 12 ?

Ans. 20.

19. What number is that, to which if you add  $\frac{3}{4}$  of  $\frac{3}{4}$  of itself, the whole will be 20 ?

Ans. 12.

20. What number is that, of which 9 is the  $\frac{3}{4}$  part ?

Ans. 13 $\frac{1}{2}$ .

21. A farmer carried a load of produce to market ; he sold 780 lbs. of pork, at 3d. per pound ; 250 lbs. of cheese, at 5d. per lb. ; 154 lbs. of butter, at 10d. per lb. : in pay he received 60 lbs. of sugar, at 7d. per lb. ; 15 gallons of molasses, at 2s. 3d. per gallon ;  $\frac{1}{2}$  barrel of mackerel, at 18s. 9d. ; 4 bushels of salt, at 6s. 4d. per bushel ; and the balance in money ; how much money did he receive ?

Ans. 15£ 14s. 8d.

22. A farmer carried his grain to market, and sold 75 bushels of wheat at 7s. 3d. per bushel ; 64 bushels of rye, at 4s. 9d. per bushel ; 142 bushels of corn, at 2s. 6d. per bushel : in exchange he received sundry articles :—3 pieces cloth, each containing 31 yards, at 8s. 9d. per yard ; 2 quintals of fish, 11s. 6d. per quintal ; 8 hhds. of salt, 1£ 1s. 6d. per hhd. and the balance in money : how much money did he receive ?

Ans. 9£ 14s.

23. A man exchanges 760 gallons of molasses, at 2s. per gallon, for 66 $\frac{1}{2}$  cwt. of cheese, at 1£ per cwt. ; how much will be the balance in his favor ?

Ans. 9£ 10s.

24. Bought 84 yards of cloth, at 6s. 3d. per yard ; how

; 8 years  
35; after  
as the fa-  
00 years.  
ied by 15  
=70, Ans.  
ied by 15,  
=05 Ans.

0285714.  
3, the sum  
Ans. 13.  
ake 3, the  
Ans. 38.  
by 3, the  
Ans. 153.  
take 3 of  
Ans. 20.  
1 2 of 3 of  
Ans. 12.  
part?

Ans. 131.  
arket; he  
of cheese,  
b.: in pay  
gallons of  
ackerel, at  
; and the  
ive?

£ 14s. 8d.  
nd sold 75  
els of rye,  
s. 6d. per  
—3 pieces  
r yard; 2  
alt, 1£ 1s.  
ch money  
s. 9£ 14s.  
es, at 2s.  
wt.; how  
s. 9£ 10s.  
ard; how

much did it come to? How many bushels of wheat, at 7s. 6d. per bushel, will it take to pay for it?

Ans. to the last, 70 bushels.

25. A man sold 342 pounds of beef, at 4d. per pound, and received his pay in molasses, at 2s. per gallon; how many gallons did he receive?

Ans. 57 gallons.

26. A man exchanged 70 bushels of rye, at 4s. 6d. per bushel, for 40 bushels of wheat, at 7s. per bushel, and received the balance in oats, at 2s. per bushel; how many bushels of oats did he receive?

Ans. 17½.

27. How many bushels of potatoes, at 1s. 6d. per bushel, must be given for 32 bushels of barley, at 2s. 6d. per bushel?

Ans. 53½ bushels.

28. How much salt, at \$1.50 per bushel, must be given in exchange for 15 bushels oats, at 2s. 3d. per bushel?

Note. It will be recollected that, when the price and cost are given, to find the quantity, they must both be reduced to the same denomination before dividing. Ans. 4½ bushels.

29. How much wine, at \$2.75 per gallon, must be given in exchange for 40 yards of cloth, at 7s. 6d. per yard.

Ans. 21 <sup>9</sup>/<sub>11</sub> gallons.

30. A had 41 cwt. of hops, at 30s. per cwt. for which B gave him 20£ in money, and the rest in prunes, at 5d. per pound; how many prunes did A receive?

Ans. 17 cwt. 3qrs. 4 lb.

31. A has linen cloth worth 2s. 6d. per yard; but, in bartering he will have 2s. 9d. per yard; B has broadcloth worth 18s. 9d. per yard, ready money; at what price ought the broadcloth to be rated in bartering with A?

30d. : 35d. :: 225d. : 262½d. Ans. Or,  $\frac{35}{30}$  of 225d. = 1£ 1s. 10½d. Ans. The two operations will be seen to be exactly alike.

32. If cloth, worth 2s. per yard, cash, be rated in barter at 2s. 6d., how should wheat, worth 8s. cash, be rated in exchange for the cloth?

Ans. 10s.

33. If 4 bushels of corn cost \$2, what is it per bushel?

Ans. \$50.

34. If 9 bushels of wheat cost 3£ 7s. 6d. what is that per bushel?

Ans. 7s. 6d.

35. If 40 sheep cost 25£, what is that per head?

36. If 3 bushels of oats cost 7s. 6d. how much are they per bushel?

Ans. 2s. 6d.

37. If 22 yards of broadcloth cost 21£ 9s. what is the price per yard ? Ans. 19s. 6d.

38. At 2s. 6d. per bushel, how much corn can be bought for 10s. ? Ans. 4 bushels.

39. A man having 25£, would lay it out in sheep, at 12s. 6d. apiece, how many can he buy ? Ans. 40.

40. If 20 cows cost 75£, what is the price of 1 cow ?  
— of 2 cows ? — of 5 cows ? — of 15 cows ?

Ans. to the last, 56£ 5s.

41. If 7 men consume 24 lbs. of meat in one week, how much would 1 man consume in the same time ? — 2 men ?  
— 5 men ? — 10 men ? Ans. to the last, 34½ lb.

*Note.* Let the pupil also perform these questions by the rule of proportion.

42. If I pay 1£ 10s. for the use of 25£, how much must I pay for the use of 18£ 15s. ? Ans. 1£. 2s. 6d.

43. What premium must I pay for the insurance of my house against loss by fire, at the rate of  $\frac{1}{2}$  per cent, that is,  $\frac{1}{2}$  pound for 100 pounds, if my house be valued at 2475£ ?  
Ans. 12£ 7s. 6d.

44. What will be the insurance, per annum, of a store and contents, valued at 9876£ 8s. at  $1\frac{1}{2}$  per centum ?

Ans. 148£ 2s. 11d.

45. What commission must I receive for selling 478£ worth of books, at 3 per cent ? Ans. 38£ 4s. 9½d.

46. A merchant bought a quantity of goods for 734£, and sold them so as to gain 21 per cent ; how much did he gain ? and for how much did he sell his goods ?

Ans. to the last, 888£ 2s. 9½d.

47. A merchant bought a quantity of goods at Montreal, for 500£, and paid 43£ for their transportation ; he sold them so as to gain 24 per cent, on the whole cost ; for how much did he sell them ?

Ans. 673£ 6s. 4½d.

48. Bought a quantity of books for 64£, but for cash a discount of 12 per cent. was made ; what did the books cost ?

Ans. 56£ 6s. 4½d.

49. Bought a book, the price of which was marked 1£ 2s. 6d. but for cash the bookseller will sell it at 33½ per cent discount ; what is the cash price ? Ans. 15s.

50. I bought a cask of liquor, containing 120 gallons, for 42£ ; for how much must I sell it to gain 15 per cent ?  
how much per gallon ? Ans. to the last, 4s. ½d.

what is the  
ns. 10s. 6d.  
n be bought  
. 4 bushels.  
n sheep, at  
Ans. 40.  
of 1 cow?  
cows?

ast, 56£ 5s.  
o week, how  
—2 men?  
last, 34£ lb.  
tions by the

y much must  
1£. 2s. 6d.  
rance of my  
cent, that is,  
l at 2475£?

12£ 7s. 6d.  
n, of a store  
ntum?

48£ 2s. 11d.  
selling 478£

38£ 4s. 9d.  
s for 734£,  
much did he

38£ 2s. 9d.  
at Montreal,  
on; he sold

ost; for how  
73£ 6s. 4d.

at for cash a  
id the books  
56£ 6s. 4d.

marked 1£  
t at 33£ per  
Ans. 15s.

0 gallons, for  
5 per cent?

last, 4s. d.

¶ 113.

51. Bought a cask of sugar, containing 740 pounds, for 59£ 4s.; how must I sell it per pound, to gain 25 per cent?

Ans. 2s.

52. What is the interest, at 6 per cent, of 71£ 0s. 4d. for 17 months 12 days?

Ans. 6£ 3s. 6d.

53. What is the interest of 487£ 0s. 3d. for 18 months?

Ans. 43£ 16s. 7d.

54. What is the interest of 88£ 50 for 7 months?

Ans. 8£ 207d.

55. What is the interest of 1000£ for 5 days?

Ans. 16s. 8d.

56. What is the interest of 10s. for 10 years?

Ans. 6s.

57. What is the interest of 884£ 25 for 15 months and 7 days, at 7 per cent?

Ans. 87£ 486+

58. What is the interest of 8154£ 01 for 2 yrs. 4 months and 3 days, at 5 per cent?

Ans. 818£ 032.

59. What sum put to interest at 6 per cent, will, in 2 years and 6 months, amount to 8150?

Note. See ¶ 79.

Ans. 8130£ 434+

60. I owe a man 475£ 10s. to be paid in 16 months without use of money being worth 6 per cent?

Ans. 440£ 5s. 6d.

61. What is the present worth of 1000£ payable in 4 years and 2 months, discounting at the rate of 6 per cent?

62. Bought articles to the amount of 500£, and sold them for 575£, how much was gained?

What per cent was gained? that is, how many pounds were gained on each 100£ laid out? If 500£ gain 75£, what does 100£ gain?

Ans. 15 per cent.

63. Bought cloth at 3£ 10s. per piece, and sold it at 4£ 5s. per piece; how much was gained per centum?

Ans. 21£.

64. A man bought a cask of liquor, containing 126 gallons, for 283£ 10s., and sold it out at the rate of 2£ 15s. per gallon; how much was his whole gain? how much per gallon? how much per cent?

Ans. His whole gain 63£; per gal. 10s. which is 22£ per centum.

65. If 100£ gain 6£ in 12 months, in what time will it gain 4£? — 10£? — 14£?

Ans. to the last, 28 months.

66. In what time will 54£ 10s. at 6 per cent, gain, 2£ 3s. 7½d. Ans. 8 months.

67. 20 men built a certain bridge in 60 days, but, it being carried away in a freshet, it is required how many men can rebuild it in 50 days.

days. days. men.

50 : 60 :: 20 : 24 men, ans.

68. If a field will feed 7 horses 8 weeks, how long will it feed 28 horses? Ans. 2 weeks.

69. If a field, 20 rods in length, must be 8 rods in width to contain an acre, how much in width must be a field, 16 rods in length, to contain the same? Ans. 10 rods.

70. If I purchase for a cloak 12 yards of plaid  $\frac{5}{8}$  of a yd. wide, how much backing  $1\frac{1}{2}$  yards wide must I buy to line it? Ans. 5 yards.

71. If a man earn 18£ 15s. in 5 months, how long must he work to earn 115£? Ans. 30¾ months.

72. B owes C 540£, but B, not being worth so much money, C agrees to take 15s. on a pound; what sum must C receive for the debt? Ans. 405£.

73. A cistern, whose capacity is 400 gallons, is supplied by a pipe which lets in 7 gallons in 5 minutes; but there is a leak in the bottom of the cistern which lets out 2 gallons in 6 minutes, supposing the cistern empty, in what time would it be filled?

In 1 minute  $\frac{7}{5}$  of a gallon is admitted, but in the same time  $\frac{2}{3}$  of a gallon leaks out. Ans. 6 hours, 15 minutes.

74. A ship has a leak which will fill it so as to make it sink in 10 hours; it has also a pump which will clear it in 15 hours; now, if they begin to pump when it begins to leak, in what time will it sink?

In one hour the ship would be  $\frac{1}{10}$  filled by the leak, but in the same time it would be  $\frac{1}{15}$  emptied by the pump.

Ans. 30 hours.

75. A cistern is supplied by a pipe which will fill it in 40 minutes; how many pipes, of the same bigness, will fill it in 5 minutes? Ans. 8.

76. Suppose I lend a friend 500£ for 4 months, he promising to do me a like favour; some time afterward, I have need of 300£; how long may I keep it to balance the former favour? Ans. 6¾ months.

77. Suppose 800 soldiers were in a garrison with pro-

vi  
pa  
be  
val  
7  
mc  
than  
80  
wat  
gall  
No  
27£  
81  
way  
after  
gain  
82.  
ceived  
away  
makes  
the co  
83.  
togethe  
In 1  
hour ha  
on the  
12 spac  
84. 7  
three m  
goes 2  
per hour  
35. T  
two men  
per hour,  
again co  
B gain

gain, 2£  
8 months.  
out, it be-  
many men

men, ans.  
long will  
2 weeks.  
s in width  
a field, 16  
s. 10 rods.  
1  $\frac{1}{2}$  of a yd.  
buy to line  
s. 5 yards.  
long must  
0  $\frac{3}{4}$  months.  
th so much  
at sum must  
Ans. 405£.  
is supplied  
s; but there  
ts out 2 gal-  
ry, in what

he same time  
15 minutes.  
s to make it  
ill clear it in  
it begins to

he leak, but  
e pump.  
s. 30 hours.  
will fill it in  
ness, will fill

Ans. 8.  
ths, he prom-  
ward, I have  
ance the for-  
6  $\frac{3}{4}$  months.  
on with pro-

visions sufficient for 2 months ; how many soldiers must depart, that the provisions may serve them 5 months ?

Ans. 480.

78. If my horse and saddle are worth 21£, and my horse be worth 6 times as much as my saddle, pray what is the value of my horse ?

Ans. 18£.

79. Bought 45 barrels of beef, at 17s. 6d. per barrel, among which are 16 barrels, whereof 4 are worth no more than 3 of the others ; how much must I pay ?

Ans. 35£ 17s. 6d.

80. Bought 126 gallons of rum for 27£ 10s. how much water must be added to reduce the first cost to 3s. 9d. per gallon ?

Note. If 3s. 9d. buy 1 gallon, how many gallons will 27£ 10s. buy ?

Ans. 20  $\frac{3}{4}$  gallons.

81. A thief, having 24 miles start of the officer, holds his way at the rate of 6 miles an hour ; the officer pressing on after him at the rate of 8 miles an hour, how much does he gain in 1 hour ? how long before he will overtake the thief ?

Ans. 12 hours.

82. A hare starts 12 rods before a hound, but is not perceived by him till she has been up 1  $\frac{1}{4}$  minutes ; she scuds away at the rate of 36 rods a minute, and the dog, on view makes after, at the rate of 40 rods a minute ; how long will the course hold ? and what distance will the dog run ?

Ans. 14  $\frac{1}{4}$  minutes, and he will run 570 rods.

83. The hour and minute hands of a watch are exactly together at 12 o'clock ; when are they next together ?

In 1 hour the minute hand passes over 12 spaces, and the hour hand over 1 space ; that is, the minute hand gains upon the hour hand 11 spaces in 1 hour ; and it must gain 12 spaces to coincide with it.

Ans. 1h. 5m. 27  $\frac{3}{4}$ s.

84. There is an island 20 miles in circumference, and three men start together to travel the same way about it ; A goes 2 miles per hour, B 4 miles per hour, and C 6 miles per hour ; in what time will they come together again ?

Ans. 10 hours.

35. There is an island 20 miles in circumference, and two men start together to travel around it ; A travels 2 miles per hour, and B 6 miles per hour ; how long before they will again come together ?

B gains 4 miles per hour, and must gain 20 miles to over-

take A ; A and B will therefore be together once in every 5 hours.

86. In a river, supposing two boats start at the same time from places 300 miles apart ; the one proceeding up stream is retarded by the current 2 miles per hour, while that moving down stream is accelerated the same ; if both be propelled by a steam engine, which would move them 8 miles per hour in still water, how far from each starting place will the boats meet ?

Ans.  $112\frac{1}{2}$  miles from the lower place, and  $187\frac{1}{2}$  miles from the upper place.

87. A man bought a pipe (126 gallons) of wine for 275£ ; he wishes to fill 10 bottles, 4 of which contain 2 quarts, and 6 of them 3 pints each, and to sell the remainder so as to make 30 per cent on the first cost ; at what rate per gallon must he sell it ?

Ans. 5'936£ +.

88. Thomas sold 150 pine apples at 1s. 3d. apiece, and received as much money as Harry received for a certain number of watermelons at 9d. apiece ; how much money did each receive, and how many melons had Harry ?

Ans. 9£ 7s. 6d. and 250 melons.

89. The third part of an army was killed, the fourth part taken prisoners, and 1000 fled ; how many were in this army ?

This and the eighteenth following questions are usually wrought by a rule called *Position*, but they are more easily solved on general principles. Thus,  $\frac{1}{3} + \frac{1}{4} = \frac{7}{12}$  of the army ; therefore, 1000 is  $\frac{5}{12}$  of the whole number of men ; and if  $\frac{5}{12}$  be 1000, how much is 12 twelfths, or the whole ?

Ans. 24000 men.

90. A farmer, being asked how many sheep he had, answered, that he had them in 5 fields ; in the first were  $\frac{1}{4}$  of his flock, in the second  $\frac{1}{4}$ , in the third  $\frac{1}{4}$ , in the fourth  $\frac{1}{4}$ , and in the fifth 450 ; how many had he ?

Ans. 1200.

91. There is a pole,  $\frac{1}{4}$  of which stands in the mud,  $\frac{1}{4}$  in the water, and the rest of it out of the water ; required the part out of the water.

Ans.  $\frac{1}{2}$ .

92. If a pole be  $\frac{1}{4}$  in the mud,  $\frac{3}{8}$  in the water, and 6 feet out of the water, what is the length of the pole ?

Ans. 90 feet.

93. The amount of a certain school is as follows :  $\frac{1}{10}$  of the pupils study grammar,  $\frac{3}{10}$  geography,  $\frac{1}{10}$  arithmetic,  $\frac{2}{10}$  learn to write, and 9 learn to read : what is the number of each ?



in every  
same time  
up stream  
that mov-  
th be pro-  
m 8 miles  
place will

187½ miles

of wine for  
contain 2  
e remainder  
at what rate  
5'936£ +  
apiece,

for a certain  
much money  
larry ?

250 melons.  
ne fourth part  
in this army ?  
s are usually  
e more easily  
= 7½ of the  
ber of men ;  
er the whole ?

24000 men.  
he had, an-  
first were ¼ of  
he fourth 1½,  
Ans. 1200.

the mud, ¼ in  
required the  
Ans. 1½  
er, and 6 feet  
Ans. 90 feet.

follows: 1½ of  
arithmetic, 2½  
he number of

Ans. 5 in grammer, 30 in geography, 24 in arithmetic ;  
12 learn to write, and 9 learn to read.

94. A man, driving his geese to market, was met by  
another, who said, "Good morrow, sir, with your hundred  
geese ;" says he, "I have not a hundred ; but if I had, in  
addition to my present number, one half as many as I now  
have, and 2½ geese more, I should have a hundred : " how  
many had he ?

100 — 2½ is what part of his present number ?

Ans. He had 65 geese.

95. In an orchard of fruit trees, ½ of them bear apples,  
¼ pears, ⅙ plums, 60 of them peaches, and 40, cherries ;  
how many trees does the orchard contain ? Ans. 1200.

96. In a certain village, ½ of the houses are painted white,  
¼ red and, ⅙ yellow, 3 are painted green, and 7 are unpaint-  
ed ; how many houses in the village ? Ans. 120.

97. Seven eights of a certain number exceed four fifths of  
the same number by 6 : required the number.

7⁄8 — 4⁄5 = 3⁄40 ; consequently, 6 is 3⁄40 of the required num-  
ber. Ans. 80.

98. What number is that to which if ¼ of itself be added,  
the sum will be 30 ? Ans. 25.

99. What number is that, to which if its ½ and ¼ be added,  
the sum will be 84 ?

84 = 1 + ½ + ¼ = 7⁄4 times the repuired number. Ans. 48.

100. What number is that, which, being increased by 2⁄3  
and 3⁄5 of itself, and by 22 more, will be made 3 times as  
much ?

The number, being taken 1, 2⁄3, and 3⁄5 times, will make  
2 4⁄15 times and 22 is evidently what that wants of 3 times.

Ans. 30.

101. What number is that, which being increased by 2⁄3,  
3⁄5 and 5⁄6 of itself, the sum will be 234¾ ? Ans. 90.

102. B, C, and D, talking of their ages, C said his age  
was once and a half the age of B, and D said his age was  
twice and one tenth the age of both, and that the sum of  
ther ages was 93 ; what was the age of each ?

Ans. B 12 years, C 18 years, D 63 years old.

103. A schoolmaster being asked how many scholars he  
had, said, " If I had as many more as I now have, 3⁄4 as ma-  
ny, ½ as many, ¼ and ⅙ as many, I should then have 435 ;"  
what was the number of his pupils ? Ans. 120.

104. B and C commenced trade with equal sums of money ; B gained a sum equal to  $\frac{1}{2}$  of his whole stock, and C lost 200£ ; then B's money was double that of C's ; what was the stock of each ?

By the condition of this question, one half of  $\frac{2}{3}$ , that is,  $\frac{1}{3}$  of the stock is equal to  $\frac{1}{3}$  of the stock, less 200£ ; consequently, 200£ is  $\frac{2}{3}$  of the stock.      Ans. 500£.

105. A man was hired 50 days on these conditions,—that for every day he worked, he should receive 3s. 9d., and for every day he was idle, he should forfeit 1s. 3d. ; at the expiration of the time, he received 2£ 17s. 6d. ; how many days did he work, and how many was he idle ?

Had he worked every day, his wages would have been 3s. 9d.  $\times 50 = 9$ £ 7s. 6d. that is 2£ 10s. more than he received ; but every day he was idle lessened his wages 3s. 9d. + 1s. 3d. = 5s. ; consequently he was idle 10 days.

Ans. He wrought 40, and was idle 10 days.

106. B and C have the same income ; B saves  $\frac{1}{2}$  of his ; but C, by spending 30£ per annum more than B, at the end of 8 years finds himself 40£ in debt ; what is their income, and what does each spend per annum ?

Ans. Their income, 200£ per annum ; B spends 175£, and C 205£ per annum.

107. A man, lying at the point of death, left his three sons his property ; to B  $\frac{1}{2}$  wanting 20£, to C  $\frac{1}{4}$ , and to D the remainder, which was 10£ less than the share of B ; what was each one's share ?      Ans. 80£, 50£, and 70£.

108. There is a fish, whose head is 4 feet long ; his tail is as long as his head and half the length of his body, and his body is as long as his head and tail ; what is the length of the fish ?

The pupil will perceive that the length of the body is  $\frac{1}{2}$  the length of the fish.      Ans. 32 feet.

109. B can do a certain piece of work in 4 days, and C can do the same work in 3 days ; in what time would both working together, perform it ?      Ans. 1 $\frac{1}{4}$  days.

110. Three persons can perform a certain piece of work in the following manner: B and C can do it in 4 days, C and D in 6 days, and B and D in 5 days : in what time can they all do it together ?      Ans. 3 $\frac{1}{3}$  days.

111. B and C can do a piece of work in 5 days ; B can do it in 7 days ; in how many days can C do it ?      Ans. 17 $\frac{1}{2}$ .

7 113.

112. A man died, leaving 1000£ to be divided between his two sons, one 14, and the other 18 years of age, in such proportion, that the share of each, being put to interest at 6 per cent, should amount to the same sum when they should arrive at the age of 21; what did each receive?

*Ans.* The elder 546£ 3s. 0 $\frac{1}{2}$ d. +; the younger, 453£ 16s. 11d.

113. A house being let upon a lease of 5 years, at 15£ per annum, and the rent being in arrear for the whole time, what is the sum due at the end of the term, simple interest being allowed at 6 per cent?

*Ans.* 84£.

114. If 3 dozen pair of gloves be equal in value to 40 yards of calico, and 100 yards of calico to three pieces of satin<sup>et</sup> of 30 yards each, and the satin<sup>et</sup> be worth 2s. 6d. per yard, how many pair of gloves can be bought for 20s.?

*Ans.* 8 pair.

115. B C and D would divide 100£ between them, so as that C may have 3£ more than B, and D 4£ more than C; how much must each man have?

*Ans.* B 30£, C 33£, and D 37£.

116. A man has pint bottles, and half pint bottles; how much wine will it take to fill one of each sort? — how much to fill 2 of each sort? — how much to fill 6 of each sort?

117. A man would draw off 30 gallons of wine into 1 pint and 2 pint bottles, of each an equal number; how many bottles will it take, of each kind, to contain the 30 gallons?

*Ans.* 80 of each.

118. A merchant has canisters, some holding 5 pounds, some 7 pounds, and some 12 pounds; how many, of each an equal number, can be filled out of 12 cwt. 3 qrs. 12 lbs. of tea?

*Ans.* 60.

119. If 18 grains of silver make a thimble, and 12 pwts. make a teaspoon, how many of each an equal number, can be made from 15 oz. 6 pwts. of silver?

*Ans.* 24 of each.

120. Let 60 pence be divided among three boys, in such a manner that, as often as the first has 3, the second shall have 5, and the third 7 pence, how many pence will each receive?

*Ans.* 12, 20, and 28 pence.

121. A gentleman, having 50 shillings to pay among his laborers for a day's work, would give to every boy 6d., to every woman 8d. and to every man 16d.; the number of

boys, women and men, was the same ; I demand the number of each. *Ans.* 20.

122. A gentleman had 7£ 17s. 6d. to pay among his laborers ; to every boy he gave 6d., to every woman 8d., and to every man 16d. ; and there were for every boy three women, for every woman two men ; I demand the number of each ? *Ans.* 15 boys, 45 women, and 90 men.

123. A farmer bought a sheep, a cow and a yoke of oxen for 20£ 12s. 6d. ; he gave for the cow 8 times as much as for the sheep, and for the oxen 3 times as much as for the cow ; how much did he give for each ?

*Ans.* For the sheep, 12s. 6d., the cow 5£, and the oxen 15£.

124. There was a farm, of which B owned  $\frac{2}{3}$ , and C  $\frac{1}{3}$  ; the farm was sold for 441£ ; what was each one's share of the money ? *Ans.* B's 126£, and C's 315£.

125. Four men traded together on a capital of 3000£, of which B put in  $\frac{1}{2}$ , C  $\frac{1}{4}$ , D  $\frac{1}{8}$ , and E  $\frac{1}{16}$  ; at the end of 3 years they had gained 2364£ ; what was each one's share of the gain ?

*Ans.* B's 1182£, C's 591£, D's 394£, E's 197£.

126. Three merchants companied ; B furnished  $\frac{2}{3}$  of the capital, C  $\frac{1}{3}$ , and D the rest ; they gain 1250£ ; what part of the capital did D furnish, and what is each one's share of the gain ?

*Ans.* D furnished  $\frac{1}{6}$  of the capital ; and B's share of the gain was 500£, C's 468£ 15s., and D's 281£ 5s.

127. B, C, and D, traded in company ; B put in 125£, C 87£ 10s., and D 120 yards of cloth ; they gained 83£ 2s. 6d., of which D's share was 30£ ; what was the value of D's cloth per yard, and what was B and C's shares of the gain ?

*Note.* D's gain being 30£ is  $\frac{600}{1662\frac{1}{2}} = \frac{1200}{3325} = \frac{48}{133}$  of the whole gain : hence the gain of B and C is readily found ; also the price at which D's cloth was valued per yard.

*Ans.* D's cloth, per yard, 1£, B's share of the gain, 31£ 5s., C's share, 21£ 17s. 6d.

128. Three gardeners, B, C, and D, having bought a piece of ground, find the profits of it amount to 120£ per annum. Now the sum of money which they laid down was in such proportion, that, as often as B paid 5£ C paid 7£,

the num-  
Ans. 20.

ng his la-  
n 8d., and  
three wo-  
number of  
d 90 men.  
ke of ox-  
as much as  
as for the

the oxen

and C  $\frac{1}{2}$ ;  
s share of  
C's 315£.  
3000£, of  
e end of 3  
ne's share

E's 197£.  
ed  $\frac{2}{3}$  of the  
what part  
ne's share

share of the  
s.  
in 125£,  
ained 83£  
the value  
shares of

48  
33 of the  
ly found;  
yard.  
gain, 31£

ght a piece  
£ per an-  
down was  
paid 7£,

and as often as C paid 4£, D paid 6£; I demand how much each man must have per annum of the gain.

Note. By the question, so often as B paid 5£, D paid  $\frac{2}{3}$  of 7£. Ans. B 26£ 13s. 4d., C 37£ 6s. 8d., D 56£.

129. A gentleman divided his fortune among his sons, giving B 9£ as often as C 5£, and D 3£ as often as C 7£; D's dividend was 1537 $\frac{1}{2}$ ; to what did the whole estate amount?

Ans. 11583£ 8s. 10d.

130. B and C undertake a piece of work for 13£ 10s. on which B employed 3 hands 5 days, and C employed 7 hands 3 days; what part of the work was done by B? what part by C? what was each one's share of the money?

Ans. B  $\frac{5}{12}$  and C  $\frac{7}{12}$ ; B's money 5£ 12s. 6d. C's 7£ 17s. 6d.

131. B and C trade in company for one year only; on the 1st of January, B put in 300£, but C could not put any money into the stock until the 1st of April; what did he then put in to have an equal share with B at the end of the year?

Ans. 400£.

132. B, C, D, and E, spent 35s. at a reckoning, and, being a little dipped, agreed that B should pay  $\frac{1}{3}$ , C  $\frac{1}{4}$ , D  $\frac{1}{5}$ , and E  $\frac{1}{6}$ ; what did each pay in this proportion?

Ans. B 13s. 4d., C 10s., D 6s. 8d., and E 5s.

133. There are 3 horses, belonging to 3 men, employed to draw a load of plaister from Montreal to Stanstead, for 6£ 12s. 3d. B and C's horses together are supposed to do  $\frac{2}{3}$  of the work, B and D's  $\frac{9}{10}$ , C and D's  $\frac{13}{10}$ ; they are to be paid proportionally; what is each one's share of the money?

Ans.  $\left\{ \begin{array}{l} \text{B's } 2£ \text{ } 17\text{s. } 6\text{d. } (= \frac{1}{3} \text{ of } 6£ \text{ } 12\text{s. } 3\text{d.}) \\ \text{C's } 1£ \text{ } 8\text{s. } 9\text{d. } (= \frac{2}{3} \text{ of } 6£ \text{ } 12\text{s. } 3\text{d.}) \\ \text{D's } 2£ \text{ } 6\text{s. } 0\text{d. } (= \frac{1}{3} \text{ of } 6£ \text{ } 12\text{s. } 3\text{d.}) \end{array} \right.$

Proof. 6£ 12s. 3d.

134. A person, who was possessed of  $\frac{2}{3}$  of a vessel, sold  $\frac{1}{3}$  of his share for 375£; what was the vessel worth?

Ans. 1500£.

135. A gay fellow soon got the better of  $\frac{2}{3}$  of his fortune; he then gave 1500£ for a commission, and his profusion continued till he had but 450£ left, which he found to be just  $\frac{1}{3}$  of his money, after he had purchased his commission; what was his fortune at first?

Ans. 3780£.

136. A younger brother received 1560£, which was just

$\frac{7}{12}$  of his elder brother's fortune, and  $5\frac{1}{2}$  times the elder brother's fortune was  $\frac{3}{4}$  as much again as the father was worth; what was the value of his estate?

Ans. 19165£ 14s. 3 $\frac{1}{2}$ d.

137. A gentleman left his son a fortune,  $\frac{1}{6}$  of which he spent in three months;  $\frac{1}{4}$  of  $\frac{5}{6}$  of the remainder lasted him 9 months longer, when he had only 537£ left; what was the sum bequeathed him by his father?

Ans. 2082£ 18s. 2 $\frac{1}{2}$ d.

138. A cannon ball, at the first discharge, flies about a mile in 8 seconds; at this rate, how long would a ball be in passing from the earth to the sun, it being 95173000 miles distant?

Ans. 24 years, 46 days, 7 hours, 33 minutes, 20 seconds.

139. A general, disposing his army into a square battalion, found he had 231 over and above, but, increasing each side with one soldier, he wanted 44 to fill up the square; of how many men did his army consist? Ans. 19000.

140. B and C cleared by an adventure at sea, 45 guineas, which was 35£ per cent. upon the money advanced, and with which they agreed to purchase a genteel horse and carriage, whereof they were to have the use in proportion to the sums adventured, which was found to be 11 to B as often as 8 to C; what money did each adventure?

Ans. B 104£ 4s. 2 $\frac{1}{2}$ d., C 75£ 15s. 9 $\frac{1}{2}$ d.

141. Tubes may be made of gold, weighing not more than at the rate of  $\frac{1}{1625}$  of a grain per foot; what would be the weight of such a tube, which would extend across the Atlantic from Quebec to London, estimating the distance at 3000 miles?

Ans. 11b. 8oz. 6pwt. 3 $\frac{2}{3}$  grs.

142. A military officer drew up his soldiers in rank and file, having the number in rank and file equal; on being reinforced with three times his first number of men, he placed them all in the same form, and then the number in rank and file was just double what it was at first; he was again reinforced with three times his whole number of men, and, after placing them all in the same form as at first, his number in rank and file was 40 men each; how many men had he at first?

Ans. 100 men.

143. Supposing a man to stand 80 feet from a steeple, and that a line reaching from the belfry to the man is just 100 feet in length the top of the spire is 3 times as high above

the elder  
rather was

14s. 3<sup>d</sup>.  
which he  
asted him  
what was

8s. 2<sup>d</sup>.  
s about a  
a ball be  
95173000

0 seconds.  
are battal-  
asing each  
he square ;  
ns. 19000.  
45 guin-  
advanced,  
steel horse  
proportion  
11 to B as  
?

15s. 9<sup>d</sup>.  
not more  
hat would  
across the  
distance at  
3<sup>d</sup> grs.  
rank and  
on being  
f men, he  
number in  
; he was  
er of men,  
first, his  
many men  
100 men.  
eeple, and  
just 100  
gh above

the ground as the steeple is ; what is the height of the spire ? and the length of a line reaching from the top of the spire to the man ? See ¶ 103.

Ans. to last, 197 feet nearly.

144. Two ships sail from the same port ; one sails directly east, at the rate of 10 miles an hour, and the other directly south, at the rate of  $7\frac{1}{2}$  miles an hour ; how many miles apart will they be at the end of 1 hour ? — 2 hours ? — 24 hours ? — 3 days ?

Ans. to last, 900 miles.

145. There is a square field, each side of which is 50 rods ; what is the distance between opposite corners ?

Ans.  $70\cdot71 +$  rods.

146. What is the area of a square field, of which the opposite corners are  $70\cdot71$  rods apart ? and what is the length of each side ?

Ans. to last, 50 rods nearly

147. There is an oblong field, 20 rods wide, and the distance of the opposite corners is  $33\frac{1}{2}$  rods ; what is the length of the field ? — its area ?

Ans. Length,  $26\frac{2}{3}$  rods ; area, 3 acres, 1 rood,  $13\frac{1}{2}$  rods.

148. There is a room 18 feet square ; how many yards of carpeting, 1 yard wide, will be required to cover the floor of it ?

$18^2 = 324$  ft. = 36 yards. Ans.


149. If the floor of a square room contain 36 square yards, how many feet does it measure on each side ?

Ans. 18 feet.

When one side of a square is given, how do you find its area, or superficial contents ?

When the area, or superficial contents, of a square is given, how do you find one side ?

150. If an oblong piece of ground be 80 rods long and 20 rods wide, what is its area ?



Note. A Parallelogram or Oblong, has its opposite sides equal and parallel, but the adjacent sides unequal. Thus, A B C D is a parallelogram, and also E F C D, and it is easy to see, that the contents of both are equal. Ans. 1600 rods, = 10 acres.

151. What is the length of an oblong, or parallelogram, whose area is 10 acres, and whose breadth is 20 rods ?

Ans. 80 rods.



152. If the area be 10 acres, and the length 80 rods, what is the other side?

When the *length* and *breadth* are given, how do you find the *area* of an oblong, or parallelogram?

When the *area* and *one side* are given, how do you find the *other side*?

153. If a board be 18 inches wide at one end, and 10 inches wide at the other, what is the *mean* or *average* width of the board? Ans. 14 inches.

When the *greatest* and *least* width are given, how do you find the *mean* width?

154. How many square feet in a board 16 feet long, 1'8 feet wide at one end, and 1'3 at the other?

Mean width,  $\frac{1'8 + 1'3}{2} = 1'55$ ; and  $1'55 \times 16 = 24'8$  feet, Ans.

155. What is the number of square feet in a board 20 feet long, 2 feet wide at one end, and running to a point at the other? Ans. 20 feet.

How do you find the contents of a straight edged board, when one end is wider than the other?

If the length be in *feet*, and the breadth in feet, in what denomination will the product be?

If the length be *feet*, and the breadth *inches*, what *parts* of a *foot* will be the product?

156. There is an oblong field, 40 rods long and 20 rods wide; if a straight line be drawn from one corner to the opposite corner, it will be divided into two equal *right-angled triangles*; what is the area of each?

Ans. 400 square rods = 2 acres 2 roods.

157. What is the area of a triangle, of which the *base* is 30 rods, and the *perpendicular* 10 rods? Ans. 150 rods.

158. If the area be 150 rods, and the base 30 rods, what is the *perpendicular*? Ans. 10 rods.

159. If the *perpendicular* be 10 rods, and the area 150 rods what is the *base*? Ans. 30 rods.

When the *legs* (the *base* and *perpendicular*) of a right-angled triangle are given, how do you find its *area*?

When the *area* and *one* of the *legs* are given, how do you find the *other leg*?

*Note.* Any triangle may be divided into two *right-angled triangles*, by drawing a perpendicular from one corner to the opposite side, as may be seen by the annexed figure.

C



Here,  $\triangle ABC$  is a triangle, divided into two right-angled triangles,  $\triangle ADC$ , and  $\triangle BDC$ ; therefore, the whole base,  $AB$ , multiplied by *one half* the perpendicular  $dC$ , will give the area of the whole. If  $AB=60$  feet, and  $dC=16$  feet, what is the area? Ans. 480 feet.

160. There is a triangle, each side of which is 10 feet; what is the length of a perpendicular from one angle to its opposite side? and what is the area of the triangle?

*Note.* It is plain, the perpendicular will divide the opposite side into two equal parts.

Ans. Perpendicular,  $8.66+$  feet; area,  $43.3+$  feet.

161. What is the solid contents of a cube measuring 6 feet on each side? Ans. 216 feet.

When *one side* of a cube is given, how do you find its *solid contents*?

When the *solid contents* of a cube are given, how do you find *one side* of it?

162. How many cubic inches in a brick which is 8 inches long, 4 inches wide, and 2 inches thick? — in 2 bricks? — in 10 bricks? Ans. to last, 640 cubic inches.

163. How many bricks in a cubic foot? — in 40 cubic feet? — in 1000 cubic feet Ans. to last, 27000.

164. How many bricks will it take to build a wall 40 feet in length, 12 feet high, and 2 feet thick? Ans. 25920.

165. If a wall be 150 bricks, = 100 feet, in length, and 4 bricks, = 16 inches, in thickness, how many bricks will lay one course? — 2 courses? — 10 courses? If the wall be 48 courses, = 8 feet high, how many bricks will build it?  $150 \times 4 = 600$ , and  $600 \times 48 = 28800$ , Ans.

166. The river Po is 1000 feet broad, and 10 feet deep, and it runs at the rate of 4 miles an hour; in what time will it discharge a cubic mile of water (reckoning 5000 feet to the mile) into the sea? Ans. 26 days, 1 hour.

167. If the country, which supplies the river Po with water, be 380 miles long, and 120 broad, and the whole land upon the surface of the earth be 62,700,000 square miles, and if the quantity of water discharged by the rivers into the sea be every where proportional to the extent of land by which the rivers are supplied; how many times

greater than the Po will the whole amount of the rivers be?   
 Ans. 1375 times.

168. Upon the same supposition, what quantity of water, altogether, will be discharged by all the rivers into the sea in a year, or 365 days.   
 Ans. 19272 cubic miles.

169. If the proportion of the sea on the surface of the earth to that of land be as  $10\frac{1}{2}$  to 5, and the mean depth of the sea be a quarter of a mile; how many years would it take, if the ocean were empty, to fill it by the rivers running at the present rate?

Ans. 1708 years, 17 days, 12 hours.

170. If a cubic foot of water weigh 1000 oz. avoirdupois, and the weight of mercury be  $13\frac{1}{2}$  times greater than of water, and the height of the mercury in the barometer (the weight of which is equal to the weight of a column of air on the same base, extending to the top of the atmosphere) be 30 inches; what will be the weight of the air upon a square foot? — a square mile? and what will be the whole weight of the atmosphere, supposing the size of the earth as in questions 166 and 168?

Ans. 2109'375 lbs. weight on a square foot.

52734375000 " " mile.

10249980468750000000 " " of the whole atmosphere.

171. If a circle be 14 feet in diameter, what is its circumference?

*Note.* It is found by calculation, that the *circumference* of a circle measures about  $3\frac{1}{2}$  times as much as its *diameter*, or, more accurately, in decimals,  $3\cdot14159$  times.   
 Ans. 44 feet.

172. If a wheel measure 4 feet across from side to side, how many feet around it?   
 Ans. 124.

173. If the diameter of a circular pond be 147 feet, what is its circumference?   
 Ans. 462 feet.

174. What is the diameter of a circle, whose circumference is 462 feet?   
 Ans. 147 feet.

175. If the distance through the centre of the earth, from side to side, be 7911 miles, how many miles around it?

$7911 \times 3\cdot14159 = 24853$  square miles, nearly,   
 Ans.

176. What is the area or contents of a circle, whose diameter is 7 feet, and its circumference 22 feet?

*Note.* The area of a circle may be found by multiplying  $\frac{1}{2}$  the diameter into  $\frac{1}{2}$  the circumference.   
 Ans.  $38\frac{1}{2}$  sq. feet.

177. What is the area of a circle, whose circumference is 176 rods?   
 Ans. 2464 rods.

178. If a circle is drawn within a square, containing 1 square rod, what is the area of this circle ?

*Note.* The diameter of the circle being 1 rod, the circumference will be  $3\cdot14159$ , Ans.  $\cdot7854$  of a sq. rod nearly.

Hence, if we square the diameter of any circle, and multiply the square by  $\cdot7854$ , the product will be the area of the circle.

179. What is the area of a circle whose diameter is 10 rods ?  $10^2 \times \cdot7854 = 78\cdot54$ . Ans.  $78\cdot54$  rods.

180. How many square inches of leather will cover a ball  $3\frac{1}{2}$  inches in diameter ?

*Note.* The area of a globe or ball is 4 times as much as the area of a circle of the same diameter, and may be found, therefore, by multiplying the whole circumference into the whole diameter.

Ans.  $38\frac{1}{2}$  square inches.

181. What is the number of square miles on the surface of the earth, supposing its diameter 7911 miles ?

$7911 \times 24853 = 196,612,083$ , Ans.

182. How many solid inches in a ball 7 inches in diameter ?

*Note.* The solid contents of a globe are found by multiplying its area by  $\frac{1}{3}$  part of its diameter.

Ans.  $179\frac{2}{3}$  solid inches.

183. What is the number of cubic miles in the earth, supposing its diameter as above ?

Ans.  $259,233,031,435$  miles.

184. What is the capacity, in cubic inches, of a hollow globe 20 inches in diameter, and how much wine will it contain, 1 gallon being 231 cubic inches ?

Ans.  $4188\cdot8+$  cubic inches, and  $18\cdot13+$  gallons.

185. There is a round log, all the way of a bigness ; the areas of the circular ends of it are each 3 square feet ; how many solid feet does 1 foot in length of this log contain ? — 2 feet in length ? — 3 feet ? — 10 feet ? A solid of this form is called a *Cylinder*.

How do you find the solid content of a cylinder, when the area of one end, and the length are given ?

186. What is the solid content of a round stick, 20 feet long and 7 inches through, that is, the ends being 7 inches in diameter ?

Find the area of one end, as before taught, and multiply it by the length.

Ans.  $5\cdot347+$  cubic feet.

If you multiply square inches by *inches in length*, what parts of a *foot* will the product be? — if square inches by *feet in length*, what part?

187. A Winchester bushel is 18½ inches in diameter, and 8 inches deep; how many cubic inches does it contain?

Ans. 2150¼.

It is plain, from the above, that the solid content of all bodies, which are of uniform bigness throughout, whatever may be the form of the ends, is found by *multiplying the area of one end into its height or length*.

Solids which decrease gradually from the base till they come to a point, are generally called *Pyramids*. If the base be a square, it is called a *square pyramid*; if a triangle, a *triangular pyramid*; if a circle, a *circular pyramid* or a *cone*. The point at the top of a pyramid is called the *vertex*, and a line, drawn from the *vertex* perpendicular to the *base*, is called the *perpendicular height* of the pyramid.

The *solid content* of any pyramid may be found by multiplying the area of the *base* by ⅓ of the *perpendicular height*.

188. What is the solid content of a pyramid whose base is 4 feet square, and the perpendicular height 9 feet?

$$4^2 \times \frac{9}{3} = 48.$$

Ans 48 feet.

189. There is a *cone*, whose height is 27 feet, and whose base is 7 feet in diameter; what is its content?

Ans. 346½ feet.

190. There is a cask, whose head diameter is 25 inches, bung diameter 31 inches, and whose length is 36 inches; how many wine gallons does it contain? — how many beer gallons;

*Note.* The *mean diameter* of the cask may be found by adding 2 thirds, or, if the staves be but little curving, 6 tenths, of the difference between the head and bung diameters, to the head diameter. The cask will then be reduced to a cylinder.

Now, if the square of the mean diameter be multiplied by 7854, (ex. 177,) the product will be the area of one end, and that, multiplied by the length, in inches, will give the solid content, in cubic inches, (ex. 185,) which, divided by 231, (note to table, wine meas.) will give the content in wine gallons, and, divided by 282, (note to table, beer meas.) the content in ale or beer measure.

In this process we see, that the square of the mean diam-

eter  
lons  
only  
by  
for  
lons,  
H  
Mul  
tity  
ing  
and  
In  
the h  
25 in

Then

191  
amete  
45 in  
19%  
prop,  
pound  
anced  
the pr

*Note.*  
feet fr  
while  
2 inch  
that th  
when  
Hence,  
will ba  
we div  
distanc  
ways  
is the  
168.

193.  
require  
194.  
power t

eter will be multiplied by '7854, and divided, for wine gallons, by 231. Hence we may contract the operation by only multiplying by their quotient, ( $\frac{7854}{231} = .0034$ ;) that is by '0034, (or by 34, pointing off 4 figures from the product for decimals.) For the same reason we may, for beer gallons, multiply by ( $\frac{7854}{282} = .0028$ , nearly.) '0028, &c.

Hence this concise RULE, for gauging or measuring casks, *Multiply the square of the mean diameter by the length; multiply this product by 34, for wine, or by 28 for beer, and pointing off four decimals, the product will be the content in gallons and decimals of a gallon.*

In the above example, the bung diameter, 31 in.—25 in. the head diameter = 6 in. difference, and  $\frac{2}{3}$  of 6 = 4 inches; 25 in. + 4 in. = 29 in. mean diameter.

Then,  $29^2 = 841$ , and  $841 \times 36 \text{ in.} = 30276$ .

Then,  $\begin{cases} 30276 \times 34 = 1029384. & \text{Ans. } 102.9384 \text{ wine gals.} \\ 30726 \times 28 = 847728. & \text{Ans. } 84.7728 \text{ beer gals.} \end{cases}$

191. How many wine gallons in a cask whose bung diameter is 36 inches, head diameter 27 inches, and length 45 inches? Ans. 166.617.

192. There is a lever 10 feet long, and the fulcrum, or prop, on which it turns is 2 feet from one end; how many pounds weight at the end, 2 feet from the prop, will be balanced by a power of 42 pounds at the other end, 8 feet from the prop?

*Note.* In turning around the prop, the end of the lever 8 feet from the prop will evidently pass over a space of 8 inches, while the end 2 feet from the prop passes over a space of 2 inches. Now it is a fundamental principle in mechanics, that the weight and power will exactly balance each other, when they are inversely as the spaces they pass over. Hence, in this example, 2 pounds, 8 feet from the prop, will balance 8 pounds 2 feet from the prop; therefore, if we divide the distance of the power from the prop by the distance of the weight from the prop, the quotient will always express the ratio of the weight to the power;  $\frac{8}{2} = 4$ , that is the weight will be 4 times as much as the power,  $42 \times 4 = 168$ . Ans. 168 lbs.

193. Supposing the lever as above, what power would it require to raise 1000 pounds? Ans.  $\frac{1000}{4} = 250$  lbs.

194. If the weight to be raised be 5 times as much as the power to be applied, and the distance of the weight from the

prop be 4 feet, how far from the prop must the power be applied ?

Ans. 20 feet.

195. If the greater distance be 40 feet, and the less  $\frac{1}{2}$  of a foot, and the power 175 pounds, what is the weight ?

Ans. 14000 pounds.

196. Two men carry a kettle, weighing 200 pounds ; the kettle is suspended on a pole, the bale being 2 feet 6 inches from the hands of one, and 3 feet 4 inches from the hands of the other ; how many pounds does each bear.

Ans. 114 $\frac{1}{2}$  lbs. and 85 $\frac{1}{2}$  lbs.

197. There is a windlass, the wheel of which is 60 inches in diameter, and the axis, around which the rope coils, is 6 inches in diameter ; how many pounds on the axle will be balanced by 240 pounds at the wheel ?

*Note.* The spaces passed over are evidently as the diameters, or the circumferences ; therefore,  $\frac{60}{6} = 10$ , ratio.

Ans. 2400 pounds.

198. If the diameter of the wheel be 60 inches, what must be the diameter of the axle, that the ratio of the weight to the power may be 10 to 1 ?

Ans. 6 inches.

*Note.* This calculation is on the supposition, that there is no friction, for which it is usual to add  $\frac{1}{3}$  to the power which is to work the machine.

199. There is a screw, whose threads are 1 inch asunder, which is turned by a lever 5 feet, = 60 inches, long ; what is the ratio of the weight to the power ?

*Note.* The power applied at the end of the lever will describe the circumference of a circle  $60 \times 2 = 120$  inches in diameter, while the weight is raised 1 inch ; therefore, the ratio will be found by dividing the circumference of a circle, whose diameter is twice the length of the lever, by the distance between the threads of the screw.  $120 \times 3\frac{1}{2} = 377\frac{1}{2}$  circumference, and  $\frac{377\frac{1}{2}}{1} = 377\frac{1}{2}$ , ratio, Answer.

200. There is a screw, whose threads are  $\frac{1}{2}$  of an inch asunder ; if it be turned by a lever 10 feet long, what weight will be balanced by 120 pounds power ?

Ans. 30171 lbs.

201. There is a machine, in which the power moves over 10 feet, while the weight is raised 1 inch ; what is the power of that machine, that is, what is the ratio of the weight to the power ?

Ans. 120.

202. A rough stone was put into a vessel, whose capacity



was 14 wine quarts, which was afterwards filled with  $2\frac{1}{2}$  qts. of water; what was the cubic content of the stone?

Ans.  $664\frac{1}{4}$  inches.

## FORMS OF NOTES, RECEIPTS, ORDERS AND BILLS OF PARCELS.

### NOTES.

#### No. I.

Montreal, Oct. 22, 1833.

For value received, I promise to pay to Oliver Bountiful, or order, two pounds, ten shillings and sixpence, on demand, with interest.

WILLIAM TRUSTY.

Attest, TIMOTHY TESTIMONY.

#### No. II.

Kingston, October 10, 1833.

For value received of A B, in goods, wares and merchandize, this day sold and delivered, I promise to pay him, or bearer, — pounds, — shillings and — pence, in ten days from date, with interest.

C — D —.

#### No. III.

By two persons.

Stanstead, October 1, 1833.

For value received, of —, in — this day sold and delivered, we jointly and severally promise to pay him, or order, — pounds, — shillings and — pence in — days from date, with interest.

B — C —,

D — E —.

### RECEIPTS.

Montreal, October, 10, 1833.

Received from Mr. Durance Adley ten pounds, in full of all accounts.

ORVAND CONSTANCY.

Receipt for money received on a Note.

York, November 1, 1833.

Received of Mr. Simon Eastly (by the hand of Titus Trusty,) sixteen pounds ten shillings and six pence, which is endorsed on his note of June 3, 1831.

X 2

SAMSON SNOW.

*A Receipt for money received on Account.*

Stanstead, June 2, 1833.

Received of Thomas Dubois twenty pounds, on account.  
 ORLANDO PROMPT.

*Receipt for money received for another person.*

Sherbrooke, June 4, 1833.

Received from P. D. twenty-five pounds for account of  
 J. T. ELI TRUMAN.

*Receipt for interest due on a Note.*

Quebec, December 12, 1833.

Received of I. S. fifteen pounds, in full of one year's interest of 250£, due to me on the — day of — last, on note from the said I. S.  
 SOLOMON GRAY.

*Receipt for Money paid before it becomes due.*

Proscot, May 3, 1833.

Received of T. Z. fifteen pounds, advanced in full for one year's rent of my farm, leased to the said T. Z. ending the first day of April next, 1834.  
 JOHN HONORUS.

**ORDERS.**

Belville, November 3, 1833.

Mr. Stephen Girard. For value received, pay to A B, or order, five pounds and six shillings, and place the same to my account.  
 SAUL MANN.

Montreal, September 1, 1830.

Mr. TIMOTHY TITUS. Please to deliver to Mr. L. D. such goods as he may call for, to the amount of seven pounds, and place the same to the account of your obedient servant,  
 NICANOR LINUS.

**BILLS OF PARCELS.**

It is usual when goods are sold for the seller to deliver to the buyer, with the goods, a bill of the articles, and their

price  
called12½ y  
8

3 hogs  
2 pipes  
1 hogsh  
3 casks  
5 bags  
1 chest

5682 feet  
2000 "  
800 "  
1500 "  
650 "  
879 "  
236 "

prices, with the amount cast up. Such bills are sometimes called bills of parcels.

Montreal, 6th May, 1833.

Mr. ABEL ATLAS.

Bought of BENJ. BUCK.

	£.	s.	d.
12½ yds. figured Satin, at 12s. 6d. per yd.	7	16	3
8 " Sprigged Tabby, - 6s. 3d. -	2	10	0

Received payment, £10 · 6 · 3

BENJ. BUCK.

Montreal, 14th May, 1833.

Mr. JOHN BURTON,

Bought of GEORGE WILLIAMS,

3 hogsheads new Rum, 118 gal. each, at 1s. 6d. per gal.			
2 pipes French Brandy, 126 & 132 gal. " 5s. 7d. "			
1 hogshead brown Sugar, 9½ cwt. " 2£ 11s. 9d. cwt.			
3 casks of Rice, 269lb. each, " 3d. lb.			
5 bags Coffee, 75lb. each, " 1s. 2d. "			
1 chest hyson Tea, 86lb. " 4s. 8d. "			

Received payment, 111£ 6s. 7½d.

For GEORGE WILLIAMS,

THOMAS ROUSSEAU.

Wilderness, 8th Feb. 1833.

Mr. SIMON JOHNSON,

Bought of ASA FULLUM,

5682 feet Boards, at 1£ 10s. per M.			
2000 " " " 2£ 1s. 8d. "			
800 " Stuff, " 3 · 3 · 2 · "			
1500 " Lathing, " 1 · "			
650 " Plank, " 1 · 10 · "			
879 " Timber, " 12 · 6 · "			
236 " " " 13 · 9 · "			

18£ 8s. ¾d.

ver to  
their

# BOOK-KEEPING.

It is necessary that every man should have some regular uniform method of keeping his accounts. What this method shall be, the law does not prescribe; but in cases of dispute it requires that the several items of the account be proved to have been sold and delivered.

For farmers and mechanics, the following method will be found both convenient and easy. It consists in having one single book, entering the name of the person, with whom an account is to be opened, at the top of the *left* hand page, *Dr.*; and at the top of the *right* hand page, his place of residence, and *Cr.* as follows:

<b>Dr. JAMES MACKAY, LENOXVILLE, Cr.</b>							
		1833.					
Jan. 4.	To 5 cords of Wood, at 8s. 9d. - -	£	2	9	By 1 Plough, - - -	£	2
May 5.	To 1 day's work, self and oxen, - -		3	6	By repairing Cart Wheels, - -		6
July 20.	To 4 bushels of Rye, at 3s. 9d. deliv- ered by your order to C. D. - -		15	6	By Cash to balance, - - -		12
			3	3			6

Another method, which some prefer is, to place the *Dr.* and *Cr.* on the same page, as follows:

<b>JOHN POPE, STANSTEAD, Cr.</b>							
1833.		<b>Dr.</b>		<b>Cr.</b>			
Jan. 24.	To 2½ tons of Hay, at 2£ per ton, - - -	£	5	10	£	1	15
Feb. 29.	By 14 bushels of Corn, at 2s. 6d. - - -						
Feb. 2.	To 30 lbs. of Flax, d'd. by your order to A B, at 7½d. - - -			18		4	13
Feb. 6.	By Cash, to balance, - - -			8		6	8
							9

Jan. 24.	To 2½ tons of Hay, at 2 <sup>s</sup> ℓ per ton,	-	-	-	-
29.	By 14 bushels of Corn, at 2s. 6d.	-	-	-	-
Feb. 2.	To 30 lbs. of Flax, d'd by your order to A B, at 7jd.	-	-	-	-
6.	By Cash, to balance,	-	-	-	-
		6	8	9	9
		5	10	1	15

